

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSPTARHH1626

PASSWORD:
TERMINAL (ENTER 1, 2, 3, OR ?):2

***** Welcome to STN International *****

NEWS 1 Web Page for STN Seminar Schedule - N. America
NEWS 2 JAN 08 CHEMLIST enhanced with New Zealand Inventory of Chemicals
NEWS 3 JAN 16 CA/Caplus Company Name Thesaurus enhanced and reloaded
NEWS 4 JAN 16 IPC version 2007.01 thesaurus available on STN
NEWS 5 JAN 16 WPIDS/WPINDEX/WPIX enhanced with IPC 8 reclassification data
NEWS 6 JAN 22 CA/Caplus updated with revised CAS roles
NEWS 7 JAN 22 CA/Caplus enhanced with patent applications from India
NEWS 8 JAN 29 PIAR reloaded with new search and display fields
NEWS 9 JAN 29 CAS Registry Number crossover limit increased to 300,000 in multiple databases
NEWS 10 FEB 15 PATDPASC enhanced with Drug Approval numbers
NEWS 11 FEB 15 RUSSAPAT enhanced with pre-1994 records
NEWS 12 FEB 23 KORAPAT enhanced with IPC 8 features and functionality
NEWS 13 FEB 26 MEDLINE reloaded with enhancements
NEWS 14 FEB 26 EMBASE enhanced with Clinical Trial Number field
NEWS 15 FEB 26 TOXCENTER enhanced with reloaded MEDLINE
NEWS 16 FEB 26 IFICDB/IFIPAT/IFIUDB reloaded with enhancements
NEWS 17 FEB 26 CAS Registry Number crossover limit increased from 10,000 to 300,000 in multiple databases
NEWS 18 MAR 15 WPIDS/WPIX enhanced with new FRAGHITSTR display format
NEWS 19 MAR 16 CASREACT coverage extended
NEWS 20 MAR 20 MARPAT now updated daily
NEWS 21 MAR 22 LWPI reloaded
NEWS 22 MAR 30 RDISCLOSURE reloaded with enhancements
NEWS 23 APR 02 JICST-EPLUS removed from database clusters and STN
NEWS 24 APR 30 GRNBANK reloaded and enhanced with Genome Project ID field
NEWS 25 APR 30 CHEMCATS enhanced with 1.2 million new records
NEWS 26 APR 30 CA/Caplus enhanced with 1870-1889 U.S. patent records
NEWS 27 APR 30 INPADOC replaced by INPADOCDB on STN
NEWS 28 MAY 01 New CAS web site launched
NEWS 29 MAY 08 CA/Caplus Indian patent publication number format defined
NEWS 30 MAY 14 RDISCLOSURE on STN Easy enhanced with new search and display fields
NEWS 31 MAY 21 BIOSIS reloaded and enhanced with archival data
NEWS 32 MAY 21 TOXCENTER enhanced with BIOSIS reload
NEWS 33 MAY 21 CA/Caplus enhanced with additional kind codes for German patents
NEWS 34 MAY 22 CA/Caplus enhanced with IPC reclassification in Japanese patents

NEWS EXPRESS NOVEMBER 10 CURRENT WINDOWS VERSION IS V8.01c, CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0c(JP), AND CURRENT DISCOVER FILE IS DATED 25 SEPTEMBER 2006.

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS LOGIN Welcome Banner and News Items

NEWS IPCS For general information regarding STN implementation of IPC 8

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

***** STN Columbus *****

FILE 'HOME' ENTERED AT 08:18:24 ON 30 MAY 2007

file reg	SINCE FILE	TOTAL
COST IN U.S. DOLLARS	ENTRY	SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'REGISTRY' ENTERED AT 08:18:37 ON 30 MAY 2007

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2007 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 28 MAY 2007 HIGHEST RN 935999-19-2
DICTIONARY FILE UPDATES: 28 MAY 2007 HIGHEST RN 935999-19-2

New CAS Information Use Policies, enter HELP USAGETERMS for details.


TSCA INFORMATION NOW CURRENT THROUGH December 2, 2006

Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stndoc/properties.html>

Uploading C:\Program Files\Stnexp\Queries\10.561754\clm14.str



chain nodes :
1 2 3 4 5 6 7 8 9 10 19 20 21 22 23 24 25 26 27 28 29 30 36
37 38
ring nodes :
11 12 13 14 15 31 32 33 34 35
chain bonds :
1-2 1-9 1-10 2-3 3-4 4-5 4-8 5-6 5-7 11-19 19-20 19-21 22-23 22-30
22-32 23-24 24-25 25-26 25-29 26-27 26-28 31-36 36-37 36-38
ring bonds :
11-12 11-15 12-13 13-14 14-15 31-32 31-35 32-33 33-34 34-35
exact/norm bonds :
1-2 1-9 1-10 2-3 3-4 4-5 4-8 5-6 5-7 11-12 11-15 11-19 12-13 13-14
14-15 19-20 19-21 22-23 22-30 22-32 23-24 24-25 25-26 25-29 26-27 26-28 31-32 31-35 31-36
32-33 33-34 34-35 36-37 36-38

G1:C,8

Match level :

1:CLASS 2:CLASS 3:CLASS 4:CLASS 5:CLASS 6:CLASS 7:CLASS 8:CLASS 9:CLASS
10:CLASS 11:Atom 12:Atom 13:Atom 14:Atom 15:Atom 19:CLASS 20:CLASS 21:CLASS 22:CLASS
23:CLASS
24:CLASS 25:CLASS 26:CLASS 27:CLASS 28:CLASS 29:CLASS 30:CLASS 31:Atom 32:Atom 33:Atom
34:Atom 35:Atom 36:CLASS 37:CLASS 38:CLASS
fragments assigned product role:
containing 22
fragments assigned reactant/reagent role:
containing 1
containing 11

L1 STRUCTURE UPLOADED

-- d
L1 HAS NO ANSWERS
L1 STR

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

Structure attributes must be viewed using STN Express query preparation.

file casreact	SINCE FILE	TOTAL
COST IN U.S. DOLLARS	ENTRY	SESSION
FULL ESTIMATED COST	0.45	0.66

FILE 'CASREACT' ENTERED AT 08:19:13 ON 30 MAY 2007
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications.

FILE CONTENT:1840 - 27 May 2007 VOL 146 ISS 23

New CAS Information Use Policies, enter HELP USAGETERMS for details.

* CASREACT now has more than 12 million reactions *

Some CASREACT records are derived from the ZIC/VINITI database (1974-1999) provided by InfoChem, INPI data prior to 1986, and Biotransformations database compiled under the direction of Professor Dr. Klaus Kieslich.

This file contains CAS Registry Numbers for easy and accurate substance identification.

-- s 11 sss sam
SAMPLE SEARCH INITIATED 08:19:21 FILE 'CASREACT'
SCREENING COMPLETE - 0 REACTIONS TO VERIFY FROM 0 DOCUMENTS
100.0% DONE 0 VERIFIED 0 HIT RXNS 0 DOCS
SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
BATCH **COMPLETE**
PROJECTED VERIFICATIONS: 0 TO 0
PROJECTED ANSWERS: 0 TO 0

L2 0 SEA SSS SAM L1 (0 REACTIONS)

-- s 11 sss full
FULL SEARCH INITIATED 08:19:30 FILE 'CASREACT'
SCREENING COMPLETE - 14 REACTIONS TO VERIFY FROM 4 DOCUMENTS
100.0% DONE 14 VERIFIED 2 HIT RXNS 1 DOCS
SEARCH TIME: 00.00.01

L3 1 SEA SSS FUL L1 (2 REACTIONS)

-- d ibib abs hitstr
'HITSTR' IS NOT A VALID FORMAT FOR FILE 'CASREACT'

The following are valid formats:

ABS ----- GI and AB
ALL ----- BIB, AB, IND, RE, Single-step Reactions

10/561,754 5/447 Robert Havlin

APPS ----- AI, PRAI
BIB ----- AN, plus Bibliographic Data
CAN ----- List of CA abstract numbers without answer numbers
CBIB ----- AN, plus Compressed Bibliographic Data
DALL ----- ALL, delimited (end of each field identified)
IABS ----- ABS, indented with text labels
IALL ----- ALL, indented with text labels
IBIB ----- BIB, indented with text labels
IND ----- Indexing data
IPC ----- International Patent Classifications
ISTD ----- STD, indented with text labels
OBIB ----- AN, plus Bibliographic Data (original)
OIBIB ----- OBIB, indented with text labels

OBIB ----- BIB, no citations
OIBIB ----- IBIB, no citations

MAX ----- Same as ALL
PATS ----- PI, SO
SCAN ----- TI and FCRD (random display, no answer number. SCAN must be entered on the same line as DISPLAY, e.g., D SCAN.)

SSRX ----- Single-Step Reactions (Map, Diagram, and Summary for all single-step reactions)
STD ----- BIB, IPC, and NCL

CRD ----- Compact Display of All Hit Reactions
CRDRF ----- Compact Reaction Display and SO, PY for Reference
FHIT ----- Reaction Map, Diagram, and Summary for first hit reaction
FHITCBIB ----- FHIT, AN plus CBIB
FCRD ----- First hit in Compact Reaction Display (CRD) format
FCRDRF ----- First hit in Compact Reaction Display (CRD) format with CA reference information (SO, PY). (Default)
PPATH ----- PATH, plus Reaction Summary for the "long path"
SPATH ----- SPATH, plus Reaction Summary for the "short path"
HIT ----- Reaction Map, Reaction Diagram, and Reaction Summary for all hit reactions and fields containing hit terms
OCC ----- All hit fields and the number of occurrences of the hit terms in each field. Includes total number of HIT, PATH, SPATH reactions. Labels reactions that have incomplete verifications.
PATH ----- Reaction Map and Reaction Diagram for the "long path". Displays all hit reactions, except those whose steps are totally included within another hit reaction which is displayed
RX ----- Hit Reactions (Map, Diagram, Summary for all hit reactions)
RXG ----- Hit Reaction Graphics (Map and Diagram for all hit reactions)
RXL ----- Hit Reaction Long (Map, Diagram, Summary for all hit reactions)
RXS ----- Hit Reaction Summaries (Map and Summary for all hit reactions)
SPATH ----- Reaction Map and Reaction Diagram for the "short path". Displays all single step reactions which contain a hit substance. Also displays those multistep reactions that have a hit substance in both the first and last steps of the reaction, except for those hit reactions whose steps are totally included within another hit reaction which is displayed

To display a particular field or fields, enter the display field codes. For a list of the display field codes, enter HELP DFIELD at an arrow prompt (=>). Examples of combinations include: D TI; D BIB RX; D TI, AU, FCRD. The information is displayed in the same order

10/561,754 6/447 Robert Havlin

as the specification. All of the formats, except CRD, CRDRF, FHIT, PATH, PPATH, SPATH, FSPATH, FCRD, FCRDRF, HIT, RX, RXG, RXS, SCAN, and OCC, may be used with the DISPLAY command to display the record for a specified Accession Number.


ENTER DISPLAY FORMAT (FCRDRF):end

=> d ibib abs hit

L3 ANSWER 1 OF 1 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 142:113909 CASREACT Full-text
TITLE: Process for preparation of N-[oxidopyridinyl] 1-pyrrolidine derivatives
INVENTOR(S): Slade, Joel; Vivalo, James Anthony; Chen, Guang-Pei; Bajwa, Joginder Singh; Parker, David John
PATENT ASSIGNEE(S): Novartis A.-G., Switz.; Novartis Pharma G.m.b.H.
SOURCE: PCT Int. Appl., 34 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005000835	A1	20050106	WO 2004-EP6915	20040625
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SI, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TO			
AU 2004251876	A1	20050106	AU 2004-251876	20040625
CA 2530142	A1	20050106	CA 2004-2530142	20040625
EP 1641778	A1	20060405	EP 2004-740324	20040625
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO, CY, TR, BG, CZ, SE, HU, PL, SK			
BR 2004011921	A	20060815	BR 2004-11921	20040625
CN 1829710	A	20060906	CN 2004-80021438	20040625
US 2007060753	A1	20070315	US 2003-561754	20051221
PRIORITY APPLN. INFO.:			US 2003-482686P	20030626
			WO 2004-EP6915	20040625
OTHER SOURCE(S):		MARPAT 142:113909		
OI				


10/561,754 7/447 Robert Havlin



AB A process for the preparation of title compds. of formula I [Y = a OH protecting group; R1 = (hetero)aryl; R2-R5 = independently H or alkyl, or R2R3 and/or R4R5 = cycloalkyl; X = CH2, S, CH(OH), etc., n = 0-3] is disclosed. For example, contacting II*2OH with 1N Na2CO3 in EtOAc to move TBOH and oxidation by H2O2 gave III (R = H). Formylation of III with formic acetic anhydride gave III (R = CHO). Reaction of III with HBr salt of N-(5-fluoro-2-pyridinyl)-2-pyrrolidinecarboxamide, followed by oxidation, gave IV. Thus, the present invention provides a process producing the title compound, which are useful to prepare certain antibacterial N-formyl hydroxylamine compds. as peptide deformylase inhibitors.

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

RX(3) OF 10 ...L + O ==> P...



10/561,754 8/447 Robert Havlin



RX(3) RCT L 821774-93-0

STAGE(1)
ROT Q 77-92-9 Citric acid
SOL 7732-18-5 Water, 141-78-6 AcOEt
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 10 minutes, room temperature

STAGE(2)
RCT O 821774-95-2
ROT R 2592-95-2 1-Benzotriazolol, S 109-02-4
N-Methylmorpholine, T 25952-53-8 KDAP
SOL 7732-18-5 Water
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) room temperature -> 5 deg C
SUBSTAGE(3) room temperature
SUBSTAGE(4) overnight

PRO P 478913-92-7

RX(7) OF 10 COMPOSED OF RX(3), RX(4)
RX(7) L + O ==> U



U

RX(3) RCT L 821774-93-0

STAGE(1)

RCT Q 77-92-9 Citric acid
 SOL 7732-18-5 Water, 141-78-6 AcOEt
 CON SUBSTAGE(1) room temperature
 SUBSTAGE(2) 10 minutes, room temperature

STAGE(2)

RCT O 821774-95-2
 RCT R 2592-95-2 1-Benzotriazolol, S 109-02-4
 N-Methylmorpholine, T 25952-53-8 EDAP
 SOL 7732-18-5 Water
 CON SUBSTAGE(1) room temperature
 SUBSTAGE(2) room temperature -> 5 deg C
 SUBSTAGE(3) room temperature
 SUBSTAGE(4) overnight

PRO P 478913-92-7

RX(4) RCT P 478913-92-7
 RCT V 109536-69-8 2-HO2CC6H4CO3H.Mg
 PRO U 478913-93-8
 SOL 108-21-4 Acetic acid, 1-methylethyl ester, 7732-18-5 Water
 CON SUBSTAGE(1) room temperature
 SUBSTAGE(2) 17 hours, room temperature
 NTE workup

=> d cost	
COST IN U.S. DOLLARS	
CONNECT CHARGES	SINCE FILE TOTAL
	ENTRY SESSION
NETWORK CHARGES	0.78 1.32
SEARCH CHARGES	0.12 0.24
DISPLAY CHARGES	113.10 113.10
	6.73 6.73
FULL ESTIMATED COST	120.73 121.39
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE TOTAL
	ENTRY SESSION
CA SUBSCRIBER PRICE	-0.73 -0.73

IN FILE 'CASREACT' AT 08:20:14 ON 30 MAY 2007

=> file reg	
COST IN U.S. DOLLARS	
CONNECT CHARGES	SINCE FILE TOTAL
	ENTRY SESSION
NETWORK CHARGES	121.63 122.29
SEARCH CHARGES	0.12 0.24
DISPLAY CHARGES	113.10 113.10
FULL ESTIMATED COST	120.73 121.39
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE TOTAL
	ENTRY SESSION
CA SUBSCRIBER PRICE	-0.73 -0.73

FILE 'REGISTRY' ENTERED AT 08:21:37 ON 30 MAY 2007
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2007 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file

provided by InfoChem.

STRUCTURE FILE UPDATES: 28 MAY 2007 HIGHEST RN 935999-19-2
 DICTIONARY FILE UPDATES: 28 MAY 2007 HIGHEST RN 935999-19-2

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH December 2, 2006


Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stndoc/properties.html>

>>

Uploading C:\Program Files\Stnexp\Queries\10.561754\clm14 crop1.str



chain nodes :
 1 2 3 12 13 14 15 16 22 23 24
 ring nodes :
 4 5 6 7 8 17 18 19 20 21
 chain bonds :
 1-3 1-2 4-12 12-13 12-14 15-18 15-16 17-22 22-23 22-24
 ring bonds :
 4-5 4-8 5-6 6-7 7-8 17-18 17-21 18-19 19-20 20-21
 exact/norm bonds :
 1-3 1-2 4-5 4-8 4-12 5-6 6-7 7-8 12-13 12-14 15-18 15-16 17-18 17-21
 17-22 18-19 19-20 20-21 22-23 22-24

G1:C,S

Match level :

1:CLASS 2:CLASS 3:CLASS 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 12:CLASS 13:CLASS
 14:CLASS 15:CLASS 16:CLASS 17:Atom 18:Atom 19:Atom 20:Atom 21:Atom 22:CLASS 23:CLASS
 24:CLASS

fragments assigned product role:
 containing 15
 fragments assigned reactant/reagent role:
 containing 1
 containing 4

L4 STRUCTURE UPLOADED

>> d
 L4 HAS NO ANSWERS
 L4 STR

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

Structure attributes must be viewed using STN Express query preparation.

=> file casreact	
COST IN U.S. DOLLARS	
CONNECT CHARGES	SINCE FILE TOTAL
	ENTRY SESSION
NETWORK CHARGES	0.45 122.74
SEARCH CHARGES	0.12 0.24
DISPLAY CHARGES	113.10 113.10
FULL ESTIMATED COST	120.73 121.39
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE TOTAL
	ENTRY SESSION
CA SUBSCRIBER PRICE	0.00 -0.73

FILE 'CASREACT' ENTERED AT 08:21:56 ON 30 MAY 2007
 USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
 COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications.

FILE CONTENT:1840 - 27 May 2007 VOL 146 ISS 23

New CAS Information Use Policies, enter HELP USAGETERMS for details.

 * CASREACT now has more than 12 million reactions *

Some CASREACT records are derived from the ZIC/VINITI database (1974-1999) provided by InfoChem, INPI data prior to 1986, and Biotransformations database compiled under the direction of Professor Dr. Klaus Kieslich.

This file contains CAS Registry Numbers for easy and accurate substance identification.

>> # 14
 SAMPLE SEARCH INITIATED 08:21:59 FILE 'CASREACT'
 SCREENING COMPLETE - 1557 REACTIONS TO VERIFY FROM 83 DOCUMENTS
 100.0% DONE 1557 VERIFIED 930 HIT RXNS 50 DOCS
 INCOMPLETE SEARCH (SYSTEM LIMIT EXCEEDED)
 SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
 BATCH **COMPLETE**
 PROJECTED VERIFICATIONS: 28776 TO 33504
 PROJECTED ANSWERS: 899 TO 1901

LS 50 SRA SSS SAM L4 (930 REACTIONS)


>> d ibib abs hit 1-10


LS ANSWER 1 OF 50 CASREACT COPYRIGHT 2007 ACS ON STN
 ACCESSION NUMBER: 146:142133 CASREACT Full-text
 TITLE: Epimerization Reaction of a Substituted Vinylcyclopropane Catalyzed by Ruthenium Carbenes: Mechanistic Analysis
 AUTHOR(S): Zeng, Xingshong; Wei, Xudong; Farina, Vittorio; Napolitano, Elio; Xu, Yibo; Zhang, Li; Haddad, Nizar; Yee, Nathan K.; Grinberg, Nelu; Shen, Sherry; Senanayake, Chris H.
 CORPORATE SOURCE: Department of Chemical Development, Boehringer Ingelheim Pharmaceuticals, Inc., Ridgefield, CT, 06877, USA
 SOURCE: Journal of Organic Chemistry (2006), 71(23), 8864-8875
 CODEN: JOCEAH; ISSN: 0022-3263
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB A novel ruthenium carbene-catalyzed epimerization of vinylcyclopropanes is reported. The reaction rate strongly depends on the presence of ruthenium ligands in solution. When the first-generation Grubbs catalyst is employed, a 5.3:1 equilibrium ratio of epimers is established quickly, but when a first-generation Hoveyda catalyst is employed, epimerization is observed only if an addl. phosphine or nitrogen ligand is added. NMR and kinetic studies suggest that the isomerization reaction occurs through the intermediacy of a ruthenacyclopentene. The observation suggests that cyclopropylmethylidene ruthenium carbenes of synthetic utility may be accessible via ruthenacyclopentenes obtained via other routes.

REFERENCE COUNT: 71 THERE ARE 71 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

RX(7) OF 77 ...X + Z ==> AA...





RX(7) RCT X 915317-30-5

STAGE(1)


RGT AB 7647-01-0 HCl
SOL 7732-18-5 Water, 123-91-1 Dioxane
CON 3 hours, room temperature

STAGE(2)


RCT Z 769167-55-7
RGT AC 125700-67-6 Benzotriazolium der, R 7087-68-5 EtN(Pr-1)2
SOL 75-09-2 CH2Cl2
CON 1 hour, room temperature

PRO AA 912291-98-6

RX(10) OF 77 3 AI ==> AJ + AF + AA...




PAGE 1-A




* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 1-A





* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *




RX(10) RCT AI 912291-94-2
PRO AJ 912291-95-3, AF 912291-99-7, AA 912291-98-6
CAT 172222-30-9 Ruthenium, dichloro(phenylmethylene)bis(tricyclohexylphosphine)-, (SP-5-31)-
SOL 108-88-3 PhMe
CON 60 deg C

RX(20) OF 77 2 AI + 2 G ==> BK + BL






* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *



RX(20) RCT AI 912291-94-2, G 172222-30-9
PRO BK 919530-94-2, BL 919530-95-3
SOL 1665-00-5 CD2Cl2

CON room temperature

RX(31) OF 77 COMPOSED OF RX(6), RX(7)
RX(31) Q + W + Z ==> AA




RX(6) RCT Q 915317-27-0, M 62-23-7

STAGE(1)

RGT H 603-35-0 PPh3
SOL 109-99-9 THF
CON room temperature -> 0 deg C

STAGE(2)

RGT Y 2446-83-5 N2(CO2CHMe2)2
SOL 109-99-9 THF
CON <5 deg C



YIELD 95%

RX(13) RCT W 653262-12-1
RGT AC 7664-41-7 NH3
PRO AH 917081-06-8
SOL 67-56-1 MeOH
CON 24 - 48 hours, room temperature

LS ANSWER 3 OF 50 CASREACT COPYRIGHT 2007 ACS ON STN

ACCESSION NUMBER: 145:419451 CASREACT [Full-text](#)

TITLE: Rapid and efficient synthesis of the pentapeptide of elastin protein and peptides containing highly hindered α,α -dialkyl amino acids employing Fmoc-amino acid chlorides under microwave irradiation in the solution phase

AUTHOR(S): Tantry, Subramanyam J.; Rao, R. V. Ramana; Babu, V. V. Suresh

CORPORATE SOURCE: Department of Studies in Chemistry, Bangalore University, Bangalore, 560 001, India

SOURCE: ARKIVOC (Gainesville, FL, United States) (2006), (1), 21-30
CODEN: AOPUAR
URL: http://www.arkat-usa.org/ark/journal/2006/101_General/1576/05-15768420ast20published20mainmanuscript.pdf

PUBLISHER: Arkat USA Inc.
DOCUMENT TYPE: Journal; (online computer file)
LANGUAGE: English

AB A rapid and efficient synthesis of peptides in solution employing Fmoc-amino acid chlorides under microwave irradiation is described. A comparison study of the microwave assisted method with those of conventional peptide synthesis using acid chlorides and various coupling additives has been performed. It has been found that, in general, the formation of a peptide bond, employing Fmoc-amino acid chloride and zinc dust or TBDMs-OBt under microwave irradiation is complete in 30-45 s with 90% yield of pure isolated peptide. Employing zinc dust as a coupling additive, the synthesis of several dipeptides,

PRO X 915317-30-5
MTE stereoselective

RX(7) RCT X 915317-30-5

STAGE(1)

RGT AB 7647-01-0 HCl
SOL 7732-18-5 Water, 123-91-1 Dioxane
CON 3 hours, room temperature

STAGE(2)

RCT Z 769167-55-7
RGT AC 125700-67-6 Benzotriazolium der. R 7087-68-5 EtN(Pr-1)2
SOL 75-09-2 CH2Cl2
CON 1 hour, room temperature

PRO AA 912291-58-6

LS ANSWER 2 OF 50 CASREACT COPYRIGHT 2007 ACS ON STN

ACCESSION NUMBER: 146:62144 CASREACT [Full-text](#)

TITLE: Synthesis and biological evaluation of a new category of purine-nucleoside analogues of purine-nucleoside analogues


AUTHOR(S): Li, Da-Liang; Bao, Hong-Li; Tan, Qi-Tao; Ke, Yu-Ping; You, Tian-Pa

CORPORATE SOURCE: Department of Chemistry, University of Science and Technology of China, Hefei, Anhui, 230026, Peop. Rep. China

SOURCE: Chinese Journal of Chemistry (2005), 23(12), 1659-1664
CODEN: CJOCEV; ISSN: 1001-604X

PUBLISHER: Shanghai Institute of Organic Chemistry
DOCUMENT TYPE: Journal
LANGUAGE: English

GI





AB Convenient procedure for coupling of 1,2,3,5-tetra-O-acetyl- β -D-ribofuranose and 4-nitroimidazole was provided to obtain β -anomer as major product. A novel category of nucleoside analogs, e.g. 1, with an imidazole base moiety bearing amino-acid residue was designed and synthesized to develop selective and effective antiviral agents. The title compds. were evaluated for the anti-HBV activity to find that only 1 exhibits cytotoxicity (MTT assay) at IC50 0.3436 μ mol/L and anti-HBV activity at HbeAg and CC50 15.21 μ mol/L.
REFERENCE COUNT: 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


RX(13) OF 51 ...W ==> AH

the pentapeptide fragment Fmoc-Val-Pro-Gly-Val-Gly-OBzl of elastin and the difficult highly hindered couplings of α,α -dialkylamino acids are reported.
REFERENCE COUNT: 27 THERE ARE 27 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

RX(14) OF 31 ...AH + AC ==> AI



AC (14)



YIELD 67%

RX(14) RCT AH 911858-48-5

STAGE(1)

RPT AG 4097-89-6 1,2-Ethanediamine, N1,N1-bis(2-aminoethyl)-
SOL 75-09-2 CH2Cl2
CON 20 minutes, room temperature

STAGE(2)

RCT AC 103321-53-5
RGT D 7440-66-6 Zn
SOL 75-09-2 CH2Cl2
CON 30 seconds

PRO AI 911056-49-6

NTB microwave irradiation in stage 2


L5 ANSWER 4 OF 50 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 145:351168 CASREACT Full-text
TITLE: Protease-Modulated Cellular Uptake of Quantum Dots
AUTHOR(S): Zhang, Yan; So, Min Kyung; Rao, Jianghong
CORPORATE SOURCE: Biophysics, Cancer Biology and Molecular Imaging
Programs, Department of Radiology, Stanford University
School of Medicine, Stanford, CA, 94305-5464, USA
SOURCE: Nano Letters (2006), 6(9), 1988-1992
CODEN: NALEPD; ISSN: 1530-6984
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English

AB Quantum dots (QDs) are often cell-impermeable and require transporters to facilitate crossing over cell membranes. Here the authors present a simple and versatile method that utilizes enzymes, matrix metalloproteinase 2 (MMP-2) and MMP-7, to modulate the cellular uptake of QDs. QD-peptide conjugates could be efficiently taken up into cells after the MMP treatment. This enzyme-modulated cellular uptake of QDs may be applied to other nanoparticles for biol. imaging and selective drug delivery into tumor cells.


REFERENCE COUNT: 38 THERE ARE 38 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

RX(1) OF 14 A + B ==> C


PAGE 1-A



PAGE 1-B




PAGE 1-C




RX(1) RCT A 910217-61-7, B 910217-62-8
RGT D 7087-68-5 EtN(Pr-1)2
PRO C 910217-63-9

RX(4) OF 14 ...A + N ==> O

PAGE 1-A




PAGE 1-B



PAGE 1-C




A




B

(1) →



PAGE 1-A


PAGE 1-B



PAGE 1-C




A




N

(4) →




PAGE 1-A

PAGE 1-B




PAGE 1-C




RX(4) RCT A 910217-61-7, N 910217-64-0
 ROT D 7087-68-5 EtN(Pr-i)
 PRO O 910217-65-1
 SOL 68-12-2 DMF


RX(6) OF 14 ... R + T ==> U




PAGE 1-A




PAGE 1-B



PAGE 1-C




PAGE 1-D




U

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *


PAGE 1-B



PAGE 1-C




PAGE 2-A



(6) →

RX(6) RCT R 35013-72-0, T 910217-66-2
 ROT S 538-75-0 DCC
 PRO U 910217-67-3
 SOL 68-12-2 DMF


RX(7) OF 14 ... R + V ==> W



R

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 1-B



PAGE 1-C






PAGE 2-A




V

(7)




RX (7) RCT R 35013-72-0, V 910217-69-5
 RGT S 538-75-0 DCC
 PRO W 910217-70-8
 SOL 68-12-2 DMF


PAGE 1-C

3
STRE2


PAGE 1-A




PAGE 1-B




RX (10) OF 14 COMPOSED OF RX (3), RX (4)
 RX (10) G + M + A ==> O




PAGE 1-A



PAGE 1-B




PAGE 1-C



RX (3) RCT G 215391-21-2, M 54907-61-8
 RGT D 7087-68-5 EtN(Pr-i)2
 PRO N 910217-64-0
 SOL 68-12-2 DMF


RX (4) RCT A 910217-61-7, N 910217-64-0
 RGT D 7087-68-5 EtN(Pr-i)2
 PRO O 910217-65-1
 SOL 68-12-2 DMF

RX (11) OF 14 COMPOSED OF RX (5), RX (6)
 RX (11) P + Q + T ==> U




* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *


PAGE 1-B




PAGE 1-C




PAGE 2-A

2
STEPS


PAGE 1-A




PAGE 1-B



PAGE 1-C




PAGE 1-D



RX(5) RCT P 58-05-5, Q 6066-82-6
 RGT S 538-75-0 DCC
 PRO R 35013-72-0
 SOL 68-12-2 DMF


RX(6) RCT R 35013-72-0, T 910217-66-2
 RGT S 538-75-0 DCC
 PRO U 910217-67-3
 SOL 68-12-2 DMF

RX(12) OF 14 COMPOSED OF RX(5), RX(7)
 RX(12) P + Q + V ==> W



* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 1-B



PAGE 1-C




PAGE 2-A




V

2
STEPS


PAGE 1-A



PAGE 1-B





PAGE 1-C




RX(5) RCT P 58-05-5, Q 6066-82-6
 RGT S 538-75-0 DCC
 PRO R 35013-72-0
 SOL 68-12-2 DMF

RX(7) RCT R 35013-72-0, V 910217-69-5
 RGT S 538-75-0 DCC
 PRO W 910217-70-8
 SOL 68-12-2 DMF


RX(14) OF 14 COMPOSED OF RX(2), RX(3), RX(4)
 RX(14) 2 E + F + M + A ==> O




PAGE 1-A



PAGE 1-B



PAGE 1-C



RX(2) RCT R 112918-82-8, F 85916-13-8

STAGE(1)

RGT H 94790-37-1 HBTU, D 7087-68-5 EtN(Pr-1)2
SOL 68-12-2 DMF

STAGE(2)

RGT I 110-89-4 Piperidine

STAGE(3)

RGT J 76-05-1 EtCO2H
SOL 75-09-2 CH2Cl2

PRO G 215391-21-2

RX(3) RCT G 215391-21-2, M 54907-61-8

RGT D 7087-68-5 EtN(Pr-1)2

PRO N 910217-64-0

SOL 68-12-2 DMF

RX(4) RCT A 910217-61-7, N 910217-64-0

RGT D 7087-68-5 EtN(Pr-1)2

PRO O 910217-65-1

SOL 68-12-2 DMF

L5 ANSWER 5 OF 50 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER:

145:124445 CASREACT Full-text

TITLE:

Process for preparation of (2S,4S)-1-[(4-nitrobenzyloxycarbonyl)-3-[(3-allyloxycarbonyl)phenylaminocarbonyl]pyrrolidine-4-thiol

INVENTOR(S):

Zhang, Wanbin; Liu, Delong; Luo, Li; Xie, Fang

PATENT ASSIGNER(S):

Shanghai Jiao Tong University, Peop. Rep. China

SOURCE:

Faming Zhuanli Shenqing Gongkai Shuomingshu, 9 pp.

DOCUMENT TYPE:

CODEN: CNXKXV

LANGUAGE:

Patent

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO. KIND DATE APPLICATION NO. DATE


CN 1752073 A 20060329 CN 2005-10030662 20051020

PRIORITY APPL. INPO:


CN 2005-10030662 20051020

AB This invention relates to a method for preparation of (2S,4S)-1-[(4-nitrobenzyloxycarbonyl)-2-[(3-allyloxycarbonyl)phenylaminocarbonyl]pyrrolidine-4-thiol,


PAGE 1-C

3
STEPS

PAGE 1-A





PAGE 1-B



which comprises (1) reacting trans-4-hydroxy-L-proline and p-nitrobenzyloxycarbonyl chloride; (2) reacting m-nitrobenzoic acid and thionyl chloride, then reacting with allyl alc. for allyl m-nitrobenzoate; (3) tin dichloride reducing allyl m-nitrobenzoate for allyl m-aminobenzoate; (4) reacting product of step 1 and allyl m-aminobenzoate; (5) methanesulfonyl chloride treating product of step 4; (6) reacting product of step 5 with potassium thioacetate; (7) hydrolyzing product of step 6 for final product. This invention provides environment friendly method for preparation of title product with low cost.

RX(5) OF 34 ...Q + T ==> U...



RX(5) RCT Q 896731-55-8, T 124-63-0


RGT L 121-44-8 Et3N


PRO U 896731-56-9

SOL 75-09-2 CH2Cl2

CON 40 minutes, room temperature

RX(6) OF 34 ...U + M ==> X...






X


RX (6) RCT U 896731-56-9, W 10387-40-3
 PRO X 153774-58-4
 SOL 68-12-2 DMP
 CON SUBSTAGE (1) 3 hours, room temperature
 SUBSTAGE (2) room temperature -> 70 deg C
 SUBSTAGE (3) 5 hours, 70 deg C

RX (7) OF 34 ... X ==> Z



X

(7)




X

RX (5) RCT Q 896731-55-8, T 124-63-0
 ROT L 121-44-8 Et3N
 PRO U 896731-56-9
 SOL 75-09-3 CH2Cl2
 CON 40 minutes, room temperature

RX (6) RCT U 896731-56-9, W 10387-40-3
 PRO X 153774-58-4
 SOL 68-12-2 DMP
 CON SUBSTAGE (1) 3 hours, room temperature
 SUBSTAGE (2) room temperature -> 70 deg C
 SUBSTAGE (3) 5 hours, 70 deg C


RX (13) OF 34 COMPOSED OF RX (6), RX (7)
 RX (13) U + W ==> Z



U




W



Z

RX (7) RCT X 153774-58-4
 ROT AA 1310-73-2 NaOH
 PRO Z 153775-54-3
 SOL 107-18-6 Allyl alcohol
 CON 20 minutes, 0 deg C
 NTS 40% overall yield from 5


RX (12) OF 34 COMPOSED OF RX (5), RX (6)
 RX (12) Q + T + W ==> X



Q




T



W

2 STEPS




Z

RX (6) RCT U 896731-56-9, W 10387-40-3
 PRO X 153774-58-4
 SOL 68-12-2 DMP
 CON SUBSTAGE (1) 3 hours, room temperature
 SUBSTAGE (2) room temperature -> 70 deg C
 SUBSTAGE (3) 5 hours, 70 deg C

RX (7) RCT X 153774-58-4
 ROT AA 1310-73-2 NaOH
 PRO Z 153775-54-3
 SOL 107-18-6 Allyl alcohol
 CON 20 minutes, 0 deg C
 NTS 40% overall yield from 5

RX (23) OF 34 COMPOSED OF RX (5), RX (6), RX (7)
 RX (23) Q + T + W ==> Z



Q




T



W

3 STEPS




Z

RX(5) RCT Q 896731-55-8, T 124-63-0
 RGT L 121-44-8 Et3N
 PRO U 896731-56-9
 SOL 75-09-2 CH2Cl2
 CON 40 minutes, room temperature

RX(6) RCT U 896731-56-9, W 10387-40-3
 PRO X 153774-58-4
 SOL 68-12-2 DMF
 CON SUBSTAGE(1) 3 hours, room temperature
 SUBSTAGE(2) room temperature -> 70 deg C
 SUBSTAGE(3) 5 hours, 70 deg C


RX(7) RCT X 153774-58-4
 RGT AA 1310-73-2 NaOH
 PRO Z 153775-54-3
 SOL 107-18-6 Allyl alcohol
 CON 20 minutes, 0 deg C
 NTE 40% overall yield from 5

L5 ANSWER 6 OF 50 CASREACT COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 145:117514 CASREACT Full-text
 TITLE: 6-N,N-Dimethylamino-2,3-naphthalimide: A New Environment-Sensitive Fluorescent Probe in δ - and μ -Selective Opioid Peptides
 AUTHOR(S): Vazquez, M. Eugenio; Blanco, Juan B.; Salvadori, Severo; Trapella, Claudio; Argazzi, Roberto; Bryant, Sharon D.; Jimenez, Yunden; Lazarus, Lawrence H.; Negri, Lucia; Giannini, Elisa; Lettanzini, Roberta; Colucci, Mar Antonella; Balboni, Gianfranco
 CORPORATE SOURCE: Departamento de Química Orgánica y Unidad Asociada al CSIC, Universidad de Santiago de Compostela, Santiago de Compostela, 15782, Spain
 SOURCE: Journal of Medicinal Chemistry (2006), 49(12), 3653-3658
 CODEN: JMCMAR; ISSN: 0022-2623
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB A new environment-sensitive fluorophore, 6-N,N-(dimethylamino)-2,3-naphthalimide (6DMN) was introduced in the δ -selective opioid peptide agonist N-Dmt-Tic-Glu-NH2 and in the μ -selective opioid peptide agonist endomorphin-2 (H-Tyr-Pro-Phe-Phe-NH2). Environment-sensitive fluorophores are a special class of chromophores that generally exhibit a low quantum yield in aqueous solution but become highly fluorescent in nonpolar solvents or when bound to hydrophobic sites in proteins or membranes. New fluorescent δ -selective irreversible antagonists (H-Dmt-Tic-Glu-NH-(CH2)5-CO-Dap(6DMN)-NH2 and H-Dmt-Tic-Glu-Dap(6DMN)-NH2) were identified as potential fluorescent probes showing good properties



AB


AF: CM 1



AF: CM 2

(7)

PAGE 1-A




PAGE 1-B

AG
 YIELD 83%

RX(7) RCT AB 897959-57-8, AF 897959-68-1
 RGT K 2592-95-2 1-Benzotriazolol, Z 109-02-4 N-Methylmorpholine, AA


for use in studies of distribution and internalization of δ receptors by confocal laser scanning microscopy.
 REFERENCE COUNT: 44 THERE ARE 44 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

RX(6) OF 25 ...Y ----> AB...



Y

(6)



AB
 YIELD 96%


RX(6) RCT Y 897959-54-5
 RGT AC 1333-74-0 H2
 PRO AB 897959-57-8
 CAT 7440-05-3 Pd
 SOL 67-56-1 MeOH
 CON 1 hour, room temperature

RX(7) OF 25 ...AB + AF ----> AG...

25952-53-8 EDAP
 PRO AG 897959-59-0
 SOL 68-12-2 DMF
 CON SUBSTAGE(1) 3 hours, 0 deg C
 SUBSTAGE(2) 24 hours, room temperature

RX(8) OF 25 ...AG ----> AH...


PAGE 1-A



PAGE 1-B

AG
 YIELD 83%


(8)




AH
 YIELD 83%

RX(8) RCT AG 897959-59-0
 RGT AC 1333-74-0 H2
 PRO AH 897959-61-4
 CAT 7440-05-3 Pd
 SOL 67-56-1 MeOH
 CON 1 hour, room temperature

RX(9) OF 25 ... AH + AI ==> AJ...




AH



AI


(9) →

PAGE 1-A




(10) →

PAGE 1-A



PAGE 1-B




AN
YIELD 90%

RX(10) RCT AJ 897959-62-5
 RGT M 76-05-1 F3CCO2H
 PRO AN 897959-46-5
 SOL 7732-18-5 Water
 CON 30 minutes, room temperature

RX(12) OF 25 COMPOSED OF RX(6), RX(7)
 RX(12) Y + AF ==> AG

PAGE 1-B




AJ
YIELD 50%


RX(9) RCT AH 897959-61-4, AI 3326-32-7
 RGT AK 121-44-8 Et3N
 PRO AJ 897959-62-5
 SOL 64-17-5 EtOH, 109-99-9 THF
 CON 24 hours, room temperature
 NTE in the dark

RX(10) OF 25 ... AJ ==> AN


PAGE 1-A



PAGE 1-B




AJ



Y




AF: CM 1



AF: CM 2

2 STEPS →

PAGE 1-A



PAGE 1-B



AG
YIELD 83%




RX(6) RCT Y 897959-54-5
 RGT AC 1333-74-0 H2
 PRO AB 897959-57-8

CAT 7440-05-3 Pd
 SOL 67-56-1 MeOH
 CON 1 hour, room temperature

RX(7) RCT AB 897959-57-8, AF 897959-68-1
 ROT K 2592-95-2 1-Benzotriazolol, Z 109-02-4 N-Methylmorpholine, AA
 25952-53-8 EDAP
 PRO AG 897959-59-0
 SOL 68-12-2 DMF
 CON SUBSTAGE(1) 3 hours, 0 deg C
 SUBSTAGE(2) 24 hours, room temperature

RX(13) OF 25 COMPOSED OF RX(7), RX(8)

RX(13) AB + AP ==> AH





RX(7) RCT AB 897959-57-8, AF 897959-68-1
 ROT K 2592-95-2 1-Benzotriazolol, Z 109-02-4 N-Methylmorpholine, AA

25952-53-8 EDAP
 PRO AG 897959-59-0
 SOL 68-12-2 DMF
 CON SUBSTAGE(1) 3 hours, 0 deg C
 SUBSTAGE(2) 24 hours, room temperature


RX(8) RCT AG 897959-59-0
 RGT AC 1333-74-0 H2
 PRO AH 897959-61-4
 CAT 7440-05-3 Pd
 SOL 67-56-1 MeOH
 CON 1 hour, room temperature

RX(14) OF 25 COMPOSED OF RX(8), RX(9)


RX(14) AG + AI ==> AJ



PAGE 1-A



PAGE 1-B




RX(8) RCT AG 897959-59-0
 RGT AC 1333-74-0 H2
 PRO AH 897959-61-4
 CAT 7440-05-3 Pd
 SOL 67-56-1 MeOH
 CON 1 hour, room temperature

RX(9) RCT AH 897959-61-4, AI 3326-32-7
 RGT AK 121-44-8 Et3N
 PRO AJ 897959-62-5
 SOL 64-17-5 EtOH, 109-99-9 THF
 CON 24 hours, room temperature
 NTE in the dark


RX(15) OF 25 COMPOSED OF RX(9), RX(10)

RX(15) AH + AI ==> AN

PAGE 1-A




PAGE 1-B




RX(9) RCT AH 897959-61-4, AI 3326-32-7
RGT AK 121-44-8 Et3N
PRO AJ 897959-62-5
SOL 64-17-5 EtOH, 109-99-9 THF
CON 24 hours, room temperature
NTE in the dark

RX(10) RCT AJ 897959-62-5
RGT M 74-05-1 F3CCO2H
PRO AH 897959-46-5
SOL 7732-18-5 Water
CON 30 minutes, room temperature

RX(17) OF 25 COMPOSED OF RX(6), RX(7), RX(8)
RX(17) Y + AF ==> AH




Y



AF: CM 1


AF: CM 2

3 STEPS


AH
YIELD 83%

3 STEPS

PAGE 1-A



PAGE 1-B

AI
YIELD 50%

RX(7) RCT AB 897959-57-8, AF 897959-68-1
RGT K 2592-95-2 1-Benzotriazolol, Z 109-02-4 N-Methylmorpholine, AA
25952-53-8 EDAP
PRO AG 897959-59-0
SOL 68-12-2 DMF
CON SUBSTAGE(1) 3 hours, 0 deg C
SUBSTAGE(2) 24 hours, room temperature

RX(8) RCT AG 897959-59-0
RGT AC 1333-74-0 H2
PRO AH 897959-61-4
CAT 7440-05-3 Pd
SOL 67-56-1 MeOH
CON 1 hour, room temperature


RX(9) RCT AH 897959-61-4, AI 3326-32-7
RGT AK 121-44-8 Et3N
PRO AJ 897959-62-5
SOL 64-17-5 EtOH, 109-99-9 THF
CON 24 hours, room temperature
NTE in the dark

RX(6) RCT Y 897959-54-5
RGT AC 1333-74-0 H2
PRO AB 897959-57-8
CAT 7440-05-3 Pd
SOL 67-56-1 MeOH
CON 1 hour, room temperature

RX(7) RCT AB 897959-57-8, AF 897959-68-1
RGT K 2592-95-2 1-Benzotriazolol, Z 109-02-4 N-Methylmorpholine, AA
25952-53-8 EDAP
PRO AG 897959-59-0
SOL 68-12-2 DMF
CON SUBSTAGE(1) 3 hours, 0 deg C
SUBSTAGE(2) 24 hours, room temperature


RX(8) RCT AG 897959-59-0
RGT AC 1333-74-0 H2
PRO AH 897959-61-4
CAT 7440-05-3 Pd
SOL 67-56-1 MeOH
CON 1 hour, room temperature

RX(19) OF 25 COMPOSED OF RX(7), RX(8), RX(9)
RX(19) AH + AF + AI ==> AJ



AB


AF: CM 1




AF: CM 2

AI

RX(20) OF 25 COMPOSED OF RX(6), RX(7), RX(8), RX(9)
RX(20) Y + AF + AI ==> AJ




Y



AF: CM 1


AF: CM 2




AI

4 STEPS

PAGE 1-A




PAGE 1-B

AJ
YIELD 50%

- RX(6) RCT Y 897959-54-5
RGT AC 1333-74-0 H2
PRO AB 897959-57-8
CAT 7440-05-3 Pd
SOL 67-56-1 MeOH
CON 1 hour, room temperature
- RX(7) RCT AB 897959-57-8, AF 897959-68-1
RGT K 2592-95-2 1-Benzotriazolol, Z 109-02-4 N-Methylmorpholine, AA
25952-53-8 EDAP
PRO AG 897959-59-0
SOL 68-12-2 DMF
CON SUBSTAGE(1) 3 hours, 0 deg C
SUBSTAGE(2) 24 hours, room temperature
- RX(8) RCT AG 897959-59-0
RGT AC 1333-74-0 H2
PRO AH 897959-61-4
CAT 7440-05-3 Pd
SOL 67-56-1 MeOH
CON 1 hour, room temperature
- RX(9) RCT AH 897959-61-4, AI 3326-32-7
RGT AK 121-44-8 Et3N
PRO AJ 897959-62-5
SOL 64-17-5 EtOH, 109-99-9 THF
CON 24 hours, room temperature
NTE in the dark

RX(21) OF 25 COMPOSED OF RX(8), RX(9), RX(10)
RX(21) AG + AI ==> AN


PAGE 1-A



PAGE 1-B




AG




AI

3
STEPS

PAGE 1-A




PAGE 1-B

AN
YIELD 90%

- RX(8) RCT AG 897959-59-0
RGT AC 1333-74-0 H2
PRO AH 897959-61-4
CAT 7440-05-3 Pd
SOL 67-56-1 MeOH
CON 1 hour, room temperature
- RX(9) RCT AH 897959-61-4, AI 3326-32-7
RGT AK 121-44-8 Et3N
PRO AJ 897959-62-5
SOL 64-17-5 EtOH, 109-99-9 THF
CON 24 hours, room temperature
NTE in the dark
- RX(10) RCT AJ 897959-62-5
RGT M 76-05-1 F3CCO2H
PRO AN 897959-66-5
SOL 7732-18-5 Water
CON 30 minutes, room temperature


RX(22) OF 25 COMPOSED OF RX(7), RX(8), RX(9), RX(10)
RX(22) AB + AP + AI ==> AN




AB



AP: CM 1




AP: CM 2




AI

4
STEPS

PAGE 1-A



PAGE 1-B

AN
YIELD 90%

- RX(7) RCT AB 897959-57-8, AF 897959-68-1
RGT K 2592-95-2 1-Benzotriazolol, Z 109-02-4 N-Methylmorpholine, AA
25952-53-8 EDAP
PRO AG 897959-59-0
SOL 68-12-2 DMF
CON SUBSTAGE(1) 3 hours, 0 deg C


SUBSTAGE(2) 24 hours, room temperature

RX(8) RCT AG 897959-59-0
 RGT AC 1333-74-0 H2
 PRO AH 897959-61-4
 CAT 7440-05-3 Pd
 SOL 67-56-1 MeOH
 CON 1 hour, room temperature


RX(9) RCT AH 897959-61-4, AI 3326-32-7
 RGT AK 121-44-8 Et3N
 PRO AJ 897959-62-5
 SOL 64-17-5 EtOH, 109-99-9 THF
 CON 24 hours, room temperature
 NTE in the dark

RX(10) RCT AJ 897959-62-5
 RGT M 76-05-1 F3CCO2H
 PRO AN 897959-46-5
 SOL 7732-18-5 Water
 CON 30 minutes, room temperature

RX(24) OF 25 COMPOSED OF RX(6), RX(7), RX(8), RX(9), RX(10)
 RX(24) Y + AF + AI ==> AN



Y



AF: CH 1

AF: CH 2

RX(8) RCT AG 897959-59-0
 RGT AC 1333-74-0 H2
 PRO AH 897959-61-4
 CAT 7440-05-3 Pd
 SOL 67-56-1 MeOH
 CON 1 hour, room temperature

RX(9) RCT AH 897959-61-4, AI 3326-32-7
 RGT AK 121-44-8 Et3N
 PRO AJ 897959-62-5
 SOL 64-17-5 EtOH, 109-99-9 THF
 CON 24 hours, room temperature
 NTE in the dark


RX(10) RCT AJ 897959-62-5
 RGT M 76-05-1 F3CCO2H
 PRO AN 897959-46-5
 SOL 7732-18-5 Water
 CON 30 minutes, room temperature

LS ANSWER 7 OF 50 CASREACT COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 144:350944 CASREACT Full-text
 TITLE: Application of triazine "superactive esters" in the
 repetitive synthesis of oligopeptides. Part 1.
 AUTHOR(S): Synthesis of [54-59] fragment of human β -casein
 Kaminski, Zbigniew J.; Saleh, Bashar; Kolesinska,
 Beata; Redlinski, Adam; Rudzinski, Juliusz
 CORPORATE SOURCE: Institute of Organic Chemistry, Technical University
 of Lodz, Lodz, 90-924, Pol.
 SOURCE: Acta Polonica Pharmaceutica (2005), 62(1), 53-57
 CODEN: APHAX; ISSN: 0001-6837
 PUBLISHER: Polish Pharmaceutical Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English


AB A new synthetic protocol which considerably improves the classic RMA (repetitive excess mixed anhydrides) procedure is proposed. The modification is based on the application of triazine "superactive esters" as superior substitutes for mixed anhydrides, which have been used as acylating reagent in the classical procedure. The improved repetitive procedure in solution was applied to the preparation of [54-59] fragment of human β -casein. The structure and high purity of the intermediates, as well as of the final products, was confirmed by FAB-MS, ¹H-NMR and HPLC.

REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RS FORMAT

RX(2) OF 9 ...C ==> G...




(2)




5 STEPS

PAGE 1-A




PAGE 1-B



AN YIELD 90%

RX(6) RCT Y 897959-54-5
 RGT AC 1333-74-0 H2
 PRO AB 897959-57-8
 CAT 7440-05-3 Pd
 SOL 67-56-1 MeOH
 CON 1 hour, room temperature


RX(7) RCT AB 897959-57-8, AF 897959-68-1
 RGT K 2592-95-2 1-Benzotriazolol, Z 109-02-4 N-Methylmorpholine, AA 25952-53-8 EDAP
 PRO AG 897959-59-0
 SOL 68-12-2 DMF
 CON SUBSTAGE(1) 3 hours, 0 deg C
 SUBSTAGE(2) 24 hours, room temperature




Q YIELD 80%

RX(2) RCT C 19669-38-6
 RGT H 7664-41-7 NH3
 PRO G 200954-13-4
 SOL 67-56-1 MeOH
 CON SUBSTAGE(1) <0 deg C
 SUBSTAGE(2) <0 deg C -> room temperature
 SUBSTAGE(3) 72 hours, room temperature

RX(3) OF 9 ...O + J ==> K



(3)



K YIELD 95%

RX(3) RCT G 200954-43-4

STAGE(1)

RGT L 7647-01-0 HCl
 SOL 64-19-7 AcOH
 CON room temperature

STAGE(2)


RCT J 13139-16-7
 RGT D 3140-73-6 Cl-(MeO)2-s-triazine, E 109-02-4
 N-Methylmorpholine
 SOL 109-99-9 THF, 68-12-3 DMF
 CON SUBSTAGE(1) room temperature -> 0 deg C
 SUBSTAGE(2) 0 deg C
 SUBSTAGE(3) 4 hours, 0 deg C
 SUBSTAGE(4) 3 hours, 0 deg C
 SUBSTAGE(5) 0 deg C -> room temperature
 SUBSTAGE(6) overnight, room temperature

STAGE(3)

RGT M 298-14-6 KHCO3
 SOL 7732-18-5 Water
 CON 1 hour, room temperature


PRO K 881492-63-3

RX(4) OF 9 A + Q ==> R



● HCl

(4) →

R
YIELD 82%

RX(4) RCT A 15761-39-4

STAGE(1)

RGT D 3140-73-6 Cl-(MeO)2-s-triazine, E 109-02-4
 N-Methylmorpholine
 SOL 109-99-9 THF
 CON SUBSTAGE(1) room temperature -> 0 deg C
 SUBSTAGE(2) 0 deg C
 SUBSTAGE(3) 4 hours, 0 deg C

STAGE(2)


RCT Q 872333-21-6
 CON SUBSTAGE(1) 3 hours, 0 deg C
 SUBSTAGE(2) 0 deg C -> room temperature
 SUBSTAGE(3) overnight, room temperature

STAGE(3)


RGT M 298-14-6 KHCO3
 SOL 7732-18-5 Water
 CON 1 hour, room temperature

PRO R 881492-64-4

RX(5) OF 9 S + T ==> U




S



● HCl

(5) →

U
YIELD 98%

RX(5) RCT S 13574-13-5

STAGE(1)

RGT D 3140-73-6 Cl-(MeO)2-s-triazine, E 109-02-4
 N-Methylmorpholine
 SOL 109-99-9 THF
 CON SUBSTAGE(1) room temperature -> 0 deg C
 SUBSTAGE(2) 0 deg C
 SUBSTAGE(3) 4 hours, 0 deg C

STAGE(2)


RCT T 881492-67-7
 CON SUBSTAGE(1) 3 hours, 0 deg C
 SUBSTAGE(2) 0 deg C -> room temperature
 SUBSTAGE(3) overnight, room temperature

STAGE(3)

RGT M 298-14-6 KHCO3
 SOL 7732-18-5 Water
 CON 1 hour, room temperature

PRO U 881492-65-5

RX(6) OF 9 V + W ==> X




V

<-----User Break----->


RX(33) RCT AO 99298-06-3, BO 874163-26-5
 PRO BR 874163-22-1
 SOL 75-05-8 MeCN
 CON 18 hours, reflux

RX(47) OF 285 COMPOSED OF RX(11), RX(15)

RX(47) Y ==> AS



Y

2
STEPS
→AS
YIELD 92%=>
=> d hist

(FILE 'HOME' ENTERED AT 08:18:24 ON 30 MAY 2007)

FILE 'REGISTRY' ENTERED AT 08:18:37 ON 30 MAY 2007
 STRUCTURE UPLOADED

FILE 'CASREACT' ENTERED AT 08:19:13 ON 30 MAY 2007

L2 0 S LI SSS SAM
L3 1 S LI SSS FULL
FILE 'REGISTRY' ENTERED AT 08:21:37 ON 30 MAY 2007
L4 STRUCTURE UPLOADED
FILE 'CASREACT' ENTERED AT 08:21:56 ON 30 MAY 2007
L5 50 S L4

=> # 15 not py>2003
COMMAND INTERRUPTED
REENTER FILE 'CASREACT'
AND TRY AGAIN, OR ENTER '?' FOR MORE INFORMATION.
Your command did not complete due to a temporary system problem. To recover, reenter the file you are in now. Then, any command that is normally available to you may be used. No cost summary for the current file will be displayed. After reentering the current file you may retry your command. Also, you may wish to SAVE your search query. This can be done in any file. If you cannot access your current file, or if your command fails a second time, notify the Help Desk. Enter "HELP STN" for information on contacting the nearest STN Help Desk by telephone or by using the SEND command in STNMAIL file.

	SINCE FILE	TOTAL
COST IN U.S. DOLLARS	ENTRY	SESSION
FULL ESTIMATED COST	1.80	124.54
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	0.00	-0.73

FILE 'CASREACT' ENTERED AT 06:24:06 ON 30 MAY 2007
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications.

FILE CONTENT:1840 - 27 May 2007 VOL 146 ISS 23

New CAS Information Use Policies, enter HELP USAGETERMS for details.

* CASREACT now has more than 12 million reactions *

Some CASREACT records are derived from the ZIC/VINITI database (1974-1999) provided by InfoChem, INPI data prior to 1986, and Biotransformations database compiled under the direction of Professor Dr. Klaus Kieslich.

This file contains CAS Registry Numbers for easy and accurate substance identification.


=> # 15 not py>2003
101431 PY>2003
L6 28 L5 NOT PY>2003
=> d ibib abs hit 1-10

L6 ANSWER 1 OF 28 CASREACT COPYRIGHT 2007 ACS ON STN
ACCESSION NUMBER: 140.287687 CASREACT Full-text
TITLE: Synthetic glycopeptides of the tandem repeat sequence of the epithelial mucin MUC4 with tumor-associated carbohydrate antigens
AUTHOR(S): Brocks, Constanze; Kunz, Horst
CORPORATE SOURCE: Institut fuer Organische Chemie, Johannes Gutenberg-Universitaet Mainz, Mainz, 55128, Germany
SOURCE: Synlett (2003), (13), 2052-2056
CODEN: SYNLES; ISSN: 0936-5214
PUBLISHER: Georg Thieme Verlag
DOCUMENT TYPE: Journal
LANGUAGE: English
AB Glycohexadecaepptides representing the tandem repeat sequence of the epithelial mucin MUC4 were prepared by applying a solid-phase methodol. The required glycosyl amino acid building blocks containing the tumor-associated saccharide antigens TN-, T-, sialyl-TN, (2,6)- and (2,3)-sialyl-T were synthesized according to a straightforward biomimetic strategy by step-wise extension of the saccharide side chain of a Fmoc-protected galactosamine threonine tert-Bu ester.
REFERENCE COUNT: 43 THERE ARE 43 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


RX(20) OF 187 ...BK ==> BL

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 1-B



PAGE 2-A




BK


(20)

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 1-B



PAGE 2-A



BL
YIELD 80%

RX(20) RCT BK 673476-54-5

STAGE(1)
ROT BL 1333-74-0 H2
CAT 7440-05-9 Pd
SOL 67-56-1 MeOH
CON room temperature


STAGE(2)
ROT C 124-41-4 NaOMe
SOL 67-56-1 MeOH
CON room temperature, pH 9

PRO BL 673476-46-5


RX(21) OF 187 ...BM ==> BN

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 1-B



PAGE 2-A



BM

(21)


* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *



PAGE 2-A

PAGE 2-B

BN
YIELD 72%

RX(21) RCT BM 688346-62-1

STAGE(1)

RGT BI 1333-74-0 H2
CAT 7440-05-3 Pd
SOL 67-56-1 MeOH
CON room temperature

STAGE(2)


RGT C 124-41-4 NaOMe
SOL 67-56-1 MeOH
CON room temperature, pH 9

PRO BN 673476-47-6

RX(22) OF 187 ...BO ==> BP

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *


PAGE 1-B



OAc


OAc

PAGE 2-A



HO

PAGE 2-B

BP
YIELD 55%

RX(22) RCT BO 673476-56-7

STAGE(1)

RGT BI 1333-74-0 H2
CAT 7440-05-3 Pd
SOL 67-56-1 MeOH
CON room temperature

STAGE(2)


RGT C 124-41-4 NaOMe
SOL 67-56-1 MeOH
CON room temperature, pH 9

PRO BP 673476-48-7

RX(23) OF 187 ...BQ ==> BR


* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 2-A



AcO

PAGE 2-B




BO

(22)

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 1-B




OH

OH


PAGE 1-B



PAGE 2-A



PAGE 2-B



BQ


(23)

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *


PAGE 1-B



PAGE 2-A




PAGE 2-B




BR
YIELD 64 %


PAGE 1-B



PAGE 2-A



PAGE 2-B



BT
YIELD 59 %

RX (24) RCT BS 673476-58-9

STAGE(1)
RGT BI 1333-74-0 H2
CAT 7440-05-3 Pd
SOL 67-56-1 MeOH
CON room temperature

RX (23) RCT BQ 600346-63-2

STAGE(1)

RGT BI 1333-74-0 H2
CAT 7440-05-3 Pd
SOL 67-56-1 MeOH
CON room temperature

STAGE(2)

RGT C 124-41-4 NaOMe
SOL 67-56-1 MeOH
CON room temperature, pH 9


PRO BR 673476-49-8

RX (24) OF 187 ...BS ==> BT

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 2-A



PAGE 2-B



* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

(24)

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

STAGE(2)

RGT C 124-41-4 NaOMe
SOL 67-56-1 MeOH
CON room temperature, pH 10

STAGE(3)


RGT BU 1310-73-2 NaOH
SOL 7732-18-5 Water
CON room temperature, pH 11.5

PRO BT 673476-50-1


RX (25) OF 187 ...BV ==> BW

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 1-B



PAGE 2-A




BV

(25)

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 1-B



* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 2-B



RX(25) RCT BV 673476-59-0

STAGE(1)

RGT BI 1333-74-0 H2
CAT 7440-05-3 Pd
SOL 67-56-1 MeOH
CON room temperature

STAGE(2)

RGT C 124-41-4 NaOMe
SOL 67-56-1 MeOH
CON room temperature, pH 9

PRO BV 673476-51-2

RX(26) OF 187 ...BX ==> BY

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(26) RCT BX 688348-64-3

STAGE(1)

RGT BI 1333-74-0 H2
CAT 7440-05-3 Pd
SOL 67-56-1 MeOH
CON room temperature

STAGE(2)

RGT C 124-41-4 NaOMe
SOL 67-56-1 MeOH
CON room temperature, pH 9

PRO BY 673476-52-3


RX(27) OF 187 ...BZ ==> CA

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *


PAGE 1-B




PAGE 2-A



PAGE 2-B



PAGE 3-A



BX

(26)

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *


PAGE 2-B



PAGE 3-A

BY
YIELD 67%

PAGE 3-A



BZ

(27)

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(27) RCT BZ 673476-61-4

STAGE(1)

RGT BI 1333-74-0 H2
CAT 7440-05-3 Pd
SOL 67-56-1 MeOH
CON room temperature

STAGE(2)

RGT C 124-41-4 NaOMe
SOL 67-56-1 MeOH
CON room temperature, pH 9

PRO CA 673476-53-4

L6 ANSWER 2 OF 28 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 140:236064 CASREACT Full-text

TITLE: Application of Pac ester in thioester method for the
synthesis of cyclopentapeptides

AUTHOR(S): Liu, Mian; Tian, Qiu-Ling; Ye, Yun-Hua
CORPORATE SOURCE: Key Laboratory of Bioorganic Chemistry and Molecular
Engineering, Ministry of Education, Department of
Chemistry, Peking University, Beijing, 100871, Peop.
Rep. China

SOURCE: Chinese Journal of Chemistry (2003), 21(7), 864-870
CODEN: CJOCV; ISSN: 1001-604X

PUBLISHER: Science Press

DOCUMENT TYPE: Journal


LANGUAGE: English

AB The thioester method for the synthesis of cyclopeptides is improved by using Pac (Pac = phenacyl, CH₂COC₆H₅) ester as a protecting group for 3-mercaptopropionic acid. The Pac group is easily removed from the C-terminal using zinc in acetic acid. The protected peptide thioesters synthesized by the improved method are easily purified for use in subsequent cyclization. Furthermore, this method is flexible for use in peptide chain


elongation, either from the C-terminal or the N-terminal. Two N-protected pentapeptide thioesters, Boc-Pro-Tyr-Leu-Ala-Gly-SCH₂CH₂COOPac and Boc-Ala-Tyr-Leu-Ala-Gly-SCH₂CH₂COOPac, were synthesized by the improved thioester method. After deprotecting the Pac ester with zinc in aqueous acetic acid and the Boc group with trifluoroacetic acid in CH₂Cl₂, two free pentapeptide thioesters were obtained. Ag-assisted cyclization in acetate buffered solution afforded cyclic pentapeptides cyclo(Pro-Tyr-Leu-Ala-Gly) and cyclo(Ala-Tyr-Leu-Ala-Gly). Effects of different buffer pH, Ag⁺ concns., etc. on the cyclization were studied.

REFERENCE COUNT: 20 THERE ARE 20 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

RX(4) OF 77 ...N + O ==> P...




N



O

(4)



P
YIELD 51%

RX(4) RCT N 503440-97-9

STAGE(1)

RGT H 7647-01-0 HCl

SOL 141-78-6 AcOEt

RX(15) RCT AK 667905-07-9

STAGE(1)

RGT H 7647-01-0 HCl

SOL 141-78-6 AcOEt

CON 20 minutes, room temperature

STAGE(2)

RCT O 132149-57-6

RGT B 165514-43-0 1,2,3-Benzotriazin-4(3H)-one,

3-[(diethoxyphosphoryl)oxy]-, Al 121-44-8 Et3N


SOL 68-12-2 DMF

CON overnight, room temperature


PRO AL 667905-09-1

NTE intermediate product could be isolated

RX(16) OF 77 ...P ==> T...



P



T
YIELD 83%

RX(16) RCT P 503440-98-0
RGT AP 64-19-7 AcOH, AQ 7440-66-6 Zn
PRO T 854749-15-8
SOL 64-19-7 AcOH, 7732-18-5 Water

CON 15 minutes, room temperature

STAGE(2)

RCT O 132149-57-6

RGT I 2592-95-2 1-Benzotriazolol, D 7087-68-5 EtN(Pr-i)2

SOL 109-99-9 THF

CON SUBSTAGE(1) room temperature, pH 7

SUBSTAGE(2) room temperature -> 0 deg C

STAGE(3)

RGT J 538-75-0 DCC


CON SUBSTAGE(1) 1 hour, 0 deg C

SUBSTAGE(2) overnight, room temperature


PRO P 503440-98-0

NTE intermediate product could be isolated

RX(15) OF 77 ...AK + O ==> AL...




AK



O

(15)




AL
YIELD 70%


CON 1 hour, room temperature

RX(20) OF 77 COMPOSED OF RX(3), RX(4)


RX(20) O + M + O ==> P



O




N



O

2
STEPS



P
YIELD 51%

RX(3) RCT O 503440-96-0

STAGE(1)

RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 20 minutes, room temperature

STAGE(2)

RCT M 3350-19-4
RGT D 7087-68-5 EtN(Pr-i)2
SOL 68-12-2 DMF
CON 3 hours, room temperature, pH 8 - 9

PRO N 503440-97-9

NTE intermediate product could be isolated

RX(4) RCT N 503440-97-9

STAGE(1)

RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 15 minutes, room temperature

STAGE(2)

RCT O 132149-57-6
RGT I 2592-95-2 1-Benzotriazolol, D 7087-68-5 EtN(Pr-i)2
SOL 109-99-9 THF
CON SUBSTAGE(1) room temperature, pH 7
SUBSTAGE(2) room temperature -> 0 deg C

STAGE(3)



RGT J 538-75-0 DCC
CON SUBSTAGE(1) 1 hour, 0 deg C
SUBSTAGE(2) overnight, room temperature

PRO P 503440-98-0


NTE intermediate product could be isolated

RX(22) OF 77 COMPOSED OF RX(4), RX(16)

RX(22) N + O ==> T



2 STEPS



YIELD 83%

RX(4) RCT N 503440-97-9

STAGE(1)

RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 15 minutes, room temperature

STAGE(2)

RCT O 132149-57-6
RGT I 2592-95-2 1-Benzotriazolol, D 7087-68-5 EtN(Pr-i)2
SOL 109-99-9 THF
CON SUBSTAGE(1) room temperature, pH 7
SUBSTAGE(2) room temperature -> 0 deg C

STAGE(3)

RGT J 538-75-0 DCC
CON SUBSTAGE(1) 1 hour, 0 deg C
SUBSTAGE(2) overnight, room temperature

PRO P 503440-98-0


NTE intermediate product could be isolated

RX(16) RCT P 503440-98-0


RGT AP 64-19-7 AcOH, AQ 7440-66-6 Zn
PRO T 854749-15-8
SOL 64-19-7 AcOH, 7732-18-5 Water
CON 1 hour, room temperature

RX(28) OF 77 COMPOSED OF RX(12), RX(15)


RX(28) AH + AJ + O ==> AL



AJ



2 STEPS



YIELD 70%

RX(12) RCT AH 667905-06-8

STAGE(1)

RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 20 minutes, room temperature

STAGE(2)

RCT AJ 13139-15-6
RGT S 165534-43-0 1,2,3-Benzotriazin-4(3H)-one, 3-[(diethoxyphosphinyl)oxy]-, AI 121-44-8 Et3N
SOL 68-12-2 DMF
CON 24 hours, room temperature

PRO AK 667905-07-9

NTE intermediate product could be isolated

RX(15) RCT AK 667905-07-9

STAGE(1)

RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 20 minutes, room temperature

STAGE(2)


RCT O 132149-57-6
RGT S 165534-43-0 1,2,3-Benzotriazin-4(3H)-one, 3-[(diethoxyphosphinyl)oxy]-, AI 121-44-8 Et3N
SOL 68-12-2 DMF
CON overnight, room temperature

PRO AL 667905-09-1



NTE intermediate product could be isolated

RX(35) OF 77 COMPOSED OF RX(2), RX(3), RX(4)


RX(35) C + F + M + O ==> P



F



3 STEPS



P
YIELD 51%

RX(2) RCT C 503440-95-7

STAGE(1)

RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 15 minutes, room temperature

STAGE(2)

RCT F 15761-38-3
RGT I 2592-95-2 1-Benzotriazolol, D 7087-68-5 EtN(Pr-i)2
SOL 109-99-9 THF
CON SUBSTAGE(1) room temperature, pH 7
SUBSTAGE(2) room temperature -> 0 deg C

STAGE(3)

RGT J 538-75-0 DCC
CON SUBSTAGE(1) 1 hour, 0 deg C
SUBSTAGE(2) overnight, room temperature

PRO G 503440-96-8

NTE intermediate product could be isolated

RX(3) RCT G 503440-96-8

STAGE(1)

RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 20 minutes, room temperature

STAGE(2)

RCT M 3350-19-4
RGT D 7087-68-5 EtN(Pr-i)2
SOL 68-12-2 DMF
CON 3 hours, room temperature, pH 8 - 9


PRO N 503440-97-9

NTE intermediate product could be isolated

RX(4) RCT N 503440-97-9

STAGE(1)

RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 15 minutes, room temperature



P
YIELD 51%

RX(1) RCT A 133367-03-0, B 70-11-1

RGT D 7087-68-5 EtN(Pr-i)2
PRO C 503440-95-7
SOL 68-12-2 DMF
CON overnight, room temperature

RX(2) RCT C 503440-95-7

STAGE(1)

RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 15 minutes, room temperature

STAGE(2)

RCT F 15761-38-3
RGT I 2592-95-2 1-Benzotriazolol, D 7087-68-5 EtN(Pr-i)2
SOL 109-99-9 THF
CON SUBSTAGE(1) room temperature, pH 7
SUBSTAGE(2) room temperature -> 0 deg C

STAGE(3)

RGT J 538-75-0 DCC
CON SUBSTAGE(1) 1 hour, 0 deg C
SUBSTAGE(2) overnight, room temperature

PRO G 503440-96-8

NTE intermediate product could be isolated

RX(3) RCT G 503440-96-8

STAGE(1)

RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 20 minutes, room temperature

STAGE(2)

RCT M 3350-19-4
RGT D 7087-68-5 EtN(Pr-i)2
SOL 68-12-2 DMF
CON 3 hours, room temperature, pH 8 - 9

PRO N 503440-97-9

NTE intermediate product could be isolated

RX(4) RCT N 503440-97-9

STAGE(2)

RCT O 132149-57-6
RGT I 2592-95-2 1-Benzotriazolol, D 7087-68-5 EtN(Pr-i)2
SOL 109-99-9 THF
CON SUBSTAGE(1) room temperature, pH 7
SUBSTAGE(2) room temperature -> 0 deg C




STAGE(3)

RGT J 538-75-0 DCC
CON SUBSTAGE(1) 1 hour, 0 deg C
SUBSTAGE(2) overnight, room temperature

PRO P 503440-98-0

NTE intermediate product could be isolated

RX(37) OF 77 COMPOSED OF RX(1), RX(2), RX(3), RX(4)
RX(37) A + B + F + M + O ==> P



STAGE(1)

RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 15 minutes, room temperature

STAGE(2)

RCT O 132149-57-6
RGT I 2592-95-2 1-Benzotriazolol, D 7087-68-5 EtN(Pr-i)2
SOL 109-99-9 THF
CON SUBSTAGE(1) room temperature, pH 7
SUBSTAGE(2) room temperature -> 0 deg C




STAGE(3)


RGT J 538-75-0 DCC
CON SUBSTAGE(1) 1 hour, 0 deg C
SUBSTAGE(2) overnight, room temperature

PRO P 503440-98-0

NTE intermediate product could be isolated

RX(39) OF 77 COMPOSED OF RX(3), RX(4), RX(16)
RX(39) G + M + O ==> T





T
YIELD 83%

RX(3) RCT G 503440-96-8

STAGE(1)

RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 20 minutes, room temperature

STAGE(2)

RCT M 3350-19-4
RGT D 7087-68-5 EtN(Pr-i)2
SOL 68-12-2 DMF
CON 3 hours, room temperature, pH 8 - 9

PRO N 503440-97-9

NTE intermediate product could be isolated

RX(4) RCT N 503440-97-9

STAGE(1)

RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 15 minutes, room temperature

STAGE(2)

RCT O 132149-57-6
RGT I 2592-95-2 1-Benzotriazolol, D 7087-68-5 EtN(Pr-i)2
SOL 109-99-9 THF
CON SUBSTAGE(1) room temperature, pH 7
SUBSTAGE(2) room temperature -> 0 deg C

STAGE(3)

RGT J 538-75-0 DCC
CON SUBSTAGE(1) 1 hour, 0 deg C
SUBSTAGE(2) overnight, room temperature

PRO P 503440-98-0

NTE intermediate product could be isolated





RX(16) RCT P 503440-98-0

RGT AP 64-19-7 AcOH, AQ 7440-66-6 Zn

PRO T 854749-15-8

SOL 64-19-7 AcOH, 7732-18-5 Water

RX(41) OF 77 COMPOSED OF RX(2), RX(3), RX(4), RX(16)
RX(41) C + F + M + O ==> T



T
YIELD 83%

RX(2) RCT C 503440-95-7

STAGE(1)

RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 15 minutes, room temperature

STAGE(2)

RCT F 15761-38-3
RGT I 2592-95-2 1-Benzotriazolol, D 7087-68-5 EtN(Pr-i)2
SOL 109-99-9 THF
CON SUBSTAGE(1) room temperature, pH 7
SUBSTAGE(2) room temperature -> 0 deg C

STAGE(3)

RGT J 538-75-0 DCC
CON SUBSTAGE(1) 1 hour, 0 deg C
SUBSTAGE(2) overnight, room temperature

PRO G 503440-96-8

NTE intermediate product could be isolated

RX(3) RCT G 503440-96-8

STAGE(1)

RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 20 minutes, room temperature

STAGE(2)

RCT M 3350-19-4
RGT D 7087-68-5 EtN(Pr-i)2
SOL 68-12-2 DMF
CON 3 hours, room temperature, pH 8 - 9

PRO N 503440-97-9

NTE intermediate product could be isolated

RX(4) RCT N 503440-97-9

STAGE(1)

RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 15 minutes, room temperature

STAGE(2)

RCT O 132149-57-6
RGT I 2592-95-2 1-Benzotriazolol, D 7087-68-5 EtN(Pr-i)2
SOL 109-99-9 THF
CON SUBSTAGE(1) room temperature, pH 7
SUBSTAGE(2) room temperature -> 0 deg C

STAGE(3)

RGT J 538-75-0 DCC
CON SUBSTAGE(1) 1 hour, 0 deg C
SUBSTAGE(2) overnight, room temperature



PRO P 503440-98-0

NTE intermediate product could be isolated


RX(16) RCT P 503440-98-0

RGT AP 64-19-7 AcOH, AQ 7440-66-6 Zn
PRO T 854749-15-8
SOL 64-19-7 AcOH, 7732-18-5 Water
CON 1 hour, room temperature

RX(48) OF 77 COMPOSED OF RX(11), RX(12), RX(15)
RX(48) AF + F + AJ + O ==> AL



3
STEPS



AL
YIELD 70%

RX(11) RCT AP 667905-05-7

STAGE(1)

RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 15 minutes, room temperature

STAGE(2)

RCT F 15761-38-3
RGT I 2592-95-2 1-Benzotriazolol, AI 121-44-8 Et3N
SOL 109-99-9 THF
CON room temperature

STAGE(3)

RGT J 538-75-0 DCC
CON overnight, room temperature

PRO AH 667905-06-8

NTE intermediate product could be isolated

RX(12) RCT AH 667905-06-8

STAGE(1)

RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 20 minutes, room temperature

STAGE(2)

RCT AJ 13139-15-6
RGT S 165534-43-0 1,2,3-Benzotriazin-4(3H)-one,
3-[(diethoxyphosphinyl)oxy]-, AI 121-44-8 Et3N
SOL 68-12-2 DMF
CON 24 hours, room temperature

PRO AK 667905-07-9

NTE intermediate product could be isolated

RX(15) RCT AK 667905-07-9

STAGE(1)

RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 20 minutes, room temperature

STAGE(2)




RCT O 132149-57-6
RGT S 165534-43-0 1,2,3-Benzotriazin-4(3H)-one,
3-[(diethoxyphosphinyl)oxy]-, AI 121-44-8 Et3N
SOL 68-12-2 DMF
CON overnight, room temperature

PRO AL 667905-09-1

NTE intermediate product could be isolated

RX(49) OF 77 COMPOSED OF RX(10), RX(11), RX(12), RX(15)

RX(49) AD + AE + F + AJ + O ==> AL

4
STEPSAL
YIELD 70%

RX(10) RCT AD 4530-20-5, AE 100-53-8

STAGE(1)

RGT AG 1122-58-3 4-DMAP
SOL 75-09-2 CH2Cl2
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) room temperature -> 0 deg C

STAGE(2)

RGT J 538-75-0 DCC
CON SUBSTAGE(1) 1 hour, 0 deg C
SUBSTAGE(2) overnight, room temperature

PRO AP 667905-05-7

RX(11) RCT AP 667905-05-7

STAGE(1)

RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 15 minutes, room temperature

STAGE(2)

RCT F 15761-38-3
RGT I 2592-95-2 1-Benzotriazolol, AI 121-44-8 Et3N
SOL 109-99-9 THF
CON room temperature

STAGE(3)

RGT J 538-75-0 DCC
CON overnight, room temperature

PRO AH 667905-06-8

NTE intermediate product could be isolated

RX(12) RCT AH 667905-06-8

STAGE(1)

RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 20 minutes, room temperature

STAGE(2)

RCT AJ 13139-15-6
RGT S 165534-43-0 1,2,3-Benzotriazin-4(3H)-one,
3-[(diethoxyphosphinyl)oxy]-, AI 121-44-8 Et3N
SOL 68-12-2 DMF
CON 24 hours, room temperature

PRO AK 667905-07-9

NTE intermediate product could be isolated

RX(15) RCT AK 667905-07-9

STAGE(1)

RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 20 minutes, room temperature

STAGE(2)





RCT O 132149-57-6
RGT S 165534-43-0 1,2,3-Benzotriazin-4(3H)-one,
3-[(diethoxyphosphinyl)oxy]-, AI 121-44-8 Et3N
SOL 68-12-2 DMF
CON overnight, room temperature

PRO AL 667905-09-1

NTE intermediate product could be isolated

RX(57) OF 77 COMPOSED OF RX(1), RX(2), RX(3), RX(4), RX(16)

RX(57) A + B + F + M + O ==> T

T
YIELD 83%

RX(1) RCT A 133367-03-0, B 70-11-1
RGT D 7087-68-5 EtN(Pr-i)2
PRO C 503440-95-7
SOL 68-12-2 DMF

CON overnight, room temperature

RX(2) RCT C 503440-95-7

STAGE(1)
RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 15 minutes, room temperature

STAGE(2)
RCT F 15761-38-3
RGT I 2592-95-2 1-Benzotriazolol, D 7087-68-5 EtN(Pr-1)2
SOL 109-99-9 THF
CON SUBSTAGE(1) room temperature, pH 7
SUBSTAGE(2) room temperature -> 0 deg C

STAGE(3)
RGT J 538-75-0 DCC
CON SUBSTAGE(1) 1 hour, 0 deg C
SUBSTAGE(2) overnight, room temperature

PRO G 503440-96-8
NTE intermediate product could be isolated

RX(3) RCT G 503440-96-8

STAGE(1)
RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 20 minutes, room temperature

STAGE(2)
RCT M 3350-19-4
RGT D 7087-68-5 EtN(Pr-1)2
SOL 68-12-2 DMF
CON 3 hours, room temperature, pH 8 - 9

PRO N 503440-97-9
NTE intermediate product could be isolated

RX(4) RCT N 503440-97-9

STAGE(1)
RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 15 minutes, room temperature

STAGE(2)
RCT O 132149-57-6
RGT I 2592-95-2 1-Benzotriazolol, D 7087-68-5 EtN(Pr-1)2
SOL 109-99-9 THF
CON SUBSTAGE(1) room temperature, pH 7
SUBSTAGE(2) room temperature -> 0 deg C






STAGE(3)
RGT J 538-75-0 DCC
CON SUBSTAGE(1) 1 hour, 0 deg C
SUBSTAGE(2) overnight, room temperature

PRO P 503440-98-0
NTE intermediate product could be isolated

RX(16) RCT P 503440-98-0

RGT AP 64-19-7 AcOH, AQ 7440-66-6 Zn
PRO T 854719-15-8
SOL 64-19-7 AcOH, 7732-18-5 Water
CON 1 hour, room temperature

RX(59) OF 77 COMPOSED OF RX(14), RX(1), RX(2), RX(3), RX(4)
RX(59) AN + AO + B + F + M + O ==>
P



RX(14) RCT AN 3655-05-8, AO 107-96-0
PRO A 133367-03-0
NTE literature prepn.

RX(1) RCT A 133367-03-0, B 70-11-1
RGT D 7087-68-5 EtN(Pr-1)2
PRO C 503440-95-7
SOL 68-12-2 DMF
CON overnight, room temperature

RX(2) RCT C 503440-95-7

STAGE(1)
RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 15 minutes, room temperature

STAGE(2)
RCT F 15761-38-3
RGT I 2592-95-2 1-Benzotriazolol, D 7087-68-5 EtN(Pr-1)2
SOL 109-99-9 THF
CON SUBSTAGE(1) room temperature, pH 7
SUBSTAGE(2) room temperature -> 0 deg C

STAGE(3)
RGT J 538-75-0 DCC
CON SUBSTAGE(1) 1 hour, 0 deg C
SUBSTAGE(2) overnight, room temperature

PRO G 503440-96-8
NTE intermediate product could be isolated

RX(3) RCT G 503440-96-8

STAGE(1)
RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 20 minutes, room temperature

STAGE(2)
RCT M 3350-19-4
RGT D 7087-68-5 EtN(Pr-1)2
SOL 68-12-2 DMF
CON 3 hours, room temperature, pH 8 - 9

PRO N 503440-97-9
NTE intermediate product could be isolated

RX(4) RCT N 503440-97-9



STAGE(1)
RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 15 minutes, room temperature



STAGE(2)
RCT O 132149-57-6
RGT I 2592-95-2 1-Benzotriazolol, D 7087-68-5 EtN(Pr-1)2
SOL 109-99-9 THF
CON SUBSTAGE(1) room temperature, pH 7
SUBSTAGE(2) room temperature -> 0 deg C

STAGE(3)
RGT J 538-75-0 DCC
CON SUBSTAGE(1) 1 hour, 0 deg C
SUBSTAGE(2) overnight, room temperature


PRO P 503440-98-0
NTE intermediate product could be isolated

RX(61) OF 77 COMPOSED OF RX(14), RX(1), RX(2), RX(3), RX(4), RX(16)
RX(61) AN + AO + B + F + M + O ==>
T





6
STEPS



T
YIELD 83%

RX(14) RCT AN 3655-05-8, AO 107-96-0
PRO A 133367-03-0
NTE literature prepn.

RX(1) RCT A 133367-03-0, B 70-11-1
RGT D 7087-68-5 EtN(Pr-1)2
PRO C 503440-95-7
SOL 68-12-2 DMF
CON overnight, room temperature

RX(2) RCT C 503440-95-7

STAGE(1)
RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 15 minutes, room temperature

STAGE(2)
RCT F 15761-38-3

CORPORATE SOURCE: Department of Pharmacology and Molecular Sciences, The
Johns Hopkins University School of Medicine,
Baltimore, MD, 21205, USA

SOURCE: Journal of the American Chemical Society (2003),
125(52), 16172-16173
CODEN: JACSAT; ISSN: 0002-7863


PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English

AB Protein kinases often show low affinity for their protein substrates, which makes it difficult to study kinase-substrate interactions. Here, the authors show using expressed protein ligation with the signaling protein Src that it is feasible to install a covalently linked ATP moiety into the tail of Src, generating a semisynthetic protein with a high affinity for its cognate tyrosine kinase, Csk. It is also established that this Src-ATP conjugate can be used to selectively pull down Csk from a complex protein mixture. This work outlines a general strategy for identifying an unknown kinase that is responsible for the phosphorylation of a protein substrate on a site of interest.


REFERENCE COUNT: 23 THERE ARE 23 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

RX(2) OF 3 ...J + S ==> T

PAGE 1-A



PAGE 2-A



J

RGT I 2592-95-2 1-Benzotriazolol, D 7087-68-5 EtN(Pr-1)2
SOL 109-99-9 THF
CON SUBSTAGE(1) room temperature, pH 7
SUBSTAGE(2) room temperature -> 0 deg C

STAGE(3)
RGT J 538-75-0 DCC
CON SUBSTAGE(1) 1 hour, 0 deg C
SUBSTAGE(2) overnight, room temperature

PRO G 503440-96-8
NTE intermediate product could be isolated

RX(3) RCT G 503440-96-8

STAGE(1)
RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 20 minutes, room temperature

STAGE(2)
RCT M 3350-19-4
RGT D 7087-68-5 EtN(Pr-1)2
SOL 68-12-2 DMF
CON 3 hours, room temperature, pH 8 - 9

PRO N 503440-97-9
NTE intermediate product could be isolated

RX(4) RCT N 503440-97-9

STAGE(1)
RGT H 7647-01-0 HCl
SOL 141-78-6 AcOEt
CON 15 minutes, room temperature


STAGE(2)
RCT O 132149-57-6
RGT I 2592-95-2 1-Benzotriazolol, D 7087-68-5 EtN(Pr-1)2
SOL 109-99-9 THF
CON SUBSTAGE(1) room temperature, pH 7
SUBSTAGE(2) room temperature -> 0 deg C

STAGE(3)
RGT J 538-75-0 DCC
CON SUBSTAGE(1) 1 hour, 0 deg C
SUBSTAGE(2) overnight, room temperature

PRO P 503440-98-0
NTE intermediate product could be isolated


RX(16) RCT P 503440-98-0
RGT AP 64-19-7 AcOH, AQ 7440-66-6 Zn
PRO T 854749-15-8
SOL 64-19-7 AcOH, 7732-18-5 Water
CON 1 hour, room temperature

L6 ANSWER 3 OF 28 CASREACT COPYRIGHT 2007 ACS ON STN
ACCESSION NUMBER: 140:89809 CASREACT [Full-text](#)
TITLE: Conversion of a Tyrosine Kinase Protein Substrate to a
High Affinity Ligand by ATP Linkage
AUTHOR(S): Shen, Kui; Cole, Philip A.



(2)

PAGE 1-A



* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *


RX(2) RCT J 643760-37-6, S 35094-46-3

STAGE(1)
SOL 7732-18-5 Water
CON overnight, room temperature, pH 7

STAGE(2)
RGT U 3483-12-3 Cleland's reagent
SOL 7732-18-5 Water
CON 3 hours, room temperature, pH 7

PRO T 643760-38-7
NTE Tris buffered soln. both stages, tris(2-carboxyethyl)phosphine alternately used in place of dithiothreitol


L6 ANSWER 4 OF 28 CASREACT COPYRIGHT 2007 ACS ON STN
ACCESSION NUMBER: 140:59928 CASREACT [Full-text](#)
TITLE: Methods to initiate synthetic re-structuring of
peptides
AUTHOR(S): Wei, Qi; Harran, Susan; Harran, Patrick G.
CORPORATE SOURCE: Department of Biochemistry, University of Texas
Southwestern Medical Center at Dallas, Dallas, TX,
75390-9038, USA
SOURCE: Tetrahedron (2003), 59(45), 8947-8954
CODEN: TETRAH; ISSN: 0040-4020
PUBLISHER: Elsevier Science B.V.
DOCUMENT TYPE: Journal
LANGUAGE: English




AB The authors present a protocol for the synthesis of macrocyclic peptide ethers via a multi-component condensation reaction followed by metal-catalyzed cycloetherification. For example, macrocycle I was obtained in two steps from the three-component condensation of reactants N-Gly-Tyr-NHBu, allyl carbonate II and isonitrile 4-FC₆H₄CH(N.tpbond.C)SO₂C₆H₄Me-4, followed by cyclization of the adduct in presence of catalysts [(η³-allyl)PdCl]₂ and van Leeuwen's Xantphos. The authors are currently studying the application of the above protocol to solid-phase synthesis.

REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


RX(24) OF 170 ...BM + B + F ==> R...



BM



B



F

(24)

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(24) RCT BM 639492-47-0, B 639491-67-1

STAGE(1)

RGT H 584-06-7 K₂CO₃

SOL 68-12-2 DMF

CON 5 hours, room temperature

STAGE(2)

RCT F 165806-95-1


CON 17 hours, room temperature

PRO R 692740-58-2


NTE mol. sieve used in first stage

RX(35) OF 170 COMPOSED OF RX(1), RX(24)


RX(35) A + BM + F ==> R



A



BM



F

2
STEPS

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(1) RCT A 692730-70-4
RGT C 110-86-1 Pyridine, D 87413-09-0 Martin's reagent
PRO B 639491-67-1
CON SUBSTAGE(1) 1 hour, 4 deg C
SUBSTAGE(2) 2 hours, room temperature

RX(24) RCT BM 639492-47-0, B 639491-67-1

STAGE(1)

RGT H 584-08-7 K₂CO₃

SOL 68-12-2 DMF

CON 5 hours, room temperature

STAGE(2)

RCT F 165806-95-1


CON 17 hours, room temperature

PRO R 692740-58-2


NTE mol. sieve used in first stage

RX(96) OF 170 COMPOSED OF RX(20), RX(1), RX(24)


RX(96) BC + BM + F ==> R



BC



BM



F

3
STEPS

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(20) RCT BC 692730-27-1
RGT C 110-86-1 Pyridine, BE 32001-55-1 4-MeOC₆H₄CPh₂Cl
PRO A 692730-70-4
SOL 109-99-9 THF
CON 4 hours, room temperature

RX(1) RCT A 692730-70-4
RGT C 110-86-1 Pyridine, D 87413-09-0 Martin's reagent
PRO B 639491-67-1
CON SUBSTAGE(1) 1 hour, 4 deg C
SUBSTAGE(2) 2 hours, room temperature

RX(24) RCT BM 639492-47-0, B 639491-67-1

STAGE(1)

RGT H 584-08-7 K₂CO₃

SOL 68-12-2 DMF

CON 5 hours, room temperature

STAGE(2)

RCT F 165806-95-1


CON 17 hours, room temperature

PRO R 692740-58-2


NTE mol. sieve used in first stage

RX(107) OF 170 COMPOSED OF RX(19), RX(20), RX(1), RX(24)


RX(107) AY + BA + BD + BM + F ==> R




AY



BA



BB



4
STEPS

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(19) RCT AY 692729-77-4, BA 1826-67-1

STAGE(1)

SOL 109-99-9 THF
CON SUBSTAGE(1) room temperature -> -70 deg C
SUBSTAGE(2) 30 minutes, -70 deg C
SUBSTAGE(3) 45 minutes, -70 deg C -> room temperature

STAGE(2)

RCT BB 24424-99-5
CAT 1122-58-3 4-DMAP
SOL 75-09-2 CH₂Cl₂
CON SUBSTAGE(1) 20 minutes, room temperature
SUBSTAGE(2) 3.5 hours, room temperature

PRO BC 692730-27-1

RX(20) RCT BC 692730-27-1
RGT C 110-86-1 Pyridine, BE 32001-55-1 4-MeOC₆H₄CPH₂Cl
PRO A 692730-70-4
SOL 109-99-9 THF
CON 4 hours, room temperature

RX(1) RCT A 692730-70-4
RGT C 110-86-1 Pyridine, D 87413-09-0 Martin's reagent
PRO B 639491-67-1
CON SUBSTAGE(1) 1 hour, 4 deg C
SUBSTAGE(2) 2 hours, room temperature

RX(24) RCT BM 639492-47-0, B 639491-67-1

STAGE(1)




RGT H 584-08-7 K₂CO₃
SOL 68-12-2 DMF
CON 5 hours, room temperature

STAGE(2)

RCT F 165806-95-1
CON 17 hours, room temperature

PRO R 692740-58-2
NTE mol. sieve used in first stage

RX(139) OF 170 COMPOSED OF RX(18), RX(19), RX(20), RX(1), RX(24)
RX(139) AW + AX + BA + BB + BM + F ==>
R



* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(18) RCT AW 100-83-4, AX 89031-83-4
RGT AZ 534-17-8 Cs₂CO₃
PRO AY 692729-77-4
SOL 68-12-2 DMF
CON SUBSTAGE(1) room temperature -> 115 deg C
SUBSTAGE(2) 4 hours, 115 deg C

SUBSTAGE(3) 1.5 hours, 115 deg C

RX(19) RCT AY 692729-77-4, BA 1826-67-1

STAGE(1)

SOL 109-99-9 THF
CON SUBSTAGE(1) room temperature -> -70 deg C
SUBSTAGE(2) 30 minutes, -70 deg C
SUBSTAGE(3) 45 minutes, -70 deg C -> room temperature

STAGE(2)

RCT BB 24424-99-5
CAT 1122-58-3 4-DMAP
SOL 75-09-2 CH₂Cl₂
CON SUBSTAGE(1) 20 minutes, room temperature
SUBSTAGE(2) 3.5 hours, room temperature

PRO BC 692730-27-1

RX(20) RCT BC 692730-27-1
RGT C 110-86-1 Pyridine, BE 32001-55-1 4-MeOC₆H₄CPH₂Cl
PRO A 692730-70-4
SOL 109-99-9 THF
CON 4 hours, room temperature

RX(1) RCT A 692730-70-4
RGT C 110-86-1 Pyridine, D 87413-09-0 Martin's reagent
PRO B 639491-67-1
CON SUBSTAGE(1) 1 hour, 4 deg C
SUBSTAGE(2) 2 hours, room temperature

RX(24) RCT BM 639492-47-0, B 639491-67-1

STAGE(1)

RGT H 584-08-7 K₂CO₃
SOL 68-12-2 DMF
CON 5 hours, room temperature

STAGE(2)

RCT F 165806-95-1
CON 17 hours, room temperature

PRO R 692740-58-2

NTE mol. sieve used in first stage

L6 ANSWER 5 OF 28 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 140:59920 CASREACT Full-text
TITLE: Synthesis of a functionalized high affinity mannose receptor ligand and its application in the construction of peptide-, polyamide- and PNA-conjugates

AUTHOR(S): Kinzel, Olaf; Fattori, Daniela; Ingallinella, Paolo; Bianchi, Elisabetta; Pessi, Antonello
CORPORATE SOURCE: Department of Molecular and Cell Biology, IRBM P. Angeletti, Pomezia, 00040, Italy
SOURCE: Journal of Peptide Science (2003), 9(6), 375-385
CODEN: JPSIEI; ISSN: 1075-2617
PUBLISHER: John Wiley & Sons Ltd.
DOCUMENT TYPE: Journal
LANGUAGE: English

AB The synthesis of a high affinity mannose receptor ligand, appropriately functionalized for chemoselective ligation with an antigen or DNA-binding moieties is described. By a


combination of solid- and solution-phase chemical a versatile synthesis of the target structure was accomplished. Examples of subsequent ligation reactions are described.
REFERENCE COUNT: 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

RX(8) OF 73 AG ==> AH...


* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *


PAGE 1-C




PAGE 1-D



PAGE 2-A












AG

(8) →

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

BN
resin-boundAH
YIELD 58%

RX(8) RCT AG 635708-52-0

STAGE(1)

RGT AI 7646-85-7 ZnCl2, AJ 7558-79-4 Na2HPO4, AK 63-68-3
L-Methionine, AL 7790-28-5 NaIO4
SOL 7732-18-5 Water
CON 15 minutes, room temperature

STAGE(2)

RGT O 76-05-1 P3CCO2H

PRO AH 635708-53-1

L6 ANSWER 6 OF 28 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 139:214703 CASREACT Full-text

TITLE: Synthesis and hydrolysis studies of a peptide
containing the reactive triad of serine proteases with
an associated linker to a dye on a solid phase support

AUTHOR(S): Clough, John M.; Jones, Ray V. H.; McCann, Hannah;
Morris, David J.; Wills, Martin

CORPORATE SOURCE: Syngenta, Jealott's Hill Research Centre, Berkshire,
RG42 6EY, UK

SOURCE: Organic & Biomolecular Chemistry (2003), 1(9),
1486-1497

CODEN: OBCRAK; ISSN: 1477-0520

PUBLISHER: Royal Society of Chemistry


DOCUMENT TYPE: Journal

LANGUAGE: English

AB The synthesis of a Tentagel-supported peptide incorporating the reactive triad of serine,
histidine and aspartic acid, found within serine protease enzymes, is described.

REFERENCE COUNT: 20 THERE ARE 20 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


RX(35) OF 256 ...BN + AK ==> BC




AK

(35) →


* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *




PAGE 1-C



PAGE 1-D



PAGE 2-A



RX(35) RCT BN 590402-32-7D, AK 590402-17-8

STAGE(1)

ROT BP 128625-52-5 Benzotriazolol P der, BQ 7087-68-5
EtN(Pr-i)2, BR 2592-95-2 1-Benzotriazolol
SOL 68-12-2 DMF
CON 3 hours, room temperature

STAGE(2)


ROT AV 76-05-1 F3CCO2H, AM 6485-79-6 Silane,
tris(1-methylethyl)-
CON 2 hours, room temperature

PRO BO 590402-41-8D

NTE solid-supported reaction, first stage attachment to resin


RX(36) OF 256 ...BN + AL ==> BS

PAGE 1-B




(36)


PAGE 1-A




PAGE 1-B




PAGE 1-A



PAGE 1-B




PAGE 1-C



* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 1-C



* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(36) RCT BN 590402-32-7D, AL 590402-19-0

STAGE(1)

ROT BP 128625-52-5 Benzotriazolol P der, BQ 7087-68-5
EtN(Pr-i)2, BR 2592-95-2 1-Benzotriazolol
SOL 68-12-2 DMF
CON 3 hours, room temperature

STAGE(2)


ROT AV 76-05-1 F3CCO2H, AM 6485-79-6 Silane,
tris(1-methylethyl)-
CON 2 hours, room temperature

PRO BS 590402-42-9D


NTE solid-supported reaction, first stage attachment to resin

RX(37) OF 256 ...BN + AM ==> BT


PAGE 1-A



PAGE 1-B




PAGE 1-C



* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 2-A




PAGE 2-B



PAGE 1-E



PAGE 2-A



RX(37) RCT BN 590402-32-7D, AM 590402-21-4

STAGE(1)

RGT BP 128625-52-5 Benzotriazolol P der, BQ 7087-68-5
EtN(Px-1)2, BR 2592-95-2 1-Benzotriazolol
SOL 68-12-2 DMF
CON 3 hours, room temperature

STAGE(2)


RGT AV 76-05-1 P3CCO2H, AM 6485-79-6 Silane,
tris(1-methylethyl)-
CON 2 hours, room temperature

PRO BT 590402-43-0D

NTE solid-supported reaction, first stage attachment to resin

RX(38) OF 256 ...BN + AN ==> BU


PAGE 1-A




(37)

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *


PAGE 1-B




PAGE 1-C




PAGE 1-D



PAGE 1-B




PAGE 1-C




* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 1-B




(38)


PAGE 1-A




PAGE 1-B




PAGE 1-C



PAGE 1-B




PAGE 1-C




* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 2-A



PAGE 1-D



* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(38) RCT BN 590402-32-7D, AN 590402-23-6

STAGE(1)

RGT BP 128625-52-5 Benzotriazolol P der, BQ 7087-68-5

EtN(Pr-i)2, BR 2592-95-2 1-Benzotriazolol

SOL 68-12-2 DMF

CON 3 hours, room temperature

STAGE(2)

RGT AV 76-05-1 P3CCO2H, AM 6485-79-6 Silane,

tris(1-methylethyl)-


CON 2 hours, room temperature

PRO BU 590402-44-1D


NTR solid-supported reaction, first stage attachment to resin

RX(39) OF 256 ...BN + AO ==> BV

PAGE 1-A




PAGE 2-B




(39)

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *


PAGE 1-B




PAGE 1-C




PAGE 1-D



PAGE 1-E



PAGE 2-A



RX(39) RCT BN 590402-32-7D, AO 590402-25-8

STAGE(1)

RT BP 126625-52-5 Benzotriazolol P der, BQ 7087-66-5
EtN(Pr-1)2, BR 2592-95-2 1-Benzotriazolol
SOL 68-12-2 DMF
CON 3 hours, room temperature

STAGE(2)


RT AV 76-05-1 P3CCO2H, AM 6485-79-6 Silane,
tris(1-methylethyl)-
CON 2 hours, room temperature

PRO BV 590402-45-2D


NTE solid-supported reaction, first stage attachment to resin

RX(40) OF 256 ...BN + AP ==> BW


PAGE 1-A



PAGE 1-B




PAGE 1-C

BN
resin-bound

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 2-A




AP


(40)

→


PAGE 1-A




PAGE 1-B



PAGE 1-C



PAGE 1-D

BN
resin-bound

RX(40) RCT BN 590402-32-7D, AP 590402-26-9

STAGE(1)

RT BP 126625-52-5 Benzotriazolol P der, BQ 7087-66-5
EtN(Pr-1)2, BR 2592-95-2 1-Benzotriazolol
SOL 68-12-2 DMF
CON 3 hours, room temperature

STAGE(2)


RT AV 76-05-1 P3CCO2H, AM 6485-79-6 Silane,
tris(1-methylethyl)-
CON 2 hours, room temperature

PRO BW 590402-46-3D


NTE solid-supported reaction, first stage attachment to resin

RX(41) OF 256 ...BN + AQ ==> BX


PAGE 1-A




PAGE 1-B




PAGE 1-C



PAGE 1-C



PAGE 1-D



RX(41) RCT BN 590402-32-7D, AQ 590402-27-0

STAGE(1)

ROT BP 128625-52-5 Benzotriazolol P der, BQ 7087-68-5
EtN(Pr-1)2, BR 2592-95-2 1-Benzotriazolol
SOL 68-12-2 DMF
CON 3 hours, room temperature

STAGE(2)


ROT AV 76-05-1 F3CCO2H, AM 6485-79-6 Silane,
tri(1-methylethyl)-
CON 2 hours, room temperature

PRO BX 590402-47-4D


NTE solid-supported reaction, first stage attachment to resin

RX(42) OF 256 ...BN + AR ==> BY

PAGE 1-A




PAGE 1-B




(41)

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *


PAGE 1-B




PAGE 1-A



PAGE 1-B



PAGE 1-C



* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *


PAGE 1-B

OBu-t


PAGE 1-D

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *



(42)



PAGE 1-B



PAGE 1-C



<-----User Break----->

=>

10/561,754

151 / 447

Robert Havlin

=> file reg
 COST IN U.S. DOLLARS
 FULL ESTIMATED COST
 DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)
 CA SUBSCRIBER PRICE

SINCE FILE ENTRY	TOTAL SESSION
5.52	130.06
0.00	-0.73

FILE 'REGISTRY' ENTERED AT 06:28:44 ON 30 MAY 2007
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2007 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file
 provided by InfoChem.

STRUCTURE FILE UPDATES: 28 MAY 2007 HIGHEST RN 935999-19-2
 DICTIONARY FILE UPDATES: 28 MAY 2007 HIGHEST RN 935999-19-2

New CAS Information Use Policies, enter HELP USAGETERMS for details.


TSCA INFORMATION NOW CURRENT THROUGH December 2, 2006

Please note that search-term pricing does apply when
 conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and
 predicted properties as well as tags indicating availability of
 experimental property data in the original document. For information
 on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stndoc/properties.html>

=>
 Uploading C:\Program Files\Stnexp\Queries\10.561754\clm14 crop1 plus.etr



chain nodes :
 1 2 3 12 13 14 15 16 22 23 24 25 26 27 28

10/561,754

152 / 447

Robert Havlin

ring nodes :
 4 5 6 7 8 17 18 19 20 21
 chain bonds :
 1-3 1-2 1-25 4-12 12-13 12-14 14-26 15-18 15-16 15-28 17-22 22-23 22-24
 24-27
 ring bonds :
 4-5 4-8 5-6 6-7 7-8 17-18 17-21 18-19 19-20 20-21
 exact/norm bonds :
 1-3 1-2 1-25 4-5 4-8 4-12 5-6 6-7 7-8 12-13 12-14 14-26 15-18 15-16
 15-28 17-18 17-21 17-22 18-19 19-20 20-21 22-23 22-24 24-27

G1:C,S

Match level :
 1:CLASS 2:CLASS 3:CLASS 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 12:CLASS 13:CLASS
 14:CLASS 15:CLASS 16:CLASS 17:Atom 18:Atom 19:Atom 20:Atom 21:Atom 22:CLASS 23:CLASS
 24:CLASS 25:CLASS 26:Atom 27:Atom 28:CLASS
 fragments assigned product role:
 containing 15
 fragments assigned reactant/reagent role:
 containing 1
 containing 4

L7 STRUCTURE UPLOADED

=> d
 L7 HAS NO ANSWERS
 L7 STR

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

Structure attributes must be viewed using STN Express query preparation.

=> file casreact
 COST IN U.S. DOLLARS
 FULL ESTIMATED COST
 DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)
 CA SUBSCRIBER PRICE

SINCE FILE ENTRY	TOTAL SESSION
0.45	130.51
0.00	-0.73

FILE 'CASREACT' ENTERED AT 08:29:13 ON 30 MAY 2007
 USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
 COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is
 held by the publishers listed in the PUBLISHER (PB) field (available
 for records published or updated in Chemical Abstracts after December
 26, 1996), unless otherwise indicated in the original publications.

FILE CONTENT:1840 - 27 May 2007 VOL 146 ISS 23

New CAS Information Use Policies, enter HELP USAGETERMS for details.

 * CASREACT now has more than 12 million reactions *

Some CASREACT records are derived from the ZIC/VINITI database (1974-1999) provided by InfoChem, INPI data prior to 1986, and Biotransformations database compiled under the direction of Professor Dr. Klaus Kieselich.

This file contains CAS Registry Numbers for easy and accurate substance identification.

==> # 17
SAMPLE SEARCH INITIATED 08:29:18 FILE 'CASREACT'
SCREENING COMPLETE - 1444 REACTIONS TO VERIFY FROM 71 DOCUMENTS
100.0% DONE 1444 VERIFIED 20 HIT RXNS 5 DOCS
SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
BATCH **COMPLETE**
PROJECTED VERIFICATIONS: 26603 TO 31157
PROJECTED ANSWERS: 5 TO 234

L8 5 SEA SSS SAM L7 (20 REACTIONS)


==> # 18 not py > 2003
101431 PY > 2003
L9 3 L8 NOT PY > 2003

==> d ibib abs fhit


L9 ANSWER 1 OF 3 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 138:385710 CASREACT Full-text
TITLE: Synthesis of N-Pmoc 3-[4-(di-tert-butylphosphonomethyl)phenyl]pipercolic acid as a conformationally constrained phosphotyrosyl mimetic suitably protected for peptide synthesis
AUTHOR(S): Liu, Ding-Quo; Wang, Xiang-Zhu; Gao, Yang; Li, Bihua; Yang, Dajun; Burke, Terrence R.
CORPORATE SOURCE: Center for Cancer Research, Laboratory of Medicinal Chemistry, MCI at Frederick, Frederick, MD, 21702, USA
SOURCE: Tetrahedron (2002), 58(52), 10423-10428
CODEN: TETRA; ISSN: 0040-4020
PUBLISHER: Elsevier Science Ltd.
DOCUMENT TYPE: Journal
LANGUAGE: English

AB Phosphonomethylphenylalanine (Pmp) has shown wide utility as a hydrolytically stable phosphotyrosyl mimetic, particularly in Src homol. 2 (SH2) domain-binding peptides. (2S,3R)-3-[4-(phosphonomethyl)phenyl]pipercolic acid (3) represents a variant of Pmp having ϕ and χ_1 torsion angles constrained through incorporation into the piperidinyl ring structure contained within pipercolic acid. Reported here is the enantioselective preparation of 3, in an orthogonally protected form (title compound, 4) suitable for use in peptide synthesis. Stereochemistries at both the 2- and 3-positions are derived inductively from a single chiral center provided by the com. available Evans chiral auxiliary, (4S)-4-benzyl-1,3-oxazolidin-2-one. Incorporation of 4 into a Grb2 SH2 domain-directed tripeptide showed that Grb2 SH2 domain-binding affinity was reduced relative to the parent Pmp-containing tripeptide. Although conformational constraint did not enhance affinity in this case, novel amino acid analog 4 may serve as a useful tool for the induction of defined phosphotyrosyl geometry in peptides directed at other signal transduction targets.

RX(9) OF 55 ...J + AP ==> AQ...




J



AP

(9)



80
YIELD 77%

RX(9) RCT J 525575-16-2

STAGE(1)
RGT AR 110-89-4 Piperidine
SOL 75-05-8 MeCN
CON 4 hours, room temperature

STAGE(2)
RCT AP 39061-59-1
RGT AS 7087-68-5 EtN(Pr-1)2
SOL 68-12-2 DMF
CON overnight, room temperature


PRO AQ 525575-19-3
REFERENCE COUNT: 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

==> d ibib abs fhit 2-
YOU HAVE REQUESTED DATA FROM 2 ANSWERS - CONTINUE? Y/(N):y

L9 ANSWER 2 OF 3 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 137:247666 CASREACT Full-text
TITLE: Bicyclic piperazinylbenzenesulphonamides are potent and selective 5-HT6 receptor antagonists
AUTHOR(S): Bromidge, Steven M.; Clarke, Stephen E.; King, Frank D.; Lovell, Peter J.; Newman, Helen; Riley, Graham; Routledge, Carol; Serafinowska, Halina T.; Smith, Douglas R.; Thomas, David R.
CORPORATE SOURCE: Department of Psychiatry, GlaxoSmithKline, Essex, Harlow, CM19 5AW, UK
SOURCE: Bioorganic & Medicinal Chemistry Letters (2002), 12(10), 1357-1360
CODEN: BMCL; ISSN: 0960-894X
PUBLISHER: Elsevier Science Ltd.
DOCUMENT TYPE: Journal
LANGUAGE: English

AB The synthesis of novel 3-(octahydropyrido[1,2-a]pyrazin-2-yl)- and 3-(hexahydropyrido[1,2-a]pyrazin-2-yl)phenyl-2-benzothiothiophene sulfonamide deriva. is described. The compds. show high affinity for the 5-HT6 receptor, excellent selectivity against a range of other receptors, and good brain penetration.

RX(10) OF 33 ...AE + Y ==> AH...




RX(10) RCT AE 239122-42-0
STAGE(1)
RGT AI 76-05-1 F3CCO2H
SOL 75-09-2 CH2Cl2

STAGE(2)
RCT Y 598-21-0
RGT D 7087-68-5 EtN(Pr-1)2
SOL 75-09-2 CH2Cl2


PRO AH 239122-44-3
REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 3 OF 3 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 126:118191 CASREACT Full-text
TITLE: Total synthesis of cyclothialidine
AUTHOR(S): Goetschi, Erwin; Janny, Christian Johannes; Reindl, Peter; Ricklin, Fabienne
CORPORATE SOURCE: Pharma Division, Hoffmann-La Roche Ltd., Basel, CH-4002, Switz.
SOURCE: Helvetica Chimica Acta (1996), 79(8), 2219-2234
CODEN: HCACAV; ISSN: 0018-019X
PUBLISHER: Verlag Helvetica Chimica Acta
DOCUMENT TYPE: Journal
LANGUAGE: English
OI



AB A total synthesis of cyclothialidine I, a DNA gyrase inhibitor isolated from *Streptomyces filipinensis*, is described. The synthetic concept was tested by preparing a lactone containing the bicyclic core entity of I. Key features of the synthesis of I are preparation of 3,5-dihydroxy-2,6-dimethylbenzoate from 3,5-dihydroxybenzoate by 2 consecutive Mannich aminomethylation/hydrogenation sequences, benzylic N-bromosuccinimide bromination of an ester derivative thereof and its subsequent coupling with Boc-Ser-Cys-OMe, cyclization of the α -hydroxy acid II (R = OH, R1 = H) to the 12-membered lactone II (RR1 = bond) using preferably Mitsunobu conditions, and completion of the peptidic side chains of I using Boc strategy. Optically pure cis-N-(tert-butoxycarbonyl)-3-hydroxy-L-proline was obtained by resolution of the racemate via an efficient reaction sequence containing a lipase-catalyzed enantiospecific acetate hydrolysis. The structure of natural I was confirmed by comparison with the synthetic material. The synthetic route described provides also easy access to analogs of I.

RX(5) OF 433 ... L + P ==> Q...



(5)

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 2-A

YIELD 73%

RX(5) RCT L 3262-72-4, P 186132-90-1
 ROT R 25952-53-8 EDAP
 PRO Q 186132-91-2
 SOL 75-05-8 MeCN

=>
=> d hist

(FILE 'HOME' ENTERED AT 08:18:24 ON 30 MAY 2007)


L1 FILE 'REGISTRY' ENTERED AT 08:18:37 ON 30 MAY 2007
 STRUCTURE UPLOADED

L2 FILE 'CASREACT' ENTERED AT 08:19:13 ON 30 MAY 2007
 0 S L1 SSS SAM
 L3 1 S L1 SSS FULL


FILE 'REGISTRY' ENTERED AT 08:21:37 ON 30 MAY 2007

halogen] or their pharmaceutically-acceptable salts or esters, including racemates, diastereoisomers and optical isomers, which are inhibitors of the hepatitis C virus (HCV). Thus, tripeptide II was prepared by a multistep synthesis involving etherification of tripeptide prolinol derivative and cyclization of 3-(bromoacetyl)quinoline derivative with tert-butylacetylthiourea as key steps. Compound II is extremely active against the HCV NS3 protease on the basis of enzymic and cellular assays.

RX(11) OF 55 COMPOSED OF RX(2), RX(4)
 RX(11) C + L ==> P




C



L

2 STEPS

P
YIELD 76%

L4 STRUCTURE UPLOADED

L5 FILE 'CASREACT' ENTERED AT 08:21:56 ON 30 MAY 2007
 50 S L4

L6 FILE 'CASREACT' ENTERED AT 08:24:06 ON 30 MAY 2007
 28 S L5 NOT PY>2003

L7 FILE 'REGISTRY' ENTERED AT 08:28:44 ON 30 MAY 2007
 STRUCTURE UPLOADED

L8 FILE 'CASREACT' ENTERED AT 08:29:13 ON 30 MAY 2007
 5 S L7
 L9 3 S L8 NOT PY > 2003

=> # 17 *** full
 FULL SEARCH INITIATED 08:47:17 FILE 'CASREACT'
 SCREENING COMPLETE - 30569 REACTIONS TO VERIFY FROM 1466 DOCUMENTS

100.0% DONE - 30569 VERIFIED 455 HIT RXNS 46 DOCS
 SEARCH TIME: 00.00.04

L10 46 SEA SSS FUL L7 (455 REACTIONS)

=> # 110 not py >2003
 101431 PY >2003
 L11 21 L10 NOT PY >2003

=> # 111 not 19
 L12 18 L11 NOT L9

=> d ibib abs hit 1-10

L12 ANSWER 1 OF 18 CASREACT. COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 142:447413 CASREACT Full-text
 TITLE: Preparation of tripeptides as hepatitis C virus inhibitors
 PATENT ASSIGNEE(S): Boehringer Ingelheim Canada Ltd., Can.
 SOURCE: Can. Pat. Appl., 33 pp.
 CODEN: CPXXEB
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
CA 2370400	A1	20030801	CA 2002-2370400	20020201
PRIORITY APPLN. INFO.:				
OTHER SOURCE(S): MARPAT 142:447413				
GI				



* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The invention relates to tripeptides I [B is H, (un)substituted aryl, aralkyl, heterocyclyl, acyl, CO₂H or ester, a (thio)amide or sulfonyl group; Y is H or alkyl; R₃ is (un)substituted alkyl, cycloalkyl or alkylcycloalkyl; R₂ is CH₂R₂₀, NHR₂₀, OR₂₀ or SR₂₀, where R₂₀ is (un)substituted (un)saturated cycloalkyl, aryl, aralkyl, heterocyclyl, etc.; R₁ is H, (un)substituted alkyl, cycloalkyl, alkenyl or alkynyl optionally substituted by


RX(2) RCT C 791835-61-5
 PRO H 791835-62-6
 SOL 101-84-8 PhOPh
 CON SUBSTAGE(1) 5 minutes, 250 deg C -> 230 deg C
 SUBSTAGE(2) 7 minutes, 230 deg C -> 245 deg C
 SUBSTAGE(3) cooled
 SUBSTAGE(4) 0 deg C
 NTS extended heating at 250 degree celsius would give decarboxylation of desired ester, reaction is done in two batches

RX(4) RCT H 791835-62-6, L 801282-34-0
 ROT Q 534-17-8 Cs2CO3
 PRO P 851009-68-2
 SOL 872-50-4 NMMP
 CON SUBSTAGE(1) room temperature
 SUBSTAGE(2) 5 hours, 72 deg C

RX(18) OF 55 COMPOSED OF RX(1), RX(2), RX(4)
 RX(18) A + B + L ==> P


A
● 2 HCl

B



L

3 STEPS



P
YIELD 76%

RX(1) RCT A 3575-32-4

STAGE(1)

RGT D 144-55-8 NaHCO₃
SOL 7932-18-5 Water
CON neutralized

STAGE(2)

RCT B 762-42-5
RGT E 62-53-3 PhNH₂
SOL 67-56-1 MeOH
CON SUBSTAGE(1) 0 deg C, neutralized
SUBSTAGE(2) heated
SUBSTAGE(3) 2 hours, 65 deg C
SUBSTAGE(4) 14 hours, room temperature

PRO C 791835-61-5

NTE exothermic reaction in second stage, incremental addition of aniline in second stage

RX(2)

RCT C 791835-61-5
PRO H 791835-62-6
SOL 101-84-8 PhOPh
CON SUBSTAGE(1) 5 minutes, 250 deg C -> 230 deg C
SUBSTAGE(2) 7 minutes, 230 deg C -> 245 deg C
SUBSTAGE(3) cooled
SUBSTAGE(4) 0 deg C



NTE extended heating at 250 degree celsius would give decarboxylation of desired ester, reaction is done in two batches

RX(4)



RCT H 791835-62-6, L 801282-34-8
RGT Q 534-17-8 Cs₂CO₃
PRO P 851009-68-2
SOL 872-50-4 NMEP
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 5 hours, 72 deg C

RX(19) OF 55 COMPOSED OF REACTION SEQUENCE RX(3), RX(4)
AND REACTION SEQUENCE RX(2), RX(4)

...J + K ==> L...
...C + L ==> P




START NEXT REACTION SEQUENCE



L

2
STEPS



P
YIELD 76%

RX(3) RCT J 572924-77-7, K 98-58-8

STAGE(1)

CAT 1122-58-3 4-DMAP
SOL 75-09-2 CH₂Cl₂
CON room temperature -> 0 deg C

STAGE(2)

RGT M 121-44-8 Et₃N
CON SUBSTAGE(1) 3 minutes, 0 deg C
SUBSTAGE(2) 1 hour, 0 deg C
SUBSTAGE(3) 0 deg C -> room temperature
SUBSTAGE(4) 18 hours, room temperature

PRO L 801282-34-8

RX(2)


RCT C 791835-61-5
PRO H 791835-62-6
SOL 101-84-8 PhOPh
CON SUBSTAGE(1) 5 minutes, 250 deg C -> 230 deg C
SUBSTAGE(2) 7 minutes, 230 deg C -> 245 deg C
SUBSTAGE(3) cooled
SUBSTAGE(4) 0 deg C

NTE extended heating at 250 degree celsius would give decarboxylation of desired ester, reaction is done in two


batches

RX(4) RCT H 791835-62-6, L 801282-34-8
RGT Q 534-17-8 Cs₂CO₃
PRO P 851009-68-2
SOL 872-50-4 NMEP
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 5 hours, 72 deg C

RX(20) OF 55 COMPOSED OF RX(2), RX(4), RX(5)
RX(20) C + L ==> S



C



L

3
STEPS

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(2) RCT C 791835-61-5
PRO H 791835-62-6
SOL 101-84-8 PhOPh
CON SUBSTAGE(1) 5 minutes, 250 deg C -> 230 deg C
SUBSTAGE(2) 7 minutes, 230 deg C -> 245 deg C
SUBSTAGE(3) cooled
SUBSTAGE(4) 0 deg C



NTE extended heating at 250 degree celsius would give decarboxylation of desired ester, reaction is done in two

RX(4) RCT H 791835-62-6, L 801282-34-8
RGT Q 534-17-8 Cs₂CO₃

PRO P 851009-68-2
 SOL 872-50-4 NMEP
 CON SUBSTAGE(1) room temperature
 SUBSTAGE(2) 5 hours, 72 deg C

RX(5) RCT P 851009-68-2
 RGT T 1310-73-2 NaOH
 PRO S 851009-69-3
 SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF
 CON SUBSTAGE(1) room temperature
 SUBSTAGE(2) 1.5 hours, room temperature

RX(21) OF 55 COMPOSED OF REACTION SEQUENCE RX(3), RX(4)
 AND REACTION SEQUENCE RX(1), RX(2), RX(4)
 ...J + K ==> L...
 ...A + B + L ==> P



START NEXT REACTION SEQUENCE

STAGE(2)
 RGT M 121-44-8 Et3N
 CON SUBSTAGE(1) 3 minutes, 0 deg C
 SUBSTAGE(2) 1 hour, 0 deg C
 SUBSTAGE(3) 0 deg C -> room temperature
 SUBSTAGE(4) 18 hours, room temperature

PRO L 801282-34-8

RX(1) RCT A 3575-32-4

STAGE(1)
 RGT D 144-55-8 NaHCO3
 SOL 7732-18-5 Water
 CON neutralized





STAGE(2)
 RCT B 762-42-5
 RGT E 62-53-3 PhNH2
 SOL 67-56-1 MeOH
 CON SUBSTAGE(1) 0 deg C, neutralized
 SUBSTAGE(2) heated
 SUBSTAGE(3) 2 hours, 65 deg C
 SUBSTAGE(4) 14 hours, room temperature

PRO C 791835-61-5
 NTE exothermic reaction in second stage, incremental addition of aniline in second stage

RX(2) RCT C 791835-61-5
 PRO H 791835-62-6
 SOL 101-84-8 PhOPh
 CON SUBSTAGE(1) 5 minutes, 250 deg C -> 230 deg C
 SUBSTAGE(2) 7 minutes, 230 deg C -> 245 deg C
 SUBSTAGE(3) cooled
 SUBSTAGE(4) 0 deg C
 NTE extended heating at 250 degree celsius would give decarboxylation of desired ester, reaction is done in two batches

RX(4) RCT H 791835-62-6, L 801282-34-8
 RGT Q 534-17-8 Cs2CO3
 PRO P 851009-68-2
 SOL 872-50-4 NMEP
 CON SUBSTAGE(1) room temperature
 SUBSTAGE(2) 5 hours, 72 deg C

RX(22) OF 55 COMPOSED OF RX(1), RX(2), RX(4), RX(5)
 RX(22) A + B + L ==> B



YIELD 76%

RX(3) RCT J 572924-77-7, K 98-58-8

STAGE(1)
 CAT 1122-58-3 4-DMAP
 SOL 75-09-2 CH2Cl2
 CON room temperature -> 0 deg C

STAGE(2)
 RGT M 121-44-8 Et3N
 CON SUBSTAGE(1) 3 minutes, 0 deg C
 SUBSTAGE(2) 1 hour, 0 deg C
 SUBSTAGE(3) 0 deg C -> room temperature
 SUBSTAGE(4) 18 hours, room temperature

PRO L 801282-34-8

RX(1) RCT A 3575-32-4


STAGE(1)
 RGT D 144-55-8 NaHCO3
 SOL 7732-18-5 Water
 CON neutralized

STAGE(2)
 RCT B 762-42-5
 RGT E 62-53-3 PhNH2
 SOL 67-56-1 MeOH
 CON SUBSTAGE(1) 0 deg C, neutralized
 SUBSTAGE(2) heated
 SUBSTAGE(3) 2 hours, 65 deg C
 SUBSTAGE(4) 14 hours, room temperature

PRO C 791835-61-5
 NTE exothermic reaction in second stage, incremental addition of aniline in second stage

RX(2) RCT C 791835-61-5
 PRO H 791835-62-6
 SOL 101-84-8 PhOPh
 CON SUBSTAGE(1) 5 minutes, 250 deg C -> 230 deg C
 SUBSTAGE(2) 7 minutes, 230 deg C -> 245 deg C
 SUBSTAGE(3) cooled
 SUBSTAGE(4) 0 deg C
 NTE extended heating at 250 degree celsius would give decarboxylation of desired ester, reaction is done in two batches




RX(4) RCT H 791835-62-6, L 801282-34-8
 RGT Q 534-17-8 Cs2CO3
 PRO P 851009-68-2
 SOL 872-50-4 NMEP
 CON SUBSTAGE(1) room temperature
 SUBSTAGE(2) 5 hours, 72 deg C



* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(5) RCT P 851009-68-2
 RGT T 1310-73-2 NaOH
 PRO S 851009-69-3
 SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF
 CON SUBSTAGE(1) room temperature
 SUBSTAGE(2) 1.5 hours, room temperature

RX(25) OF 55 COMPOSED OF RX(2), RX(4), RX(5), RX(6)
 RX(25) C + L + V ==> W



RX(2) RCT C 791835-61-5
 PRO H 791835-62-6
 SOL 101-84-8 PhOPh
 CON SUBSTAGE(1) 5 minutes, 250 deg C -> 230 deg C
 SUBSTAGE(2) 7 minutes, 230 deg C -> 245 deg C
 SUBSTAGE(3) cooled
 SUBSTAGE(4) 0 deg C
 NTE extended heating at 250 degree celsius would give decarboxylation of desired ester, reaction is done in two batches

RX(4) RCT H 791835-62-6, L 801282-34-8
 RGT Q 534-17-8 Cs2CO3
 PRO P 851009-68-2
 SOL 872-50-4 NMEP
 CON SUBSTAGE(1) room temperature
 SUBSTAGE(2) 5 hours, 72 deg C

RX(5) RCT P 851009-68-2
 RGT T 1310-73-2 NaOH
 PRO S 851009-69-3
 SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF
 CON SUBSTAGE(1) room temperature
 SUBSTAGE(2) 1.5 hours, room temperature





RX(6) RCT S 851009-69-3

STAGE(1)
 RGT M 121-44-8 Et3N, X 543-27-1 ClCO2Bu-i
 SOL 109-99-9 THF
 CON SUBSTAGE(1) 0 deg C
 SUBSTAGE(2) 1 hour, 0 deg C

STAGE(2)
 RCT V 334-88-3
 SOL 60-29-7 Et2O
 CON SUBSTAGE(1) 1 minute, 0 deg C
 SUBSTAGE(2) 30 minutes, 0 deg C
 SUBSTAGE(3) 45 minutes, room temperature

PRO W 851009-70-6

RX(33) OF 55 COMPOSED OF RX(1), RX(2), RX(4), RX(5), RX(6)
 RX(33) A + B + L + V ==> W



RX(1) RCT A 3575-32-4

STAGE(1)
 RGT D 144-55-5 NaHCO3
 SOL 7732-18-5 Water
 CON neutralized

STAGE(2)
 RCT B 762-42-5
 RGT E 62-53-3 PhNH2
 SOL 67-56-1 MeOH
 CON SUBSTAGE(1) 0 deg C, neutralized
 SUBSTAGE(2) heated
 SUBSTAGE(3) 2 hours, 65 deg C
 SUBSTAGE(4) 14 hours, room temperature

PRO C 791835-61-5
 NTE exothermic reaction in second stage, incremental addition of aniline in second stage

RX(2) RCT C 791835-61-5
 PRO H 791835-62-6
 SOL 101-84-8 PhOPh

CON SUBSTAGE(1) 5 minutes, 250 deg C -> 230 deg C
 SUBSTAGE(2) 7 minutes, 230 deg C -> 245 deg C
 SUBSTAGE(3) cooled
 SUBSTAGE(4) 0 deg C
 NTE extended heating at 250 degree celsius would give decarboxylation of desired ester, reaction is done in two batches

RX(4) RCT H 791835-62-6, L 801282-34-8
 RGT Q 534-17-8 Cs2CO3
 PRO P 851009-68-2
 SOL 872-50-4 NMEP
 CON SUBSTAGE(1) room temperature
 SUBSTAGE(2) 5 hours, 72 deg C

RX(5) RCT P 851009-68-2
 RGT T 1310-73-2 NaOH
 PRO S 851009-69-3
 SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF
 CON SUBSTAGE(1) room temperature
 SUBSTAGE(2) 1.5 hours, room temperature



RX(6) RCT S 851009-69-3

STAGE(1)
 RGT M 121-44-8 Et3N, X 543-27-1 ClCO2Bu-i
 SOL 109-99-9 THF
 CON SUBSTAGE(1) 0 deg C
 SUBSTAGE(2) 1 hour, 0 deg C


STAGE(2)
 RCT V 334-88-3
 SOL 60-29-7 Et2O
 CON SUBSTAGE(1) 1 minute, 0 deg C
 SUBSTAGE(2) 30 minutes, 0 deg C
 SUBSTAGE(3) 45 minutes, room temperature

PRO W 851009-70-6



RX(34) OF 55 COMPOSED OF REACTION SEQUENCE RX(3), RX(4), RX(5)
 AND REACTION SEQUENCE RX(2), RX(4), RX(5)
 ...J + K ==> L...
 ... C + L ==> S



3
STEPS



START NEXT REACTION SEQUENCE

3
STEPS

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(3) RCT J 572924-77-7, K 98-58-8

STAGE(1)

CAT 1122-58-3 4-DMAP

SOL 75-09-2 CH₂Cl₂

CON room temperature -> 0 deg C

STAGE(2)

RGT M 121-44-8 Et₃N

CON SUBSTAGE(1) 3 minutes, 0 deg C

SUBSTAGE(2) 1 hour, 0 deg C

SUBSTAGE(3) 0 deg C -> room temperature

SUBSTAGE(4) 18 hours, room temperature

PRO L 801282-34-8

RX(2)

RCT C 791835-61-5

PRO H 791835-62-6

SOL 101-84-8 PhOPh

CON SUBSTAGE(1) 5 minutes, 250 deg C -> 230 deg C

SUBSTAGE(2) 7 minutes, 230 deg C -> 245 deg C

SUBSTAGE(3) cooled

SUBSTAGE(4) 0 deg C

NTS extended heating at 250 degree celsius would give decarboxylation of desired ester, reaction is done in two batches

RX(4)

RCT H 791835-62-6, L 801282-34-8

RGT Q 534-17-8 Cs₂CO₃

PRO P 851009-68-2

SOL 872-50-4 NMRP

CON SUBSTAGE(1) room temperature

SUBSTAGE(2) 5 hours, 72 deg C

RX(5)

RCT P 851009-68-2

RGT T 1310-73-2 NaOH

PRO S 851009-69-3

SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF

CON SUBSTAGE(1) room temperature



SUBSTAGE(2) 1.5 hours, room temperature

RX(35) OF 55 COMPOSED OF REACTION SEQUENCE RX(3), RX(4), RX(5), RX(6)




AND REACTION SEQUENCE RX(2), RX(4), RX(5), RX(6)

...J + K ==> L...

... C + L + V ==> W



START NEXT REACTION SEQUENCE



RX(3) RCT J 572924-77-7, K 98-58-8

STAGE(1)

CAT 1122-58-3 4-DMAP

SOL 75-09-2 CH₂Cl₂

CON room temperature -> 0 deg C

STAGE(2)

RGT M 121-44-8 Et₃N

CON SUBSTAGE(1) 3 minutes, 0 deg C

SUBSTAGE(2) 1 hour, 0 deg C

SUBSTAGE(3) 0 deg C -> room temperature

SUBSTAGE(4) 18 hours, room temperature

PRO L 801282-34-8

RX(2)

RCT C 791835-61-5

PRO H 791835-62-6

SOL 101-84-8 PhOPh

CON SUBSTAGE(1) 5 minutes, 250 deg C -> 230 deg C

SUBSTAGE(2) 7 minutes, 230 deg C -> 245 deg C

SUBSTAGE(3) cooled

SUBSTAGE(4) 0 deg C

NTS extended heating at 250 degree celsius would give decarboxylation of desired ester, reaction is done in two batches

RX(4)

RCT H 791835-62-6, L 801282-34-8

RGT Q 534-17-8 Cs₂CO₃

PRO P 851009-68-2

SOL 872-50-4 NMRP

CON SUBSTAGE(1) room temperature

SUBSTAGE(2) 5 hours, 72 deg C

RX(5)

RCT P 851009-68-2

RGT T 1310-73-2 NaOH

PRO S 851009-69-3

SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF

CON SUBSTAGE(1) room temperature

SUBSTAGE(2) 1.5 hours, room temperature

RX(6)

RCT S 851009-69-3

STAGE(1)

RGT M 121-44-8 Et₃N, X 543-27-1 ClCO₂Bu-1



SOL 109-99-9 THF
CON SUBSTAGE(1) 0 deg C
SUBSTAGE(2) 1 hour, 0 deg C

STAGE(2)

RCT V 334-88-3
SOL 60-29-7 Et2O
CON SUBSTAGE(1) 1 minute, 0 deg C
SUBSTAGE(2) 30 minutes, 0 deg C
SUBSTAGE(3) 45 minutes, room temperature

PRO W 851009-70-6

RX(36) OF 55 COMPOSED OF RX(2), RX(4), RX(5), RX(6), RX(7)
RX(36) C + L + V ==> Z





STAGE(1)
RGT AA 10035-10-6 HBr
SOL 7732-18-5 Water, 109-99-9 THF
CON SUBSTAGE(1) room temperature -> 0 deg C
SUBSTAGE(2) 0 deg C
SUBSTAGE(3) 1 hour, 0 deg C


STAGE(2)
RGT D 144-55-8 NaHCO3
SOL 7732-18-5 Water

PRO Z 851009-71-7

RX(37) OF 55 COMPOSED OF REACTION SEQUENCE RX(3), RX(4), RX(5)
AND REACTION SEQUENCE RX(1), RX(2), RX(4), RX(5)
...J + K ==> L...
...A + B + L ==> S




4 STEPS



START NEXT REACTION SEQUENCE

5 STEPS



YIELD 100%

RX(2) RCT C 791835-61-5
PRO H 791835-62-6
SOL 101-84-8 PhOPh
CON SUBSTAGE(1) 5 minutes, 250 deg C -> 230 deg C
SUBSTAGE(2) 7 minutes, 230 deg C -> 245 deg C
SUBSTAGE(3) cooled
SUBSTAGE(4) 0 deg C
NTE extended heating at 250 degree celsius would give decarboxylation of desired ester, reaction is done in two batches

RX(4) RCT H 791835-62-6, L 801282-34-8
RGT Q 534-17-8 Ca2CO3
PRO P 851009-68-2
SOL 872-50-4 NMBP
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 5 hours, 72 deg C


RX(5) RCT P 851009-68-2
RGT T 1310-73-2 NaOH
PRO S 851009-69-3
SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 1.5 hours, room temperature

RX(6) RCT S 851009-69-3
STAGE(1)
RGT M 121-44-8 Et3N, X 543-27-1 ClCO2Bu-1
SOL 109-99-9 THF
CON SUBSTAGE(1) 0 deg C
SUBSTAGE(2) 1 hour, 0 deg C



STAGE(2)
RCT V 334-88-3
SOL 60-29-7 Et2O
CON SUBSTAGE(1) 1 minute, 0 deg C
SUBSTAGE(2) 30 minutes, 0 deg C
SUBSTAGE(3) 45 minutes, room temperature

PRO W 851009-70-6

RX(7) RCT W 851009-70-6



2 HCl



4 STEPS

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(3) RCT J 572924-77-7, K 98-58-8

STAGE(1)
CAT 1122-58-3 4-DMAP
SOL 75-09-2 CH2Cl2
CON room temperature -> 0 deg C

STAGE(2)
RGT M 121-44-8 Et3N
CON SUBSTAGE(1) 3 minutes, 0 deg C
SUBSTAGE(2) 1 hour, 0 deg C
SUBSTAGE(3) 0 deg C -> room temperature
SUBSTAGE(4) 18 hours, room temperature

PRO L 801282-34-8

RX(1) RCT A 3575-32-4

STAGE(1)
RGT D 144-55-8 NaHCO3
SOL 7732-18-5 Water
CON neutralized

STAGE(2)

RCT B 762-42-5
 RGT E 62-53-3 PhNH₂
 SOL 67-56-1 MeOH
 CON SUBSTAGE(1) 0 deg C, neutralized
 SUBSTAGE(2) heated
 SUBSTAGE(3) 2 hours, 65 deg C
 SUBSTAGE(4) 14 hours, room temperature



PRO C 791835-61-5
 NTE exothermic reaction in second stage, incremental addition of aniline in second stage

RX(2) RCT C 791835-61-5
 PRO H 791835-62-6
 SOL 101-84-8 PhOPh
 CON SUBSTAGE(1) 5 minutes, 250 deg C -> 230 deg C
 SUBSTAGE(2) 7 minutes, 230 deg C -> 245 deg C
 SUBSTAGE(3) cooled
 SUBSTAGE(4) 0 deg C
 NTE extended heating at 250 degree celsius would give decarboxylation of desired ester, reaction is done in two batches




RX(4) RCT H 791835-62-6, L 801282-34-8
 RGT Q 534-17-8 Cs₂CO₃
 PRO P 851009-68-2
 SOL 872-50-4 NMEP
 CON SUBSTAGE(1) room temperature
 SUBSTAGE(2) 5 hours, 72 deg C

RX(5) RCT P 851009-68-2
 RGT T 1310-73-2 NaOH
 PRO S 851009-69-3
 SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF
 CON SUBSTAGE(1) room temperature
 SUBSTAGE(2) 1.5 hours, room temperature

RX(38) OF 55 COMPOSED OF REACTION SEQUENCE RX(3), RX(4), RX(5), RX(6)
 AND REACTION SEQUENCE RX(1), RX(2), RX(4), RX(5), RX(6)
 ...J + K ==> L...
 ...A + B + L + V ==> W



START NEXT REACTION SEQUENCE



RX(3) RCT J 572924-77-7, K 98-58-8
 STAGE(1)
 CAT 1122-58-3 4-DMAP
 SOL 75-09-2 CH₂Cl₂
 CON room temperature -> 0 deg C
 STAGE(2)
 RGT M 121-44-8 Et₃N
 CON SUBSTAGE(1) 3 minutes, 0 deg C
 SUBSTAGE(2) 1 hour, 0 deg C
 SUBSTAGE(3) 0 deg C -> room temperature
 SUBSTAGE(4) 16 hours, room temperature

PRO L 801282-34-8

RX(1) RCT A 3575-32-4
 STAGE(1)
 RGT D 144-55-8 NaHCO₃
 SOL 7732-18-5 Water
 CON neutralized
 STAGE(2)
 RCT B 762-42-5
 RGT E 62-53-3 PhNH₂
 SOL 67-56-1 MeOH
 CON SUBSTAGE(1) 0 deg C, neutralized
 SUBSTAGE(2) heated
 SUBSTAGE(3) 2 hours, 65 deg C
 SUBSTAGE(4) 14 hours, room temperature

PRO C 791835-61-5
 NTE exothermic reaction in second stage, incremental addition of aniline in second stage

RX(2) RCT C 791835-61-5
 PRO H 791835-62-6
 SOL 101-84-8 PhOPh
 CON SUBSTAGE(1) 5 minutes, 250 deg C -> 230 deg C
 SUBSTAGE(2) 7 minutes, 230 deg C -> 245 deg C
 SUBSTAGE(3) cooled
 SUBSTAGE(4) 0 deg C
 NTE extended heating at 250 degree celsius would give decarboxylation of desired ester, reaction is done in two

batches


RX(4) RCT H 791835-62-6, L 801282-34-8
 RGT Q 534-17-8 Cs₂CO₃
 PRO P 851009-68-2
 SOL 872-50-4 NMEP
 CON SUBSTAGE(1) room temperature
 SUBSTAGE(2) 5 hours, 72 deg C



RX(5) RCT P 851009-68-2
 RGT T 1310-73-2 NaOH
 PRO S 851009-69-3
 SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF
 CON SUBSTAGE(1) room temperature
 SUBSTAGE(2) 1.5 hours, room temperature

RX(6) RCT S 851009-69-3
 STAGE(1)
 RGT M 121-44-8 Et₃N, X 543-27-1 ClCO₂Bu-1
 SOL 109-99-9 THF
 CON SUBSTAGE(1) 0 deg C
 SUBSTAGE(2) 1 hour, 0 deg C
 STAGE(2)
 RCT V 334-86-3
 SOL 60-29-7 Et₂O
 CON SUBSTAGE(1) 1 minute, 0 deg C
 SUBSTAGE(2) 30 minutes, 0 deg C
 SUBSTAGE(3) 45 minutes, room temperature

PRO W 851009-70-6

RX(39) OF 55 COMPOSED OF RX(1), RX(2), RX(4), RX(5), RX(6), RX(7)
 RX(39) A + B + L + V ==> Z





RX(1) RCT A 3575-32-4

STAGE(1)

RGT D 144-55-8 NaHCO₃
SOL 7732-18-5 Water
CON neutralized

STAGE(2)

RCT B 762-42-5
RGT E 62-53-3 PhNH₂
SOL 67-56-1 MeOH
CON SUBSTAGE(1) 0 deg C, neutralized
SUBSTAGE(2) heated
SUBSTAGE(3) 2 hours, 65 deg C
SUBSTAGE(4) 14 hours, room temperature





PRO C 791835-61-5

NTE exothermic reaction in second stage, incremental addition of aniline in second stage

RX(2) RCT C 791835-61-5

PRO H 791835-62-6
SOL 101-84-8 PhOPh

CON SUBSTAGE(1) 5 minutes, 250 deg C -> 230 deg C



SUBSTAGE(2) 7 minutes, 230 deg C -> 245 deg C
SUBSTAGE(3) cooled
SUBSTAGE(4) 0 deg C
NTE extended heating at 250 degree celsius would give decarboxylation of desired ester, reaction is done in two batches

RX(4) RCT H 791835-62-6, L 801282-34-8
RGT Q 534-17-8 Ca2CO₃
PRO P 851009-68-2
SOL 872-50-4 NMEP
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 5 hours, 72 deg C

RX(5) RCT P 851009-68-2
RGT T 1310-73-2 NaOH
PRO S 851009-69-3
SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 1.5 hours, room temperature

RX(6) RCT S 851009-69-3

STAGE(1)

RGT M 121-44-8 Et3N, X 543-27-1 ClCO2Bu-1
SOL 109-99-9 THF
CON SUBSTAGE(1) 0 deg C
SUBSTAGE(2) 1 hour, 0 deg C

STAGE(2)

RCT V 334-88-3
SOL 60-29-7 Et2O
CON SUBSTAGE(1) 1 minute, 0 deg C
SUBSTAGE(2) 30 minutes, 0 deg C
SUBSTAGE(3) 45 minutes, room temperature

PRO W 851009-70-6

RX(7) RCT W 851009-70-6

STAGE(1)

RGT AA 10035-10-6 HBr
SOL 7732-18-5 Water, 109-99-9 THF
CON SUBSTAGE(1) room temperature -> 0 deg C
SUBSTAGE(2) 0 deg C
SUBSTAGE(3) 1 hour, 0 deg C

STAGE(2)

RGT D 144-55-8 NaHCO₃
SOL 7732-18-5 Water

PRO Z 851009-71-7

RX(42) OF 55 COMPOSED OF RX(2), RX(4), RX(5), RX(6), RX(7), RX(8)

RX(42) C + L + V + AB ==> AC

RX(2) RCT C 791835-61-5
PRO H 791835-62-6
SOL 101-84-8 PhOPh
CON SUBSTAGE(1) 5 minutes, 250 deg C -> 230 deg C
SUBSTAGE(2) 7 minutes, 230 deg C -> 245 deg C
SUBSTAGE(3) cooled
SUBSTAGE(4) 0 deg C

NTE extended heating at 250 degree celsius would give decarboxylation of desired ester, reaction is done in two batches

RX(4) RCT H 791835-62-6, L 801282-34-8
RGT Q 534-17-8 Ca2CO₃
PRO P 851009-68-2
SOL 872-50-4 NMEP
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 5 hours, 72 deg C

RX(5) RCT P 851009-68-2
RGT T 1310-73-2 NaOH
PRO S 851009-69-3
SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 1.5 hours, room temperature

RX(6) RCT S 851009-69-3

STAGE(1)

RGT M 121-44-8 Et3N, X 543-27-1 ClCO2Bu-1
SOL 109-99-9 THF
CON SUBSTAGE(1) 0 deg C
SUBSTAGE(2) 1 hour, 0 deg C

STAGE(2)

RCT V 334-88-3
SOL 60-29-7 Et2O
CON SUBSTAGE(1) 1 minute, 0 deg C
SUBSTAGE(2) 30 minutes, 0 deg C
SUBSTAGE(3) 45 minutes, room temperature

PRO W 851009-70-6

RX(7) RCT W 851009-70-6

STAGE(1)

RGT AA 10035-10-6 HBr
SOL 7732-18-5 Water, 109-99-9 THF
CON SUBSTAGE(1) room temperature -> 0 deg C
SUBSTAGE(2) 0 deg C
SUBSTAGE(3) 1 hour, 0 deg C

STAGE(2)

RGT D 144-55-8 NaHCO₃
SOL 7732-18-5 Water


PRO Z 851009-71-7

RX(8) RCT Z 851009-71-7, AB 572923-98-9


PRO AC 851009-72-8
SOL 67-63-0 Me2CHOH
CON SUBSTAGE(1) 5 minutes, heated
SUBSTAGE(2) 1.5 hours

NTE overall yield over 4 steps is 53%


RX(44) OF 55 COMPOSED OF RX(1), RX(2), RX(4), RX(5), RX(6), RX(7), RX(8)
 RX(44) A + B + L + V + AB ==> AC




A



B




L



AB

7
STEPS



AC

RX(1) RCT A 3575-32-4

STAGE(1)

ROT D 144-55-8 NaHCO3
 SOL 7732-18-5 Water
 CON neutralized

STAGE(2)

RCT B 762-42-5
 ROT E 62-53-3 PhNH2
 SOL 67-56-1 MeOH
 CON SUBSTAGE(1) 0 deg C, neutralized
 SUBSTAGE(2) heated
 SUBSTAGE(3) 2 hours, 65 deg C
 SUBSTAGE(4) 14 hours, room temperature

PRO C 791835-61-5

NTE exothermic reaction in second stage, incremental addition of aniline in second stage

RX(2)

RCT C 791835-61-5
 PRO H 791835-62-6
 SOL 101-84-8 PhOPh
 CON SUBSTAGE(1) 5 minutes, 250 deg C -> 230 deg C
 SUBSTAGE(2) 7 minutes, 230 deg C -> 245 deg C
 SUBSTAGE(3) cooled
 SUBSTAGE(4) 0 deg C

NTE extended heating at 250 degrees celsius would give decarboxylation of desired ester, reaction is done in two batches

RX(4)

RCT H 791835-62-6, L 801282-34-8
 ROT Q 534-17-8 Ca2CO3
 PRO P 851009-68-2
 SOL 872-80-4 NMP
 CON SUBSTAGE(1) room temperature
 SUBSTAGE(2) 5 hours, 72 deg C

RX(5)

RCT P 851009-68-2
 ROT T 1310-73-2 NaOH
 PRO S 851009-69-3
 SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF
 CON SUBSTAGE(1) room temperature
 SUBSTAGE(2) 1.5 hours, room temperature

RX(6) RCT S 851009-69-3

STAGE(1)

ROT M 121-44-8 Et3N, X 543-27-1 ClCO2Bu-t
 SOL 109-99-9 THF
 CON SUBSTAGE(1) 0 deg C
 SUBSTAGE(2) 1 hour, 0 deg C

STAGE(2)

RCT V 334-88-3
 SOL 60-29-7 Et2O
 CON SUBSTAGE(1) 1 minute, 0 deg C
 SUBSTAGE(2) 30 minutes, 0 deg C
 SUBSTAGE(3) 45 minutes, room temperature

PRO W 851009-70-6

RX(7) RCT W 851009-70-6

STAGE(1)

ROT AA 10035-10-6 HBr
 SOL 7732-18-5 Water, 109-99-9 THF
 CON SUBSTAGE(1) room temperature -> 0 deg C
 SUBSTAGE(2) 0 deg C
 SUBSTAGE(3) 1 hour, 0 deg C

STAGE(2)


ROT D 144-55-8 NaHCO3
 SOL 7732-18-5 Water

PRO Z 851009-71-7


RX(8) RCT Z 851009-71-7, AB 572923-98-9

PRO AC 851009-72-8
 SOL 67-63-0 Me2CHOH
 CON SUBSTAGE(1) 5 minutes, heated
 SUBSTAGE(2) 1.5 hours
 NTE overall yield over 4 steps is 53%

RX(45) OF 55 COMPOSED OF REACTION SEQUENCE RX(1), RX(4), RX(5), RX(6), RX(7)
 AND REACTION SEQUENCE RX(2), RX(4), RX(5), RX(6), RX(7)
 ...J + K ==> L...
 ... C + L + V ==> Z




J




K

5
STEPS




L

START NEXT REACTION SEQUENCE



C



L



V

5
STEPS

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(3) RCT J 572924-77-7, K 98-58-8

STAGE(1)

CAT 1122-58-3 4-DMAP
SOL 75-09-2 CH₂Cl₂
CON room temperature -> 0 deg C

STAGE(2)

RGT M 121-44-8 Et₃N
CON SUBSTAGE(1) 3 minutes, 0 deg C
SUBSTAGE(2) 1 hour, 0 deg C
SUBSTAGE(3) 0 deg C -> room temperature
SUBSTAGE(4) 18 hours, room temperature

PRO L 801282-34-8

RX(2) RCT C 791835-61-5

PRO H 791835-62-6
SOL 101-84-8 PhOPh
CON SUBSTAGE(1) 5 minutes, 250 deg C -> 230 deg C
SUBSTAGE(2) 7 minutes, 230 deg C -> 245 deg C
SUBSTAGE(3) cooled
SUBSTAGE(4) 0 deg C

NTE extended heating at 250 degree celsius would give decarboxylation of desired ester, reaction is done in two batches

RX(4) RCT H 791835-62-6, L 801282-34-8

RGT Q 534-17-8 Ca₂CO₃
PRO P 851009-68-2
SOL 872-50-4 NMSP
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 5 hours, 72 deg C

RX(5) RCT P 851009-68-2

RGT T 1310-73-2 NaOH
PRO S 851009-69-3
SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 1.5 hours, room temperature

RX(6) RCT S 851009-69-3

STAGE(1)

RGT M 121-44-8 Et₃N, X 543-27-1 ClCO₂Bu-i
SOL 109-99-9 THF
CON SUBSTAGE(1) 0 deg C
SUBSTAGE(2) 1 hour, 0 deg C

STAGE(2)

RCT V 334-88-3
SOL 60-29-7 Et₂O
CON SUBSTAGE(1) 1 minute, 0 deg C
SUBSTAGE(2) 30 minutes, 0 deg C
SUBSTAGE(3) 45 minutes, room temperature

PRO W 851009-70-6

RX(7) RCT W 851009-70-6

STAGE(1)

RGT AA 10035-10-6 HBr
SOL 7732-18-5 Water, 109-99-9 THF
CON SUBSTAGE(1) room temperature -> 0 deg C
SUBSTAGE(2) 0 deg C
SUBSTAGE(3) 1 hour, 0 deg C

STAGE(2)

RGT D 144-55-8 NaHCO₃
SOL 7732-18-5 Water



PRO Z 851009-71-7

RX(46) OF 55 COMPOSED OF REACTION SEQUENCE RX(3), RX(4), RX(5), RX(6), RX(7),
RX(8)




AND REACTION SEQUENCE RX(2), RX(4), RX(5), RX(6), RX(7), RX(8)

...J + K ==> L...

... C + L + V + AB ==> AC



START NEXT REACTION SEQUENCE



* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(3) RCT J 572924-77-7, K 98-58-8

STAGE(1)

CAT 1122-58-3 4-DMAP
SOL 75-09-2 CH₂Cl₂
CON room temperature -> 0 deg C

STAGE(2)

RGT M 121-44-8 Et₃N
CON SUBSTAGE(1) 3 minutes, 0 deg C
SUBSTAGE(2) 1 hour, 0 deg C
SUBSTAGE(3) 0 deg C -> room temperature
SUBSTAGE(4) 18 hours, room temperature

PRO L 801282-34-8

RX(2) RCT C 791835-61-5

PRO H 791835-62-6
SOL 101-84-8 PhOPh
CON SUBSTAGE(1) 5 minutes, 250 deg C -> 230 deg C
SUBSTAGE(2) 7 minutes, 230 deg C -> 245 deg C
SUBSTAGE(3) cooled
SUBSTAGE(4) 0 deg C

NTE extended heating at 250 degree celsius would give decarboxylation of desired ester, reaction is done in two batches

RX(4) RCT H 791835-62-6, L 801282-34-8

RGT Q 534-17-8 Ca₂CO₃
PRO P 851009-68-2
SOL 872-50-4 NMSP
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 5 hours, 72 deg C

RX(5) RCT P 851009-68-2

RGT T 1310-73-2 NaOH
PRO S 851009-69-3
SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 1.5 hours, room temperature

RX(6) RCT S 851009-69-3

STAGE(1)

RGT M 121-44-8 Et₃N, X 543-27-1 ClCO₂Bu-i
SOL 109-99-9 THF
CON SUBSTAGE(1) 0 deg C
SUBSTAGE(2) 1 hour, 0 deg C

STAGE(2)

RCT V 334-88-3
SOL 60-29-7 Et₂O
CON SUBSTAGE(1) 1 minute, 0 deg C
SUBSTAGE(2) 30 minutes, 0 deg C
SUBSTAGE(3) 45 minutes, room temperature

PRO W 851009-70-6

RX(7) RCT W 851009-70-6

STAGE(1)

RGT AA 10035-10-6 HBr
SOL 7732-18-5 Water, 109-99-9 THF
CON SUBSTAGE(1) room temperature -> 0 deg C
SUBSTAGE(2) 0 deg C
SUBSTAGE(3) 1 hour, 0 deg C

STAGE(2)



RGT D 144-55-8 NaHCO₃
SOL 7732-18-5 Water

PRO Z 851009-71-7


RX(8) RCT Z 851009-71-7, AB 572923-98-9

PRO AC 851009-72-8
SOL 67-63-0 Me₂CHOH
CON SUBSTAGE(1) 5 minutes, heated
SUBSTAGE(2) 1.5 hours
NTE overall yield over 4 steps is 53%

RX(47) OF 55 COMPOSED OF REACTION SEQUENCE RX(3), RX(4), RX(5), RX(6), RX(7)
AND REACTION SEQUENCE RX(1), RX(2), RX(4), RX(5), RX(6), RX(7)
...J + K ==> L...
...A + B + L + V ==> Z



START NEXT REACTION SEQUENCE



NTE exothermic reaction in second stage, incremental addition of aniline in second stage

RX(2) RCT C 791835-61-5
PRO H 791835-62-6
SOL 101-84-8 PhOPh
CON SUBSTAGE(1) 5 minutes, 250 deg C -> 230 deg C
SUBSTAGE(2) 7 minutes, 230 deg C -> 245 deg C
SUBSTAGE(3) cooled
SUBSTAGE(4) 0 deg C

NTE extended heating at 250 degree celsius would give decarboxylation of desired ester, reaction is done in two batches

RX(4) RCT H 791835-62-6, L 801202-34-8
ROT Q 534-17-8 Ca2CO3
PRO P 851009-68-2
SOL 872-50-4 NMSP
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 5 hours, 72 deg C

RX(5) RCT P 851009-68-2
ROT T 1310-73-2 NaOH
PRO S 851009-69-3
SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 1.5 hours, room temperature

RX(6) RCT S 851009-69-3
STAGE(1)
ROT M 121-44-8 Et3N, X 543-27-1 ClCO2Bu-i
SOL 109-99-9 THF
CON SUBSTAGE(1) 0 deg C
SUBSTAGE(2) 1 hour, 0 deg C
STAGE(2)
RCT V 334-88-3
SOL 60-29-7 Et2O
CON SUBSTAGE(1) 1 minute, 0 deg C
SUBSTAGE(2) 30 minutes, 0 deg C
SUBSTAGE(3) 45 minutes, room temperature


PRO W 851009-70-6

RX(7) RCT W 851009-70-6
STAGE(1)
ROT AA 10035-10-6 HBr
SOL 7732-18-5 Water, 109-99-9 THF
CON SUBSTAGE(1) room temperature -> 0 deg C
SUBSTAGE(2) 0 deg C
SUBSTAGE(3) 1 hour, 0 deg C

STAGE(2)
ROT D 144-55-8 NaHCO3
SOL 7732-18-5 Water

PRO Z 851009-71-7

RX(48) OF 55 COMPOSED OF REACTION SEQUENCE RX(3), RX(4), RX(5), RX(6), RX(7),
RX(8)



6
STEPS

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(3) RCT J 572924-77-7, K 98-58-8

STAGE(1)
CAT 1122-58-3 4-DMAP
SOL 75-09-2 CH2Cl2
CON room temperature -> 0 deg C

STAGE(2)
ROT M 121-44-8 Et3N
CON SUBSTAGE(1) 3 minutes, 0 deg C
SUBSTAGE(2) 1 hour, 0 deg C
SUBSTAGE(3) 0 deg C -> room temperature
SUBSTAGE(4) 18 hours, room temperature

PRO L 801202-34-8



RX(1) RCT A 3575-32-4

STAGE(1)
ROT D 144-55-8 NaHCO3
SOL 7732-18-5 Water
CON neutralized


STAGE(2)
RCT B 762-42-5
ROT E 62-53-3 PhNH2
SOL 67-56-1 MeOH
CON SUBSTAGE(1) 0 deg C, neutralized
SUBSTAGE(2) heated
SUBSTAGE(3) 2 hours, 65 deg C
SUBSTAGE(4) 14 hours, room temperature



PRO C 791835-61-5

RX(8) AND REACTION SEQUENCE RX(1), RX(2), RX(4), RX(5), RX(6), RX(7),
...J + K ==> L...
...A + B + L + V + AB ==> AC



START NEXT REACTION SEQUENCE





* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(3) RCT J 572924-77-7, K 98-58-8

STAGE(1)
CAT 1122-58-3 4-DMAP
SOL 75-09-2 CH₂Cl₂
CON room temperature -> 0 deg C

STAGE(2)
RGT M 121-44-8 Et₃N
CON SUBSTAGE(1) 3 minutes, 0 deg C
SUBSTAGE(2) 1 hour, 0 deg C
SUBSTAGE(3) 0 deg C -> room temperature
SUBSTAGE(4) 18 hours, room temperature

PRO L 801282-34-8




RX(1) RCT A 3575-32-4

STAGE(1)
RGT D 144-55-8 NaHCO₃
SOL 7732-18-5 Water
CON neutralized

STAGE(2)
RCT B 762-42-5
RGT E 62-53-3 PhNH₂
SOL 67-56-1 MeOH
CON SUBSTAGE(1) 0 deg C, neutralized
SUBSTAGE(2) heated
SUBSTAGE(3) 2 hours, 65 deg C

RX(8) RCT Z 851009-71-7, AB 572923-98-9
PRO AC 851009-72-8
SOL 67-63-0 Me₂CHOH
CON SUBSTAGE(1) 5 minutes, heated
SUBSTAGE(2) 1.5 hours
NTE overall yield over 4 steps is 53%

RX(51) OF 55 COMPOSED OF RX(2), RX(4), RX(5), RX(6), RX(7), RX(8), RX(9)
RX(51) C + L + V + AB ==> AS



* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(2) RCT C 791835-61-5
PRO H 791835-62-6
SOL 101-84-8 PhOPh
CON SUBSTAGE(1) 5 minutes, 250 deg C -> 230 deg C

SUBSTAGE(4) 14 hours, room temperature
PRO C 791835-61-5
NTE exothermic reaction in second stage, incremental addition of aniline in second stage
RX(2) RCT C 791835-61-5
PRO H 791835-62-6
SOL 101-84-8 PhOPh
CON SUBSTAGE(1) 5 minutes, 250 deg C -> 230 deg C
SUBSTAGE(2) 7 minutes, 230 deg C -> 245 deg C
SUBSTAGE(3) cooled
SUBSTAGE(4) 0 deg C
NTE extended heating at 250 degree celsius would give decarboxylation of desired ester, reaction is done in two batches
RX(4) RCT H 791835-62-6, L 801282-34-8
RGT Q 534-17-8 Ca₂CO₃
PRO P 851009-68-2
SOL 872-50-4 NMEP
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 5 hours, 72 deg C

RX(5) RCT P 851009-68-2
RGT T 1310-73-2 NaOH
PRO S 851009-69-3
SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 1.5 hours, room temperature

RX(6) RCT S 851009-69-3

STAGE(1)
RGT M 121-44-8 Et₃N, X 543-27-1 ClCO₂Bu-1
SOL 109-99-9 THF
CON SUBSTAGE(1) 0 deg C
SUBSTAGE(2) 1 hour, 0 deg C

STAGE(2)
RCT V 334-88-3
SOL 60-29-7 Et₂O
CON SUBSTAGE(1) 1 minute, 0 deg C
SUBSTAGE(2) 30 minutes, 0 deg C
SUBSTAGE(3) 45 minutes, room temperature

PRO W 851009-70-6

RX(7) RCT W 851009-70-6

STAGE(1)
RGT AA 10035-10-6 HBr
SOL 7732-18-5 Water, 109-99-9 THF
CON SUBSTAGE(1) room temperature -> 0 deg C
SUBSTAGE(2) 0 deg C
SUBSTAGE(3) 1 hour, 0 deg C

STAGE(2)
RGT D 144-55-8 NaHCO₃
SOL 7732-18-5 Water

PRO Z 851009-71-7

SUBSTAGE(2) 7 minutes, 230 deg C -> 245 deg C
SUBSTAGE(3) cooled
SUBSTAGE(4) 0 deg C
NTE extended heating at 250 degree celsius would give decarboxylation of desired ester, reaction is done in two batches

RX(4) RCT H 791835-62-6, L 801282-34-8
RGT Q 534-17-8 Ca₂CO₃
PRO P 851009-68-2
SOL 872-50-4 NMEP
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 5 hours, 72 deg C

RX(5) RCT P 851009-68-2
RGT T 1310-73-2 NaOH
PRO S 851009-69-3
SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 1.5 hours, room temperature

RX(6) RCT S 851009-69-3

STAGE(1)
RGT M 121-44-8 Et₃N, X 543-27-1 ClCO₂Bu-1
SOL 109-99-9 THF
CON SUBSTAGE(1) 0 deg C
SUBSTAGE(2) 1 hour, 0 deg C

STAGE(2)
RCT V 334-88-3
SOL 60-29-7 Et₂O
CON SUBSTAGE(1) 1 minute, 0 deg C
SUBSTAGE(2) 30 minutes, 0 deg C
SUBSTAGE(3) 45 minutes, room temperature

PRO W 851009-70-6

RX(7) RCT W 851009-70-6

STAGE(1)
RGT AA 10035-10-6 HBr
SOL 7732-18-5 Water, 109-99-9 THF
CON SUBSTAGE(1) room temperature -> 0 deg C
SUBSTAGE(2) 0 deg C
SUBSTAGE(3) 1 hour, 0 deg C

STAGE(2)
RGT D 144-55-8 NaHCO₃
SOL 7732-18-5 Water

PRO Z 851009-71-7

RX(8) RCT Z 851009-71-7, AB 572923-98-9
PRO AC 851009-72-8
SOL 67-63-0 Me₂CHOH
CON SUBSTAGE(1) 5 minutes, heated
SUBSTAGE(2) 1.5 hours
NTE overall yield over 4 steps is 53%

RX(9) RCT AC 851009-72-8

STAGE(1)

RGT T 1310-73-2 NaOH
SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 5 hours, room temperature

STAGE(2)



RGT AF 7647-01-0 HCl
SOL 7732-18-5 Water
CON room temperature, pH 6

STAGE(3)

RGT T 1310-73-2 NaOH
SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF
CON 15 minutes, room temperature


PRO AB 851009-74-0

RX(52) OF 55 COMPOSED OF RX(1), RX(2), RX(4), RX(5), RX(6), RX(7), RX(8), RX(9)
RX(52) A + B + L + V + AB ==> AE



A ●3 HCl


B



L



V



AB

8
STEPS

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(1) RCT A 3575-32-4

STAGE(1)

RGT D 144-55-8 NaHCO3
SOL 7732-18-5 Water
CON neutralized

STAGE(2)

RGT B 762-42-5
RGT E 62-53-3 PhNH2
SOL 67-56-1 MeOH
CON SUBSTAGE(1) 0 deg C, neutralized
SUBSTAGE(2) heated
SUBSTAGE(3) 2 hours, 65 deg C
SUBSTAGE(4) 14 hours, room temperature

PRO C 791835-61-5

NTE exothermic reaction in second stage, incremental addition of aniline in second stage

RX(2) RCT C 791835-61-5

PRO H 791835-62-6

SOL 101-84-8 PhOPh

CON SUBSTAGE(1) 5 minutes, 250 deg C -> 230 deg C
SUBSTAGE(2) 7 minutes, 230 deg C -> 245 deg C
SUBSTAGE(3) cooled
SUBSTAGE(4) 0 deg C

NTE extended heating at 250 degree celsius would give decarboxylation of desired ester, reaction is done in two batches

RX(4) RCT H 791835-62-6, L 601262-34-8

RGT Q 534-17-8 Cs2CO3

PRO P 851009-68-2

SOL 872-50-4 NMMP

CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 5 hours, 72 deg C

RX(5) RCT P 851009-68-2

RGT T 1310-73-2 NaOH

PRO S 851009-69-3

SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 1.5 hours, room temperature

RX(6) RCT S 851009-69-3

STAGE(1)

RGT M 121-44-8 Et3N, X 543-27-1 ClCO2Bu-1
SOL 109-99-9 THF

CON SUBSTAGE(1) 0 deg C
SUBSTAGE(2) 1 hour, 0 deg C

STAGE(2)

RCT V 334-88-3
SOL 60-29-7 Et2O
CON SUBSTAGE(1) 1 minute, 0 deg C
SUBSTAGE(2) 30 minutes, 0 deg C
SUBSTAGE(3) 45 minutes, room temperature

PRO W 851009-70-6

RX(7) RCT W 851009-70-6

STAGE(1)

RGT AA 10035-10-6 HBr
SOL 7732-18-5 Water, 109-99-9 THF
CON SUBSTAGE(1) room temperature -> 0 deg C
SUBSTAGE(2) 0 deg C
SUBSTAGE(3) 1 hour, 0 deg C

STAGE(2)

RGT D 144-55-8 NaHCO3
SOL 7732-18-5 Water

PRO Z 851009-71-7

RX(8) RCT Z 851009-71-7, AB 572923-98-9

PRO AC 851009-72-8

SOL 67-63-0 Me2CHOH

CON SUBSTAGE(1) 5 minutes, heated
SUBSTAGE(2) 1.5 hours

NTE overall yield over 4 steps is 53%

RX(9) RCT AC 851009-72-8

STAGE(1)

RGT T 1310-73-2 NaOH
SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 5 hours, room temperature

STAGE(2)

RGT AF 7647-01-0 HCl
SOL 7732-18-5 Water
CON room temperature, pH 6

STAGE(3)

RGT T 1310-73-2 NaOH
SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF
CON 15 minutes, room temperature

PRO AB 851009-74-0

RX(54) OF 55 COMPOSED OF REACTION SEQUENCE RX(3), RX(4), RX(5), RX(6), RX(7),


RX(8), RX(9)

AND REACTION SEQUENCE RX(2), RX(4), RX(5), RX(6), RX(7), RX(8),


RX(9)

...J + K ==> L...


... C + L + V + AB ==> AE



J




K





L

START NEXT REACTION SEQUENCE



C

7
STEPS

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(3) RCT J 572924-77-7, K 98-58-8

STAGE(1)
CAT 1122-58-3 4-DMAP
SOL 75-09-2 CH2Cl2
CON room temperature -> 0 deg C

STAGE(2)
RGT M 121-44-8 Et3N
CON SUBSTAGE(1) 3 minutes, 0 deg C
SUBSTAGE(2) 1 hour, 0 deg C
SUBSTAGE(3) 0 deg C -> room temperature
SUBSTAGE(4) 18 hours, room temperature

PRO L 801282-34-8



RX(2) RCT C 791835-61-5
PRO H 791835-62-6
SOL 101-84-8 PhOPh
CON SUBSTAGE(1) 5 minutes, 250 deg C -> 230 deg C
SUBSTAGE(2) 7 minutes, 230 deg C -> 245 deg C
SUBSTAGE(3) cooled
SUBSTAGE(4) 0 deg C
NTE extended heating at 250 degrees celsius would give decarboxylation of desired ester, reaction is done in two batches

RX(4) RCT H 791835-62-6, L 801282-34-8
RGT Q 534-17-8 Ca2CO3
PRO P 851009-68-2

STAGE(3)
RGT T 1310-73-2 NaOH
SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF
CON 15 minutes, room temperature

PRO AE 851009-74-0

RX(55) OF 55 COMPOSED OF REACTION SEQUENCE RX(3), RX(4), RX(5), RX(6), RX(7),
RX(8), RX(9)
AND REACTION SEQUENCE RX(1), RX(2), RX(4), RX(5), RX(6), RX(7),
RX(8), RX(9)
...J + K ==> L...
...A + D + L + V + AB ==> AE



START NEXT REACTION SEQUENCE

SOL 872-50-4 NMEP
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 5 hours, 72 deg C

RX(5) RCT P 851009-68-2
RGT T 1310-73-2 NaOH
PRO S 851009-69-3
SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 1.5 hours, room temperature

RX(6) RCT S 851009-69-3

STAGE(1)
RGT M 121-44-8 Et3N, X 543-27-1 ClCO2Bu-1
SOL 109-99-9 THF
CON SUBSTAGE(1) 0 deg C
SUBSTAGE(2) 1 hour, 0 deg C

STAGE(2)
RGT V 334-88-3
SOL 60-29-7 Et2O
CON SUBSTAGE(1) 1 minute, 0 deg C
SUBSTAGE(2) 30 minutes, 0 deg C
SUBSTAGE(3) 45 minutes, room temperature

PRO W 851009-70-6

RX(7) RCT W 851009-70-6

STAGE(1)
RGT AA 10035-10-6 HBr
SOL 7732-18-5 Water, 109-99-9 THF
CON SUBSTAGE(1) room temperature -> 0 deg C
SUBSTAGE(2) 0 deg C
SUBSTAGE(3) 1 hour, 0 deg C

STAGE(2)
RGT D 144-55-8 NaHCO3
SOL 7732-18-5 Water


PRO Z 851009-71-7

RX(8) RCT Z 851009-71-7, AB 572923-98-9
PRO AC 851009-72-8
SOL 67-63-0 Me2CHOH
CON SUBSTAGE(1) 5 minutes, heated
SUBSTAGE(2) 1.5 hours
NTE overall yield over 4 steps is 53%



RX(9) RCT AC 851009-72-8

STAGE(1)
RGT T 1310-73-2 NaOH
SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF
CON SUBSTAGE(1) room temperature
SUBSTAGE(2) 5 hours, room temperature

STAGE(2)
RGT AF 7647-01-0 HCl
SOL 7732-18-5 Water
CON room temperature, pH 6



2 HCl

8
STEPS

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(3) RCT J 572924-77-7, K 98-58-8

STAGE(1)
CAT 1122-58-3 4-DMAP
SOL 75-09-2 CH2Cl2
CON room temperature -> 0 deg C

STAGE(2)
RGT M 121-44-8 Et3N
CON SUBSTAGE(1) 3 minutes, 0 deg C
SUBSTAGE(2) 1 hour, 0 deg C
SUBSTAGE(3) 0 deg C -> room temperature
SUBSTAGE(4) 18 hours, room temperature

PRO L 801282-34-8

RX(1) RCT A 3575-32-4

STAGE(1)

RGT D 144-55-8 NaHCO₃

SOL 7732-18-5 Water

CON neutralized

STAGE(2)

RCT B 762-42-5

RGT E 62-53-3 PhNH₂

SOL 67-56-1 MeOH

CON SUBSTAGE(1) 0 deg C, neutralized

SUBSTAGE(2) heated

SUBSTAGE(3) 2 hours, 65 deg C

SUBSTAGE(4) 14 hours, room temperature

PRO C 791835-61-5

NTE exothermic reaction in second stage, incremental addition of aniline in second stage

RX(2) RCT C 791835-61-5

PRO H 791835-62-6

SOL 101-84-8 PhOPh

CON SUBSTAGE(1) 5 minutes, 250 deg C -> 230 deg C

SUBSTAGE(2) 7 minutes, 230 deg C -> 245 deg C

SUBSTAGE(3) cooled

SUBSTAGE(4) 0 deg C

NTE extended heating at 250 degree celsius would give decarboxylation of desired ester, reaction is done in two batches

RX(4) RCT H 791835-62-6, L 801282-34-8

RGT G 534-17-8 Ca₂CO₃

PRO P 851009-68-2

SOL 872-50-4 NMEP

CON SUBSTAGE(1) room temperature

SUBSTAGE(2) 5 hours, 72 deg C

RX(5) RCT P 851009-68-2

RGT T 1310-73-2 NaOH

PRO S 851009-69-3

SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF

CON SUBSTAGE(1) room temperature

SUBSTAGE(2) 1.5 hours, room temperature

RX(6) RCT S 851009-69-3

STAGE(1)

RGT M 121-44-8 Et₃N, X 543-27-1 ClCO₂Bu-t

SOL 109-99-9 THF

CON SUBSTAGE(1) 0 deg C

SUBSTAGE(2) 1 hour, 0 deg C

STAGE(2)

RCT V 334-88-3

SOL 60-29-7 Et₂O

CON SUBSTAGE(1) 1 minute, 0 deg C

SUBSTAGE(2) 30 minutes, 0 deg C

SUBSTAGE(3) 45 minutes, room temperature

PRO W 851009-70-6

RX(7) RCT W 851009-70-6

STAGE(1)

RGT AA 10035-10-6 HBr

SOL 7732-18-5 Water, 109-99-9 THF

CON SUBSTAGE(1) room temperature -> 0 deg C

SUBSTAGE(2) 0 deg C

SUBSTAGE(3) 1 hour, 0 deg C

STAGE(2)

RGT D 144-55-8 NaHCO₃

SOL 7732-18-5 Water

PRO Z 851009-71-7

RX(8) RCT Z 851009-71-7, AB 572923-98-9

PRO AC 851009-72-8

SOL 67-63-0 Me₂CHOH

CON SUBSTAGE(1) 5 minutes, heated

SUBSTAGE(2) 1.5 hours

NTE overall yield over 4 steps is 53%

RX(9) RCT AC 851009-72-8

STAGE(1)

RGT T 1310-73-2 NaOH

SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF

CON SUBSTAGE(1) room temperature

SUBSTAGE(2) 5 hours, room temperature

STAGE(2)

RGT AF 7647-01-0 HCl

SOL 7732-18-5 Water

CON room temperature, pH 6

STAGE(3)

RGT T 1310-73-2 NaOH

SOL 7732-18-5 Water, 67-56-1 MeOH, 109-99-9 THF

CON 15 minutes, room temperature

PRO AB 851009-74-0

L12 ANSWER 2 OF 18 CASREACT COPYRIGHT 2007 ACS on STM

ACCESSION NUMBER: 142:94101 CASREACT Pull-text

TITLE: Design and synthesis of dipeptide-type HIV-1 protease inhibitors with high antiviral activity

AUTHOR(S): Kimura, Tooru; Hidaka, Kouichi; Abdel-Rahman, Hamdy M.; Matsumoto, Hikaru; Tanaka, Yoshiaki; Matsui, Yasuko; Hayashi, Yoshio; Kiso, Yoshiaki

CORPORATE SOURCE: Department of Medicinal Chemistry, Center for Frontier Research in Medicinal Science, Kyoto Pharmaceutical University, Kyoto, 607-8412, Japan

SOURCE: Peptide Science (2003), Volume Date 2004, 40th, 241-244


CODEN: PSCIFQ; ISSN: 1344-7661

PUBLISHER: Japanese Peptide Society

DOCUMENT TYPE: Journal

LANGUAGE: English


GI




AB A symposium report. A series of peptidomimetic HIV protease inhibitors, e.g. I, containing allophenylisoretinate [Aps, (28,38)-3-amino-2-hydroxy-4-phenylbutyric acid] with a hydroxymethylcarbonyl (HMC) isostere as a transition-state mimetic was designed and synthesized. From the structure-activity relationship studies, potent dipeptide-type inhibitors having high antiviral activity either in the absence or in the presence of 50% human serum were discovered.

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

RX(9) OF 108 ...X + R ==> Y



(9) →



Y

RX(9) RCT X 470697-22-2, R 467446-90-8

STAGE(1)

RGT D 25952-53-8 EDAP, E 2592-95-2 1-Benzotriazolol

SOL 68-12-2 DMF


STAGE(2)

RGT O 7647-01-0 HCl


SOL 123-91-1 Dioxane, 7732-18-5 Water

PRO Y 819083-80-2

RX(10) OF 108 ...Z + R ==> AA



(10) →



AA

RX(10) RCT Z 819083-90-4, R 467446-90-8

STAGE(1)

RGT D 25952-53-8 EDAP, E 2592-95-2 1-Benzotriazolol

SOL 68-12-2 DMF


STAGE(2)

RGT O 7647-01-0 HCl

SOL 123-91-1 Dioxane, 7732-18-5 Water

PRO AA 819083-81-3



RX(24) OF 108 ...AO + AZ ==> X...



RX(24) RCT AO 819083-85-7, AZ 116661-86-0

STAGE(1)
RGT D 25952-53-8 EDAPSTAGE(2)
RGT O 7647-01-0 HCl
SOL 123-91-1 Dioxane, 7732-18-5 Water
PRO X 478697-22-2

RX(25) OF 108 ...AQ + AZ ==> Z...



RX(24) RCT AO 819083-85-7, AZ 116661-86-0

STAGE(1)
RGT D 25952-53-8 EDAPSTAGE(2)
RGT O 7647-01-0 HCl
SOL 123-91-1 Dioxane, 7732-18-5 Water

PRO X 478697-22-2


RX(9) RCT X 478697-22-2, R 467446-90-8

STAGE(1)
RGT D 25952-53-8 EDAP, E 2592-95-2 1-Benzotriazolol
SOL 68-12-2 DMFSTAGE(2)
RGT O 7647-01-0 HCl
SOL 123-91-1 Dioxane, 7732-18-5 Water

PRO Y 819083-80-2


RX(47) OF 108 COMPOSED OF RX(25), RX(10)

RX(47) AQ + AZ + R ==> AA



AQ

AZ




RX(25) RCT AQ 819083-86-8, AZ 116661-86-0

STAGE(1)
RGT D 25952-53-8 EDAPSTAGE(2)
RGT O 7647-01-0 HCl
SOL 123-91-1 Dioxane, 7732-18-5 Water

PRO Z 819083-90-4


RX(46) OF 108 COMPOSED OF RX(24), RX(9)

RX(46) AO + AZ + R ==> Y




R

2 STEPS



R

2 STEPS



AA

RX(25) RCT AQ 819083-86-8, AZ 116661-86-0

STAGE(1)
RGT D 25952-53-8 EDAPSTAGE(2)
RGT O 7647-01-0 HCl
SOL 123-91-1 Dioxane, 7732-18-5 Water

PRO Z 819083-90-4




RX(10) RCT Z 819083-90-4, R 467446-90-8

STAGE(1)
RGT D 25952-53-8 EDAP, E 2592-95-2 1-Benzotriazolol
SOL 68-12-2 DMFSTAGE(2)
RGT O 7647-01-0 HCl
SOL 123-91-1 Dioxane, 7732-18-5 Water

PRO AA 819083-81-3

RX(65) OF 108 COMPOSED OF RX(35), RX(9)

RX(65) BI + BB + X ==> Y



RX(35) RCT BI 15980-22-0, BB 105-36-2

STAGE(1)
RGT BC 584-08-7 K2CO3

STAGE(2)
RGT BD 1310-73-2 NaOH
SOL 7732-18-5 Water

PRO R 467446-90-8

RX(9) RCT X 478697-22-2, R 467446-90-8

STAGE(1)
RGT D 25952-53-8 EDAP, E 2592-95-2 1-Benzotriazolol
SOL 68-12-2 DMF

STAGE(2)

PRO R 467446-90-8

RX(10) RCT Z 819083-90-4, R 467446-90-8



STAGE(1)
RGT D 25952-53-8 EDAP, E 2592-95-2 1-Benzotriazolol
SOL 68-12-2 DMF

STAGE(2)
RGT O 7647-01-0 HCl
SOL 123-91-1 Dioxane, 7732-18-5 Water


PRO AA 819083-81-3

RX(87) OF 108 COMPOSED OF REACTION SEQUENCE RX(35), RX(9) AND REACTION SEQUENCE RX(24), RX(9)

...BI + BB ==> R...
...AO + AZ + R ==> Y






START NEXT REACTION SEQUENCE



RGT O 7647-01-0 HCl
SOL 123-91-1 Dioxane, 7732-18-5 Water

PRO Y 819083-80-2



RX(66) OF 108 COMPOSED OF RX(35), RX(10)
RX(66) BI + BB + Z ==> AA



RX(35) RCT BI 15980-22-0, BB 105-36-2

STAGE(1)
RGT BC 584-08-7 K2CO3

STAGE(2)
RGT BD 1310-73-2 NaOH
SOL 7732-18-5 Water



RX(35) RCT BI 15980-22-0, BB 105-36-2

STAGE(1)
RGT BC 584-08-7 K2CO3

STAGE(2)
RGT BD 1310-73-2 NaOH
SOL 7732-18-5 Water

PRO R 467446-90-8

RX(24) RCT AO 819083-85-7, AZ 116661-86-0

STAGE(1)
RGT D 25952-53-8 EDAP

STAGE(2)
RGT O 7647-01-0 HCl
SOL 123-91-1 Dioxane, 7732-18-5 Water

PRO X 478697-22-2



RX(9) RCT X 478697-22-2, R 467446-90-8

STAGE(1)
RGT D 25952-53-8 EDAP, E 2592-95-2 1-Benzotriazolol
SOL 68-12-2 DMF




STAGE(2)
RGT O 7647-01-0 HCl

SOL 123-91-1 Dioxane, 7732-18-5 Water

PRO Y 819083-80-2

RX(89) OF 108 COMPOSED OF REACTION SEQUENCE RX(35), RX(10)
AND REACTION SEQUENCE RX(25), RX(10)...BI + BB ==> R...
...AQ + AZ + R ==> AA

START NEXT REACTION SEQUENCE



RX(35) RCT BI 15980-22-0, BB 105-36-2

STAGE(1)
ROT BC 584-08-7 K2CO3STAGE(2)
ROT BD 1310-73-2 NaOH
SOL 7732-18-5 Water

PRO R 467446-90-8

RX(25) RCT AQ 819083-86-8, AZ 116661-86-0

STAGE(1)
ROT D 25952-53-8 EDAPSTAGE(2)
ROT O 7647-01-0 HCl
SOL 123-91-1 Dioxane, 7732-18-5 Water

PRO Z 819083-90-4

RX(10) RCT Z 819083-90-4, R 467446-90-8

STAGE(1)
ROT D 25952-53-8 EDAP, E 2592-95-2 1-Benzotriazolol
SOL 68-12-2 DMFSTAGE(2)
ROT O 7647-01-0 HCl
SOL 123-91-1 Dioxane, 7732-18-5 Water

PRO AA 819083-81-3

L12 ANSWER 3 OF 18 CASREACT COPYRIGHT 2007 ACS ON STN
 ACCESSION NUMBER: 140:263738 CASREACT Full-text
 TITLE: Synthesis and opioid activity of N,N-dimethyl-Dmt-Tic-NH-CH(R)-R' analogues: acquisition of potent δ antagonist

AUTHOR(S): Balboni, Gianfranco; Salvadori, Severo; Guerrini, Remo; Negri, Lucia; Giannini, Elisa; Bryant, Sharon D.; Jinmaa, Yunden; Lazarus, Lawrence H.

CORPORATE SOURCE: Department of Toxicology, University of Cagliari, Cagliari, I-09126, Italy

SOURCE: Bioorganic & Medicinal Chemistry (2003), 11(24), 5435-5441

PUBLISHER: CODEN: BMECEP; ISSN: 0968-0896




DOCUMENT TYPE: Elsevier Ltd.

LANGUAGE: English

AB N,N-Dimethylation of the H-Dmt-Tic-NH-CH(R)-R' series of compds. produced no significant effect on the high δ -opioid receptor affinity (K_i =0.035-0.454 nM), but dramatically decreased that for the μ -opioid receptor. The effect of N-methylation was independent of the length of the linker (R); however, the bioactivities were affected by the chemical composition of the third aromatic group (R'): Ph (Ph) (5'-S') elicited a greater reduction in μ -affinity (40-70-fold) compared to analogs containing 1H-benzimidazole-2-yl (Bid) (9-fold). The major consequences of N,N-dimethylation on in vitro bioactivity were: (i) a loss of δ -agonism coupled with the appearance of potent δ antagonism; and (ii) a consistent loss of μ -affinity resulted in enhanced δ -opioid receptor selectivity. With the exception of one compound, the change in the hydrophobic environment at the N-terminus and formation of a tertiary amine by N,N-dimethylation in analogs of the Dmt-Tic pharmacophore produced potent δ -selective antagonists.

REFERENCE COUNT: 36 THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

RX(9) OF 62 ...Z + 2 K ==> AA






RX(9) RCT Z 673461-36-4, K 50-00-0

STAGE(1)
ROT O 64-19-7 AcOH, M 25895-60-7 NaBH3CN, N 109-02-4 N-Methylmorpholine
SOL 7732-18-5 Water, 75-05-8 MeCN
CON SUBSTAGE(1) 10 minutes, room temperature
SUBSTAGE(2) 15 minutes, room temperatureSTAGE(2)
ROT E 76-05-1 F3CCO2H
CON room temperature, acidify

PRO AA 859231-90-6

RX(19) OF 62 ...D + AM ==> AN...






AN: CM 2
YIELD 82%


RX(19) RCT D 99953-00-1, AM 673461-29-5
RGT H 2592-95-2 1-Benzotriazolol, I 25952-53-8 EDAP
PRO AN 673461-31-9
CON room temperature

RX(23) OF 62 ...AN ==> Z...



AN: CM 2

(23)




Z: CM 2
YIELD 97%

RX(23) RCT AN 673461-31-9
RGT E 76-05-1 F3CCO2H

PRO Z 673461-36-4
CON room temperature


RX(34) OF 62 COMPOSED OF RX(18), RX(19)
RX(34) AL + E + D ==> AN




AL

E

D



AN: CM 1
YIELD 82%




AN: CM 2
YIELD 82%

RX(18) RCT AL 673461-28-4, E 76-05-1
PRO AM 673461-29-5
CON room temperature


RX(19) RCT D 99953-00-1, AM 673461-29-5
RGT H 2592-95-2 1-Benzotriazolol, I 25952-53-8 EDAP
PRO AN 673461-31-9
CON room temperature

RX(35) OF 62 COMPOSED OF RX(19), RX(23)
RX(35) D + AN ==> Z




D

AN: CM 1



AN: CM 2

Z: CM 1
YIELD 97%




Z: CM 2
YIELD 97%

RX(19) RCT D 99953-00-1, AM 673461-29-5
RGT H 2592-95-2 1-Benzotriazolol, I 25952-53-8 EDAP
PRO AN 673461-31-9
CON room temperature

RX(23) RCT AN 673461-31-9
RGT E 76-05-1 F3CCO2H
PRO Z 673461-36-4
CON room temperature


RX(39) OF 62 COMPOSED OF RX(23), RX(9)
RX(39) AN + 2 K ==> AA



AN: CM 1

AN: CM 2

H2C=O
2 K
2 STEPS



AA: CM 1
YIELD 89%

AA: CM 2
YIELD 89%

RX(23) RCT AN 673461-31-9
RGT E 76-05-1 F3CCO2H
PRO Z 673461-36-4
CON room temperature




RX(9) RCT Z 673461-36-4, K 50-00-0

STAGE(1)
RGT Q 64-19-7 AcOH, M 25895-60-7 NaBH3CN, N 109-02-4
N-Methylmorpholine
SOL 7732-18-5 Water, 75-05-8 MeCN
CON SUBSTAGE(1) 10 minutes, room temperature
SUBSTAGE(2) 15 minutes, room temperature

STAGE(2)
RGT E 76-05-1 F3CCO2H
CON room temperature, acidify

PRO AA 859231-90-6

RX(47) OF 62 COMPOSED OF RX(18), RX(19), RX(23)
RX(47) AL + E + D ==> Z





RX(18) RCT AL 673461-28-4, E 76-05-1
PRO AM 673461-29-5
CON room temperature

RX(19) RCT D 99953-00-1, AM 673461-29-5
RGT H 2592-95-2 1-Benzotriazolol, I 25952-53-8 EDAP
PRO AN 673461-31-9
CON room temperature

RX(23) RCT AN 673461-31-9
RGT E 76-05-1 F3CCO2H
PRO Z 673461-36-4
CON room temperature

RX(49) OF 62 COMPOSED OF RX(19), RX(23), RX(9)
RX(49) D + AM + 2 K ==> AA



* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(19) RCT D 99953-00-1, AM 673461-29-5
RGT H 2592-95-2 1-Benzotriazolol, I 25952-53-8 EDAP
PRO AN 673461-31-9
CON room temperature

RX(23) RCT AN 673461-31-9
RGT E 76-05-1 F3CCO2H
PRO Z 673461-36-4
CON room temperature

RX(9) RCT Z 673461-36-4, K 50-00-0

STAGE(1)


RGT G 64-19-7 AcOH, M 25895-60-7 NaBH3CN, N 109-02-4
N-Methylmorpholine
SOL 7732-18-5 Water, 75-05-8 MeCN
CON SUBSTAGE(1) 10 minutes, room temperature
SUBSTAGE(2) 15 minutes, room temperature

STAGE(2)

RGT E 76-05-1 F3CCO2H
CON room temperature, acidify

PRO AA 859231-90-6

RX(50) OF 62 COMPOSED OF RX(18), RX(19), RX(23), RX(9)
RX(50) AL + E + D + 2 K ==> AA



* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(18) RCT AL 673461-28-4, E 76-05-1
PRO AM 673461-29-5
CON room temperature

RX(19) RCT D 99953-00-1, AM 673461-29-5
RGT H 2592-95-2 1-Benzotriazolol, I 25952-53-8 EDAP
PRO AN 673461-31-9
CON room temperature

RX(23) RCT AN 673461-31-9
RGT E 76-05-1 F3CCO2H
PRO Z 673461-36-4
CON room temperature

RX(9) RCT Z 673461-36-4, K 50-00-0

STAGE(1)

RGT G 64-19-7 AcOH, M 25895-60-7 NaBH3CN, N 109-02-4
N-Methylmorpholine
SOL 7732-18-5 Water, 75-05-8 MeCN
CON SUBSTAGE(1) 10 minutes, room temperature
SUBSTAGE(2) 15 minutes, room temperature

STAGE(2)

RGT E 76-05-1 F3CCO2H
CON room temperature, acidify

PRO AA 859231-90-6


L12 ANSWER 4 OF 18 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 138:183296 CASREACT Full-text
TITLE: Transcellular Proteolysis Demonstrated by Novel Cell
Surface-associated Substrates of Dipeptidyl Peptidase
IV (CD26)
AUTHOR(S): Lorey, Susan; Faust, Juergen; Mrestani-Klaus, Carmen;

Kaehne, Thilo; Ansorge, Siegfried; Neubert, Klaus;
Buehling, Frank
CORPORATE SOURCE: Institute of Biochemistry, Department of Biochemistry
and Biotechnology, Martin-Luther-University
Halle-Wittenberg, Halle (Saale), D-06120, Germany
SOURCE: Journal of Biological Chemistry (2002), 277(36),
33170-33177
CODEN: JBCHAJ; ISSN: 0021-9258
PUBLISHER: American Society for Biochemistry and Molecular
Biology
DOCUMENT TYPE: Journal
LANGUAGE: English


AB Proteolytic enzymes contribute to the regulation of cellular functions such as cell proliferation and death, cytokine production, and matrix remodeling. Dipeptidyl peptidase IV (DP IV) catalyzes the cleavage of several cytokines and thereby contributes to the regulation of cytokine production and the proliferation of immune cells. Here we show for the first time that cell surface-bound DP IV catalyzes the cleavage of specific substrates that are associated with the cellular surface of neighboring cells. Rhodamine 110 (R110), a highly fluorescent xanthene dye, was used to synthesize dipeptidyl peptidase IV (DP IV/CD26) substrates Gly(Ala)-Pro-R110-R, thus facilitating a stable binding of the fluorescent moiety on the cell surface. The fixation resulted from the interaction with the reactive anchor rhodamine and allowed the quantification of cellular DP IV activity on single cells. The reactivity, length, and hydrophobicity of Rhodamine was characterized as the decisive factor that facilitated the determination of cellular DP IV activity. Using fluorescence microscopy, it was possible to differentiate between different DP IV activities. The hydrolysis of cell-bound substrates Xaa-Pro-R110-R by DP IV of neighboring cells and by soluble DP IV was shown using flow cytometry. These data demonstrate that ectopeptidases such as DP IV may be involved in communication between blood cells via proteolysis of cell-associated substrates.

REFERENCE COUNT: 44 THERE ARE 44 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

RX(3) OF 50 ...C + I ==> J...




(3)


J
YIELD 96%

RX(3) RCT C 498539-64-3, I 108-24-7
PRO J 586961-97-2
SOL 110-86-1 Pyridine
CON 24 hours, room temperature

RX(10) OF 50 ...W + H ==> Y...




W




H

(10) →

Y
YIELD 35%


RX(10) RCT W 101310-84-3, H 498539-65-4
RGT E 25952-53-8 EDAP
PRO Y 586961-92-4
SOL 68-12-2 DMF
CON SUBSTAGE(1) 1 hour, 0 deg C
SUBSTAGE(2) 20 hours, room temperature
SUBSTAGE(3) 6 hours, room temperature

RX(13) OF 50 ...M ==> AD



M

(13) →




AD

RX(13) RCT M 586961-19-5
RGT AE 76-05-1 F3CCO2H
PRO AD 498539-66-5


SOL 75-09-2 CH2Cl2
CON 3 hours, room temperature

RX(14) OF 50 ...O ==> AG



O


(14) →



AG


RX(14) RCT O 586961-24-2
RGT AE 76-05-1 F3CCO2H
PRO AG 498539-67-6
SOL 75-09-2 CH2Cl2
CON 3 hours, room temperature

RX(15) OF 50 AH ==> AI



AH


(15) →



AI


RX(15) RCT AH 586961-51-5
RGT AE 76-05-1 F3CCO2H
PRO AI 498539-68-7
SOL 75-09-2 CH2Cl2
CON 3 hours, room temperature

RX(16) OF 50 ...R ==> AJ



R


(16)




AJ

RX(16) RCT R 586961-54-0
 RGT AR 76-05-1 F3CCO2H
 PRO AJ 498539-69-8
 SOL 75-09-2 CH2Cl2
 CON 3 hours, room temperature

RX(17) OF 50 AK + S ==> T




AK



(17)


RX(18) RCT AA 586961-80-0
 RGT AR 76-05-1 F3CCO2H
 PRO AL 498539-71-2
 SOL 75-09-2 CH2Cl2
 CON 3 hours, room temperature

RX(19) OF 50 ...AC ==> AM



AC


(19)



AM

RX(19) RCT AC 586961-89-9
 RGT AR 76-05-1 F3CCO2H
 PRO AM 498539-72-3
 SOL 75-09-2 CH2Cl2
 CON 3 hours, room temperature


RX(20) OF 50 ...Y ==> AN



T


RX(17) RCT AK 586961-74-2, S 876-08-4
 RGT AR 76-05-1 F3CCO2H
 PRO T 498539-70-1
 SOL 75-09-2 CH2Cl2
 CON 3 hours, room temperature

RX(18) OF 50 ...AA ==> AL




AA

(18)




AL



Y


(20)



AN


RX(20) RCT Y 586961-92-4
 RGT AO 7647-01-0 HCl, AP 64-19-7 AcOH
 PRO AN 498539-73-4
 CON 30 minutes, room temperature

RX(21) OF 50 ...J ==> AQ



J


(21)



AQ

RX (21) RCT J 586962-97-2
 RGT AO 7647-01-0 HCl, AP 64-19-7 AcOH
 PRO AQ 498539-74-5
 CON 30 minutes, room temperature

RX (31) OF 50 COMPOSED OF RX (3), RX (21)
 RX (31) C + I ==> AQ




C



I

2
 STEPS



AQ


STAGE (2)

RCT L 79-04-9
 CON SUBSTAGE (1) 1 hour, 4 deg C
 SUBSTAGE (2) 1 hour, room temperature


PRO M 586961-19-5

RX (13) RCT M 586961-19-5
 RGT AR 76-05-1 F3CCO2H
 PRO AD 498539-66-5
 SOL 75-09-2 CH2Cl2
 CON 3 hours, room temperature

RX (33) OF 50 COMPOSED OF RX (5), RX (14)
 RX (33) C + N ==> AG




C



N

2
 STEPS




AG

RX (5) RCT C 498539-64-3


RX (3) RCT C 498539-64-3, I 108-24-7
 PRO J 586962-97-2
 SOL 110-86-1 Pyridine
 CON 24 hours, room temperature

RX (21) RCT J 586962-97-2
 RGT AO 7647-01-0 HCl, AP 64-19-7 AcOH
 PRO AQ 498539-74-5
 CON 30 minutes, room temperature

RX (32) OF 50 COMPOSED OF RX (4), RX (13)
 RX (32) C + L ==> AD




C



L

2
 STEPS



AD

RX (4) RCT C 498539-64-3
 STAGE (1)
 SOL 68-12-2 DMF
 CON room temperature -> 4 deg C

STAGE (1)

SOL 68-12-2 DMF
 CON room temperature -> 4 deg C


STAGE (2)

RCT N 1575-61-7
 CON SUBSTAGE (1) 1 hour, 4 deg C
 SUBSTAGE (2) 1 hour, room temperature


PRO O 586961-24-2

RX (14) RCT O 586961-24-2
 RGT AR 76-05-1 F3CCO2H
 PRO AG 498539-67-6
 SOL 75-09-2 CH2Cl2
 CON 3 hours, room temperature

RX (34) OF 50 COMPOSED OF RX (6), RX (14)
 RX (34) C + P ==> AG




C



P

2
 STEPS



AG

RX(6) RCT C 498539-64-3

STAGE(1)

SOL 68-12-2 DMF

CON room temperature -> 4 deg C

STAGE(2)

RCT P 4635-59-0

CON SUBSTAGE(1) 1 hour, 4 deg C

SUBSTAGE(2) 1 hour, room temperature

PRO O 586961-24-2

RX(14) RCT O 586961-24-2

RGT AR 76-05-1 F3CCO2H


PRO AQ 498539-67-6

SOL 75-09-2 CH2Cl2


CON 3 hours, room temperature

RX(35) OF 50 COMPOSED OF RX(7), RX(16)


RX(35) C + Q ==> AJ



C



2 STEPS



AJ

RX(7) RCT C 498539-64-3

STAGE(1)

SOL 68-12-2 DMF

CON room temperature -> 4 deg C

STAGE(2)

RCT Q 4509-90-4

CON SUBSTAGE(1) 1 hour, 4 deg C

SUBSTAGE(2) 1 hour, room temperature

PRO R 586961-54-8

RX(16) RCT R 586961-54-8

RGT AR 76-05-1 F3CCO2H


PRO AJ 498539-69-8

SOL 75-09-2 CH2Cl2


CON 3 hours, room temperature

RX(36) OF 50 COMPOSED OF RX(9), RX(10)


RX(36) U + V + H ==> Y



U




V



H

2 STEPS



Y YIELD 35%

RX(9) RCT U 1132-17-4, V 60-32-2

PRO W 101310-84-3

SOL 109-99-9 THF

CON 3 hours, reflux

RX(10) RCT W 101310-84-3, H 498539-65-4

RGT E 25952-53-8 EDAP

PRO Y 586961-92-4

SOL 68-12-2 DMF


CON SUBSTAGE(1) 1 hour, 0 deg C

SUBSTAGE(2) 20 hours, room temperature


SUBSTAGE(3) 6 hours, room temperature

RX(37) OF 50 COMPOSED OF RX(10), RX(20)

RX(37) W + H ==> AN




W



H

2 STEPS



AN

RX(10) RCT W 101310-84-3, H 498539-65-4

RGT E 25952-53-8 EDAP

PRO Y 586961-92-4

SOL 68-12-2 DMF

CON SUBSTAGE(1) 1 hour, 0 deg C

SUBSTAGE(2) 20 hours, room temperature

SUBSTAGE(3) 6 hours, room temperature

RX(20) RCT Y 586961-92-4


RGT AQ 7647-01-0 HCl, AP 64-19-7 AcOH

PRO AN 498539-73-4


CON 30 minutes, room temperature

RX(38) OF 50 COMPOSED OF RX(11), RX(18)

RX(38) Z + H ==> AL




Z



H

2 STEPS



AL

RX(11) RCT Z 116965-29-8

STAGE(1)

SOL 68-12-2 DMF

CON room temperature -> 4 deg C

STAGE(2)

RCT H 498539-65-4

ROT D 100-74-3 4-Ethylmorpholine, E 25952-53-8 EDAP

CON SUBSTAGE(1) 1 hour, 4 deg C

SUBSTAGE(2) 6 hours, room temperature

PRO AA 586961-80-0

RX(10) RCT AA 586961-80-0

ROT AR 76-05-1 F3CCO2H


PRO AL 498539-71-2

SOL 75-09-2 CH2Cl2


CON 3 hours, room temperature

RX(39) OF 50 COMPOSED OF RX(12), RX(19)

RX(39) AB + H ==> AM



AB

2
STEPS

AM

RX(12) RCT AB 82333-93-5

STAGE(1)

SOL 68-12-2 DMF

CON room temperature -> 4 deg C

STAGE(2)

RCT H 498539-65-4

ROT D 100-74-3 4-Ethylmorpholine, E 25952-53-8 EDAP

CON SUBSTAGE(1) 1 hour, 4 deg C

SUBSTAGE(2) 6 hours, room temperature

PRO AC 586961-89-9

RX(19) RCT AC 586961-89-9

ROT AR 76-05-1 F3CCO2H


PRO AM 498539-72-3

SOL 75-09-2 CH2Cl2


CON 3 hours, room temperature

RX(49) OF 50 COMPOSED OF RX(9), RX(10), RX(20)


RX(49) U + V + H ==> AN




U



V



H

3
STEPS

AN

RX(9) RCT U 1122-17-4, V 60-32-2

PRO W 101310-84-3

SOL 109-99-9 THF

CON 3 hours, reflux

RX(10) RCT W 101310-84-3, H 498539-65-4

ROT E 25952-53-8 EDAP

PRO Y 586961-92-4

SOL 68-12-2 DMF

CON SUBSTAGE(1) 1 hour, 0 deg C

SUBSTAGE(2) 20 hours, room temperature

SUBSTAGE(3) 6 hours, room temperature

RX(20) RCT Y 586961-92-4

ROT AO 7647-01-0 HCl, AP 64-19-7 AcOH

PRO AN 490539-73-4

CON 30 minutes, room temperature

L12 ANSWER 5 OF 18 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER:

136:247867 CASREACT

TITLE:

Full-text: Polymer-assisted solution-phase parallel synthesis of dipeptide p-nitroanilides and dipeptide diphenyl phosphonates

AUTHOR(S):

Senten, Kristel; Van der Veken, Pieter; Bal, Gunther; Haemers, Achiel; Augustyns, Koen

CORPORATE SOURCE:


Department of Medicinal Chemistry, University of

Antwerp (UIA), Antwerp, B-2610, Belg.
 SOURCE: Tetrahedron Letters (2001), 42(52), 9135-9138
 CODEN: TETRAY; ISSN: 0040-4039
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English


AB This letter describes the parallel synthesis of dipeptide p-nitroanilides and dipeptide di-Ph phosphonates, compds. that can be used as substrates and irreversible inhibitors for the rapid profiling of dipeptidyl peptidases. A polymer-assisted solution-phase synthesis was used for a rapid and clean coupling between easily available building blocks.

REFERENCE COUNT: 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

RX(1) OF 35 A + B ==> C




A



B

(1)

C
YIELD 98%

RX(1) RCT A 132388-68-2

STAGE(1)

ROT D 2592-95-2 1-Benzotriazolol, E 538-75-0 DCC

SOL 75-09-2 CH2Cl2

STAGE(2)

RCT B 7369-91-7

SOL 75-09-2 CH2Cl2

STAGE(3)


ROT F 76-05-1 F3CCO2H

SOL 75-09-2 CH2Cl2



PRO C 90145-75-8

NTE polymer-assisted solution-phase synthesis, solid-supported reagent, methylpolyethylene resin used

RX(2) OF 35 H + B ==> I



(2) →

I
YIELD 87%

RX(2) RCT H 1676-90-0

STAGE(1)

RGT D 2592-95-2 1-Benzotriazolol, E 538-75-0 DCC
SOL 75-09-2 CH2Cl2

STAGE(2)

RCT B 7369-91-7
SOL 75-09-2 CH2Cl2



STAGE(3)

RGT F 76-05-1 F3CCO2H
SOL 75-09-2 CH2Cl2

PRO I 60189-48-2

NTE polymer-assisted solution-phase synthesis, solid-supported reagent, methylpolystyrene resin used

RX(3) OF 35 J + B ==> K

K
YIELD 42%

RX(3) RCT J 20866-46-0

STAGE(1)

RGT D 2592-95-2 1-Benzotriazolol, E 538-75-0 DCC
SOL 75-09-2 CH2Cl2

STAGE(2)

RCT B 7369-91-7
SOL 75-09-2 CH2Cl2


STAGE(3)

RGT F 76-05-1 F3CCO2H
SOL 75-09-2 CH2Cl2



PRO K 99264-68-3

NTE polymer-assisted solution-phase synthesis, solid-supported reagent, methylpolystyrene resin used

RX(4) OF 35 L + B ==> M



(4) →

H
YIELD 10%

RX(4) RCT L 13139-16-7

STAGE(1)

RGT D 2592-95-2 1-Benzotriazolol, E 538-75-0 DCC
SOL 75-09-2 CH2Cl2

STAGE(2)

RCT B 7369-91-7
SOL 75-09-2 CH2Cl2


STAGE(3)

RGT F 76-05-1 F3CCO2H
SOL 75-09-2 CH2Cl2


PRO M 90145-77-0

NTE polymer-assisted solution-phase synthesis, solid-supported reagent, methylpolystyrene resin used

RX(5) OF 35 N + B ==> O



(5) →

O
YIELD 16%

RX(5) RCT N 13734-34-4

STAGE(1)

RGT D 2592-95-2 1-Benzotriazolol, E 538-75-0 DCC
SOL 75-09-2 CH2Cl2

STAGE(2)

RCT B 7369-91-7
SOL 75-09-2 CH2Cl2


STAGE(3)

RGT F 76-05-1 F3CCO2H
SOL 75-09-2 CH2Cl2


PRO O 90145-72-5

NTE polymer-assisted solution-phase synthesis, solid-supported reagent, methylpolystyrene resin used

RX(7) OF 35 R + B ==> S



(7) →



RX(7) RCT R 13734-39-8

STAGE(1)

ROT D 2592-95-2 1-Benzotriazolol, R 538-75-0 DCC
SOL 75-09-2 CH2Cl2

STAGE(2)

RCT B 7369-91-7
SOL 75-09-2 CH2Cl2



STAGE(3)

ROT F 76-05-1 F3CCO2H
SOL 75-09-2 CH2Cl2

PRO S 90145-70-3

NTE polymer-assisted solution-phase synthesis, solid-supported reagent, methylpolystyrene resin used

RX(8) OF 35 T + B ==> U



RX(8) RCT T 47375-34-8

STAGE(1)

ROT D 2592-95-2 1-Benzotriazolol, R 538-75-0 DCC
SOL 75-09-2 CH2Cl2

STAGE(2)

RCT B 7369-91-7
SOL 75-09-2 CH2Cl2



STAGE(3)

ROT F 76-05-1 F3CCO2H
SOL 75-09-2 CH2Cl2

PRO U 90145-69-0

NTE polymer-assisted solution-phase synthesis, solid-supported reagent, methylpolystyrene resin used

RX(9) OF 35 V + B ==> W



RX(9) RCT V 13734-41-3

STAGE(1)

ROT D 2592-95-2 1-Benzotriazolol, R 538-75-0 DCC
SOL 75-09-2 CH2Cl2

STAGE(2)

RCT B 7369-91-7
SOL 75-09-2 CH2Cl2

STAGE(3)

ROT F 76-05-1 F3CCO2H
SOL 75-09-2 CH2Cl2


PRO W 90145-74-7

NTE polymer-assisted solution-phase synthesis, solid-supported reagent, methylpolystyrene resin used

L12 ANSWER 6 OF 18 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 135:344726 CASREACT Full-text
TITLE: Process for preparation of substituted aspartic acid
inventor(s): Wannamaker, Marion W.; Forster, Cornelia
PATENT ASSIGNEE(S): Vertex Pharmaceuticals Incorporated, USA
SOURCE: PCT Int. Appl., 59 pp.
CODEN: PXXXX2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001081330	A2	20011101	WO 2001-0812769	20010419
WO 2001081330	A3	20020307		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW			
RW:	OH, OM, OS, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IS, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CP, CO, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
CA 2402128	A1	20011101	CA 2001-2402128	20010419
EP 1278737	A2	20030129	EP 2001-927217	20010419
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
ZA 2002006658	A	20040428	ZA 2002-6658	20020820
BO 107028	A	20030630	BO 2002-107028	20020822
BO 2002004111	A	20020828	NO 2002-4111	20020828
US 2003119899	A1	20030626	US 2003-239981	20020828
US 7109357	B2	20060919		
BR 2002004426	A	20040608	BR 2002-204426	20021029
PRIORITY APPLN. INFO.:			US 2000-199329P	20000424
			WO 2001-0812769	20010419

OTHER SOURCE(S): MARPAT 135:344726
OI




AB Title compds. [I; R1 = (substituted) aliphatic, aralkyl, heterocyclylalkyl, aryl; R2 = organic radical, preferably a P2-P4 moiety of a caspase inhibitor], were prepared by treatment of butenolactones (II; R1 as above) with N3Y (Y = H, silyl, counterion) to give azidolactones (III; R1 as above) followed by conversion of III to aminolactones (IV) or iminophosphoranes (V; R undefined) and coupling of either with R2CO2H (R2 as above) or a reactive equivalent thereof. Thus, Me3SiN3, HOAc, DBU, and 5-ethoxy-5H-furan-2-one were stirred 24 h in CH2Cl2 to give 73% 4-azido-5-ethoxydihydrofuran-2-one. The latter with (S)-pyrrolidine-1,2-dicarboxylic acid 1-tert-Bu ester was hydrogenated in EtOAc over Pd/C under 1 atm H2 for 1 h; the mixture was diluted with CH2Cl2, filtered, and evaporated. The crude mixture was stirred with diisopropylethylamine, EDC, and HOBT in CH2Cl2 for 24 h to give 56% (R)-2-(2-ethoxy-5-oxo-tetrahydrofuran-3-ylcarbamoyl)pyrrolidine-1-carboxylic acid tert-Bu ester.

RX(11) OF 28 ...AH + AN ==> AK...



(11)



AK
YIELD 59%

RX(11) RCT AH 370877-09-1

STAGE(1)

RGT AO 108-48-5 2,6-Lutidine
SOL 75-09-2 CH₂Cl₂

STAGE(2)

RGT AP 27607-77-8 Me₃SiSO₃CF₃

STAGE(3)

RGT AQ 144-55-8 NaHCO₃

STAGE(4)

SOL 75-09-2 CH₂Cl₂

STAGE(5)

RCT AN 68222-59-3
RGT W 25952-53-8 EDAP, X 2592-95-2 1-Benzotriazolol
SOL 75-09-2 CH₂Cl₂


STAGE(6)

SOL 141-78-6 AcOEt

PRO AK 371126-01-1



RX(20) OF 28 COMPOSED OF RX(11), RX(10)

RX(20) AH + AN + AL ==> AM



AH

AN



AM
YIELD 62%

RX(11) RCT AH 370877-09-1

STAGE(1)

RGT AO 108-48-5 2,6-Lutidine
SOL 75-09-2 CH₂Cl₂

STAGE(2)

RGT AP 27607-77-8 Me₃SiSO₃CF₃

STAGE(3)

RGT AQ 144-55-8 NaHCO₃

STAGE(4)

SOL 75-09-2 CH₂Cl₂

STAGE(5)

RCT AN 68222-59-3
RGT W 25952-53-8 EDAP, X 2592-95-2 1-Benzotriazolol
SOL 75-09-2 CH₂Cl₂

STAGE(6)

SOL 141-78-6 AcOEt

PRO AK 371126-01-1

RX(10) RCT AK 371126-01-1

STAGE(1)

RGT K 1333-74-0 H₂
CAT 7440-05-3 Pd
SOL 141-78-6 AcOEt

STAGE(2)

SOL 75-09-2 CH₂Cl₂

STAGE(3)

RCT AL 2486-71-7

RGT O 7087-68-5 EtN(Pr-1)₂, W 25952-53-8 EDAP

STAGE(4)

SOL 141-78-6 AcOEt

PRO AM 371126-03-3

L12 ANSWER 7 OF 18 CASREACT COPYRIGHT 2007 ACS ON STN

ACCESSION NUMBER: 135:195782 CASREACT Full-text

TITLE: Solid-Phase Synthesis of Peptidomimetic Inhibitors for the Hepatitis C Virus NS3 Protease

AUTHOR(S): Poupard, Marc-Andre; Cameron, Dale R.; Chabot, Catherine; Ghio, Elise; Goudreau, Nathalie; Goulet, Sylvie; Poirier, Martin; Tsantrizos, Youla S. Department of Chemistry, Boehringer Ingelheim (Canada) Ltd., QC, H7S 2G5, Can.

SOURCE: Journal of Organic Chemistry (2001), 66(14), 4743-4751

PUBLISHER: CODEN: JOCEAH; ISSN: 0022-3263


DOCUMENT TYPE: American Chemical Society

LANGUAGE: English

AB The NS3 serine protease enzyme of the hepatitis C virus (HCV) is essential for viral replication. Short peptides mimicking the N-terminal substrate cleavage products of the NS3 protease are known to act as weak inhibitors of the enzyme and have been used as templates for the design of peptidomimetic inhibitors. Automated solid-phase synthesis of a small library of compds. based on such a peptidomimetic scaffold has led to the identification of potent and highly selective inhibitors of the NS3 protease enzyme.


REFERENCE COUNT: 42 THERE ARE 42 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

RX(1) OF 32 A + B ==> C



A

B



C
YIELD 45%

RX(1) RCT A 357292-85-4

STAGE(1)

RGT D 603-35-0 PPh₃, E 2446-83-5 N₂(CO₂CHMe₂)₂
SOL 75-09-2 CH₂Cl₂, 109-99-9 THF

STAGE(2)

RCT B 108-95-2
SOL 109-99-9 THF


STAGE(3)

RGT F 76-05-1 F₃CCO₂H
SOL 75-09-2 CH₂Cl₂

PRO C 357292-85-8

NTS stereoselective, solid-supported reactant, Wang resin used


RX(2) OF 32 A + I ==> J



A

I

(2)



J
YIELD 50%

RX(2) RCT A 357292-85-4

STAGE(1)

RGT D 603-35-0 PPh₃, E 2446-83-5 N₂(CO₂CHMe₂)₂
SOL 75-09-2 CH₂Cl₂, 109-99-9 THF

STAGE(2)

RCT 1 371-41-5
SOL 109-99-9 THF



STAGE(3)

RGT F 76-05-1 F3CCO2H
SOL 75-09-2 CH2Cl2

PRO J 357292-90-1

NTE stereoselective, solid-supported reactant, Wang resin used

RX(3) OF 32 A + K ==> L



RX(3) RCT A 357292-85-4

STAGE(1)

RGT D 603-35-0 PPh3, E 2446-83-5 N2(CO2CHMe2)2
SOL 75-09-2 CH2Cl2, 109-99-9 THF

STAGE(2)



RCT K 106-48-9
SOL 109-99-9 THF

STAGE(3)

RGT F 76-05-1 F3CCO2H
SOL 75-09-2 CH2Cl2

PRO L 357292-91-2

NTE stereoselective, solid-supported reactant, Wang resin used



RX(5) RCT A 357292-85-4

STAGE(1)

RGT D 603-35-0 PPh3, E 2446-83-5 N2(CO2CHMe2)2
SOL 75-09-2 CH2Cl2, 109-99-9 THF

STAGE(2)

RCT O 540-38-5
SOL 109-99-9 THF

STAGE(3)



RGT F 76-05-1 F3CCO2H
SOL 75-09-2 CH2Cl2

PRO P 357292-92-3

NTE stereoselective, solid-supported reactant, Wang resin used

RX(6) OF 32 A + Q ==> R

RX(4) OF 32 A + M ==> N



RX(4) RCT A 357292-85-4

STAGE(1)

RGT D 603-35-0 PPh3, E 2446-83-5 N2(CO2CHMe2)2
SOL 75-09-2 CH2Cl2, 109-99-9 THF

STAGE(2)

RCT M 106-41-2
SOL 109-99-9 THF



STAGE(3)

RGT F 76-05-1 F3CCO2H
SOL 75-09-2 CH2Cl2

PRO N 357292-86-5

NTE stereoselective, solid-supported reactant, Wang resin used

RX(5) OF 32 A + O ==> P



RX(6) RCT A 357292-85-4

STAGE(1)

RGT D 603-35-0 PPh3, E 2446-83-5 N2(CO2CHMe2)2
SOL 75-09-2 CH2Cl2, 109-99-9 THF

STAGE(2)

RCT Q 106-53-6
SOL 109-99-9 THF



STAGE(3)

RGT F 76-05-1 F3CCO2H
SOL 75-09-2 CH2Cl2

PRO R 357292-93-4

NTE stereoselective, solid-supported reactant, Wang resin used

RX(7) OF 32 A + S ==> T



YIELD 53%

RX(7) RCT A 357292-95-4

STAGE(1)

RGT D 603-35-0 PPh3, E 2446-83-5 N2(CO2CHMe2)2
SOL 75-09-2 CH2Cl2, 109-99-9 THF

STAGE(2)

RCT S 591-20-8
SOL 109-99-9 THF



STAGE(3)

RGT F 76-05-1 F3CCO2H
SOL 75-09-2 CH2Cl2

PRO T 357292-87-6

NTE stereoselective, solid-supported reactant, Wang resin used

RX(8) OF 32 A + U ==> V



YIELD 82%

RX(8) RCT A 357292-85-4

STAGE(1)

RGT D 603-35-0 PPh3, E 2446-83-5 N2(CO2CHMe2)2
SOL 75-09-2 CH2Cl2, 109-99-9 THF

STAGE(2)

RCT U 95-56-7
SOL 109-99-9 THF



STAGE(3)

RGT F 76-05-1 F3CCO2H
SOL 75-09-2 CH2Cl2

PRO V 357292-88-7

NTE stereoselective, solid-supported reactant, Wang resin used

RX(9) OF 32 A + W ==> X



YIELD 30%

RX(9) RCT A 357292-85-4

STAGE(1)

RGT D 603-35-0 PPh3, E 2446-83-5 N2(CO2CHMe2)2
SOL 75-09-2 CH2Cl2, 109-99-9 THF

STAGE(2)

RCT W 92-69-3
SOL 109-99-9 THF



STAGE(3)

RGT F 76-05-1 F3CCO2H
SOL 75-09-2 CH2Cl2

PRO X 357292-94-5

NTE stereoselective, solid-supported reactant, Wang resin used

RX(10) OF 32 A + M + Y ==> Z



YIELD 23%

RX(10) RCT A 357292-85-4

STAGE(1)

RGT D 603-35-0 PPh3, E 2446-83-5 N2(CO2CHMe2)2
SOL 75-09-2 CH2Cl2, 109-99-9 THF

STAGE(2)

RCT M 106-41-2
SOL 109-99-9 THF

STAGE(3)

RCT Y 5720-07-0
RGT AA 497-19-8 Na2CO3
CAT 14221-01-3 Pd(PPh3)4
SOL 110-71-4 (CH2OMe)2

STAGE(4)



RGT F 76-05-1 F3CCO2H

SOL 75-09-2 CH₂Cl₂

PRO Z 357292-95-6

NTE stereoselective, solid-supported reactant, Wang resin used

RX(11) OF 32 A + S + AD ==> AE



RX(11) RCT A 357292-85-4

STAGE(1)

RGT D 603-35-0 PPh₃, E 2446-83-5 N₂(CO₂CHMe₂)₂
SOL 75-09-2 CH₂Cl₂, 109-99-9 THF

STAGE(2)

RCT S 591-20-8
SOL 109-99-9 THF

STAGE(3)

RCT AD 98-80-6
RGT AA 497-19-8 Na₂CO₃
CAT 14221-01-3 Pd(PPh₃)₄
SOL 110-71-4 (CH₂OMe)₂

STAGE(4)

RGT F 76-05-1 F₃CCO₂H
SOL 75-09-2 CH₂Cl₂

PRO AE 357292-96-7

NTE stereoselective, solid-supported reactant, Wang resin used

CAT 14221-01-3 Pd(PPh₃)₄
SOL 110-71-4 (CH₂OMe)₂




STAGE(4)

RGT F 76-05-1 F₃CCO₂H
SOL 75-09-2 CH₂Cl₂

PRO AG 357292-97-8

NTE stereoselective, solid-supported reactant, Wang resin used

RX(13) OF 32 A + S + AH ==> AI






RX(13) RCT A 357292-85-4

STAGE(1)

RGT D 603-35-0 PPh₃, E 2446-83-5 N₂(CO₂CHMe₂)₂
SOL 75-09-2 CH₂Cl₂, 109-99-9 THF

RX(12) OF 32 A + S + AF ==> AG



RX(12) RCT A 357292-85-4

STAGE(1)

RGT D 603-35-0 PPh₃, E 2446-83-5 N₂(CO₂CHMe₂)₂
SOL 75-09-2 CH₂Cl₂, 109-99-9 THF

STAGE(2)

RCT S 591-20-8
SOL 109-99-9 THF

STAGE(3)

RCT AF 13331-27-6
RGT AA 497-19-8 Na₂CO₃

STAGE(2)

RCT S 591-20-8
SOL 109-99-9 THF

STAGE(3)

RCT AH 78887-39-5
RGT AA 497-19-8 Na₂CO₃
CAT 14221-01-3 Pd(PPh₃)₄
SOL 110-71-4 (CH₂OMe)₂



STAGE(4)

RGT F 76-05-1 F₃CCO₂H
SOL 75-09-2 CH₂Cl₂

PRO AI 357292-98-9

NTE stereoselective, solid-supported reactant, Wang resin used

RX(14) OF 32 A + U + AD ==> AJ



RX(14) RCT A 357292-85-4

STAGE(1)

RGT D 603-35-0 PPh₃, E 2446-83-5 N₂(CO₂CHMe₂)₂
SOL 75-09-2 CH₂Cl₂, 109-99-9 THF

STAGE(2)

RCT U 95-56-7

SOL 109-99-9 THF

STAGE(3)

RCT AD 98-80-6

RGT AA 497-19-8 Na2CO3

CAT 14221-01-3 Pd(PPh3)4

SOL 110-71-4 (CH2OMe)2

STAGE(4)



RGT F 76-05-1 F3CCO2H

SOL 75-09-2 CH2Cl2

PRO AJ 357292-99-0

NTE stereoselective, solid-supported reactant, Wang resin used

RX(15) OF 32 A + U + Y ==> AK



* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(15) RCT A 357292-85-4

STAGE(1)

RGT D 603-35-0 PPh3, E 2446-83-5 N2(CO2CHMe2)2

SOL 75-09-2 CH2Cl2, 109-99-9 THF

STAGE(2)

RCT U 95-56-7

SOL 109-99-9 THF

STAGE(3)

RCT Y 5720-07-0

RGT AA 497-19-8 Na2CO3

CAT 14221-01-3 Pd(PPh3)4

SOL 110-71-4 (CH2OMe)2

STAGE(4)



RGT F 76-05-1 F3CCO2H

SOL 75-09-2 CH2Cl2

PRO AK 357293-00-6

NTE stereoselective, solid-supported reactant, Wang resin used

RX(16) OF 32 A + U + AL ==> AM



* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(16) RCT A 357292-85-4

STAGE(1)

RGT D 603-35-0 PPh3, E 2446-83-5 N2(CO2CHMe2)2

SOL 75-09-2 CH2Cl2, 109-99-9 THF

STAGE(2)

RCT U 95-56-7

SOL 109-99-9 THF

STAGE(3)

RCT AL 14047-39-1

RGT AA 497-19-8 Na2CO3

CAT 14221-01-3 Pd(PPh3)4

SOL 110-71-4 (CH2OMe)2

STAGE(4)



RGT F 76-05-1 F3CCO2H

SOL 75-09-2 CH2Cl2

PRO AM 357293-01-7

NTE stereoselective, solid-supported reactant, Wang resin used

RX(17) OF 32 A + AN + AO ==> AP

AP
YIELD 40%

RX(17) RCT A 357292-85-4

STAGE(1)

RGT D 603-35-0 PPh3, E 2446-83-5 N2(CO2CHMe2)2

SOL 75-09-2 CH2Cl2, 109-99-9 THF

STAGE(2)

RCT AN 6602-32-0

SOL 109-99-9 THF

STAGE(3)

RCT AO 6165-69-1

RGT AA 497-19-8 Na2CO3

CAT 14221-01-3 Pd(PPh3)4

SOL 110-71-4 (CH2OMe)2

STAGE(4)



RGT F 76-05-1 F3CCO2H

SOL 75-09-2 CH2Cl2

PRO AP 357293-02-8

NTE stereoselective, solid-supported reactant, Wang resin used

RX(18) OF 32 A + AQ ==> AR

AR
YIELD 23%

RX(18) RCT A 357292-85-4

STAGE(1)

RGT D 603-35-0 PPh3, E 2446-83-5 N2(CO2CHMe2)2

SOL 75-09-2 CH2Cl2, 109-99-9 THF

STAGE(2)

RCT AQ 27292-49-5

SOL 109-99-9 THF

STAGE(3)


RGT F 76-05-1 F3CCO2H

SOL 75-09-2 CH2Cl2

PRO AR 357293-03-9

NTE stereoselective, solid-supported reactant, Wang resin used

RX(19) OF 32 A + AS ==> AT



AT
YIELD 60%

RX (19) RCT A 357292-85-4

STAGE (1)

RGT D 603-35-0 PPh₃, E 2446-83-5 N₂(CO₂CHMe₂)₂
SOL 75-09-2 CH₂Cl₂, 109-99-9 THF

STAGE (2)

RCT AS 934-34-9
SOL 109-99-9 THF


STAGE (3)

RGT F 76-05-1 F₃CCO₂H
SOL 75-09-2 CH₂Cl₂

PRO AT 357293-04-0

NTS stereoselective, solid-supported reactant, Wang resin used

RX (20) OF 32 A + AU ==> AV



AV
YIELD 77%

RX (20) RCT A 357292-85-4

STAGE (1)

RGT D 603-35-0 PPh₃, E 2446-83-5 N₂(CO₂CHMe₂)₂
SOL 75-09-2 CH₂Cl₂, 109-99-9 THF

STAGE (2)

RCT AU 149-30-4
SOL 109-99-9 THF


STAGE (3)

RGT F 76-05-1 F₃CCO₂H
SOL 75-09-2 CH₂Cl₂

PRO AV 357293-05-1

NTS stereoselective, solid-supported reactant, Wang resin used

RX (21) OF 32 A + AW ==> AX



AX
YIELD 26%

RX (21) RCT A 357292-85-4

STAGE (1)

RGT D 603-35-0 PPh₃, E 2446-83-5 N₂(CO₂CHMe₂)₂
SOL 75-09-2 CH₂Cl₂, 109-99-9 THF

STAGE (2)

RCT AW 626-64-2
SOL 109-99-9 THF


STAGE (3)

RGT F 76-05-1 F₃CCO₂H
SOL 75-09-2 CH₂Cl₂

PRO AX 357293-06-2

NTS stereoselective, solid-supported reactant, Wang resin used

RX (22) OF 32 A + AY ==> AZ



AZ
YIELD 16%

RX (22) RCT A 357292-85-4

STAGE (1)

RGT D 603-35-0 PPh₃, E 2446-83-5 N₂(CO₂CHMe₂)₂
SOL 75-09-2 CH₂Cl₂, 109-99-9 THF

STAGE (2)

RCT AY 109-00-2
SOL 109-99-9 THF


STAGE (3)

RGT F 76-05-1 F₃CCO₂H
SOL 75-09-2 CH₂Cl₂

PRO AZ 357293-07-3

NTS stereoselective, solid-supported reactant, Wang resin used

RX (23) OF 32 A + BA ==> BB



BB
YIELD 21%

RX(23) RCT A 357292-05-4

STAGE(1)

RGT D 603-35-0 PPh3, E 2446-83-5 N2(CO2CHMe2)2
SOL 75-09-2 CH2Cl2, 109-99-9 THF

STAGE(2)

RCT BA 148-24-3
SOL 109-99-9 THF


STAGE(3)

RGT F 76-05-1 F3CCO2H
SOL 75-09-2 CH2Cl2

PRO BB 357293-08-4

NTE stereoselective, solid-supported reactant, Wang resin used

RX(24) OF 32 A + BC ==> BD



BD
YIELD 50%

RX(24) RCT A 357292-05-4

STAGE(1)

RGT D 603-35-0 PPh3, E 2446-83-5 N2(CO2CHMe2)2
SOL 75-09-2 CH2Cl2, 109-99-9 THF

STAGE(2)

RCT BC 130-16-5
SOL 109-99-9 THF


STAGE(3)

RGT F 76-05-1 F3CCO2H
SOL 75-09-2 CH2Cl2

PRO BD 357293-09-5

NTE stereoselective, solid-supported reactant, Wang resin used

RX(25) OF 32 A + BE ==> BF



BF
YIELD 41%

RX(25) RCT A 357292-05-4

STAGE(1)

RGT D 603-35-0 PPh3, E 2446-83-5 N2(CO2CHMe2)2
SOL 75-09-2 CH2Cl2, 109-99-9 THF

STAGE(2)

RCT BE 491-30-5
SOL 109-99-9 THF


STAGE(3)

RGT F 76-05-1 F3CCO2H
SOL 75-09-2 CH2Cl2

PRO BF 357293-10-8

NTE stereoselective, solid-supported reactant, Wang resin used

RX(26) OF 32 A + BG ==> BH



BH
YIELD 33%

RX(26) RCT A 357292-05-4

STAGE(1)

RGT D 603-35-0 PPh3, E 2446-83-5 N2(CO2CHMe2)2
SOL 75-09-2 CH2Cl2, 109-99-9 THF

STAGE(2)

RCT BG 3336-49-0
SOL 109-99-9 THF



STAGE(3)

RGT F 76-05-1 F3CCO2H
SOL 75-09-2 CH2Cl2

PRO BH 357293-11-9

NTE stereoselective, solid-supported reactant, Wang resin used

RX(27) OF 32 A + BI ==> BJ



BJ
YIELD 75%

RX(27) RCT A 357292-85-4

STAGE(1)

ROT D 603-35-0 PPh3, E 2446-83-5 N2(CO2CHMe2)2
SOL 75-09-2 CH2Cl2, 109-99-9 THF

STAGE(2)

RCT BI 611-36-9
SOL 109-99-9 THF



STAGE(3)

ROT F 76-05-1 F3CCO2H
SOL 75-09-2 CH2Cl2

PRO BJ 357293-12-0

NTE stereoselective, solid-supported reactant, Wang resin used

RX(28) OF 32 A + BK ==> BL



BL
YIELD 57%

RX(28) RCT A 357292-85-4

STAGE(1)

ROT D 603-35-0 PPh3, E 2446-83-5 N2(CO2CHMe2)2
SOL 75-09-2 CH2Cl2, 109-99-9 THF

STAGE(2)

RCT BK 322-97-4
SOL 109-99-9 THF



STAGE(3)

ROT F 76-05-1 F3CCO2H
SOL 75-09-2 CH2Cl2

PRO BL 357293-13-1

NTE stereoselective, solid-supported reactant, Wang resin used

RX(29) OF 32 A + BM ==> BN



BN
YIELD 61%

RX(29) RCT A 357292-85-4

STAGE(1)

ROT D 603-35-0 PPh3, E 2446-83-5 N2(CO2CHMe2)2
SOL 75-09-2 CH2Cl2, 109-99-9 THF

STAGE(2)

RCT BM 64415-07-2
SOL 109-99-9 THF



STAGE(3)

ROT F 76-05-1 F3CCO2H
SOL 75-09-2 CH2Cl2

PRO BN 357293-14-2

NTE stereoselective, solid-supported reactant, Wang resin used

RX(30) OF 32 A + BC ==> BO



BO
YIELD 39%

RX(30) RCT A 357292-85-4

STAGE(1)

ROT D 603-35-0 PPh3, E 2446-83-5 N2(CO2CHMe2)2
SOL 75-09-2 CH2Cl2, 109-99-9 THF

STAGE(2)

RCT BC 130-16-5
SOL 109-99-9 THF


STAGE(3)

ROT F 76-05-1 F3CCO2H
SOL 75-09-2 CH2Cl2

PRO BO 357293-15-3

NTE stereoselective, solid-supported reactant, Wang resin used

RX(31) OF 32 A + BP ==> BQ



RX(31) RCT A 357292-85-4

STAGE(1)

RGT D 603-35-0 PPh₃, R 2446-83-5 N₂(CO₂CHMe₂)₂
 SOL 75-09-2 CH₂Cl₂, 109-99-9 THF

STAGE(2)

RCT BP 82121-05-9
 SOL 109-99-9 THF


STAGE(3)

RGT F 76-05-1 F₃CCO₂H
 SOL 75-09-2 CH₂Cl₂

PRO BQ 357293-16-4

NTE stereoselective, solid-supported reactant, Wang resin used

RX(32) OF 32 A + BR ==> BS



RX(32) RCT A 357292-85-4

STAGE(1)

RGT D 603-35-0 PPh₃, R 2446-83-5 N₂(CO₂CHMe₂)₂
 SOL 75-09-2 CH₂Cl₂, 109-99-9 THF

STAGE(2)

RCT BR 23432-39-5
 SOL 109-99-9 THF

STAGE(3)

RGT F 76-05-1 F₃CCO₂H
 SOL 75-09-2 CH₂Cl₂


PRO BS 357293-17-5

NTE stereoselective, solid-supported reactant, Wang resin used

L12 ANSWER 8 OF 18 CASREACT COPYRIGHT 2007 ACS ON STM
 ACCESSION NUMBER: 133:335259 CASREACT Full-text
 TITLE: 1-Aminocyclopropaneboronic Acid: Synthesis and
 Incorporation into an Inhibitor of Hepatitis C Virus
 NS3 Protease
 AUTHOR(S): Priestley, E. Scott; Decicco, Carl P.
 CORPORATE SOURCE: Department of Chemical and Physical Sciences, DuPont
 Pharmaceuticals Company, Wilmington, DE, 19880, USA
 SOURCE: Organic Letters (2000), 2(20), 3095-3097
 CODEN: ORLE77; ISSN: 1523-7060
 PUBLISHER: American Chemical Society
 DOCUMENT TYPE: Journal

LANGUAGE: English
 AB The previously unreported α,α-disubstituted 1-aminoboronic esters have potential utility in peptidomimetic design, particularly against serine protease targets. A concise synthesis of 1-aminocyclopropaneboronic pinanediol ester is reported, and a peptidyl derivative has modest affinity (K_i = 1.6 μM) for hepatitis C NS3 protease. Analogs with iso-Pr and cyclohexyl in place of cyclopropyl were also prepared and tested.
 REFERENCE COUNT: 22 THERE ARE 22 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

RX(10) OF 27 ...U ==> Y



* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(10) RCT U 303191-77-7
 RGT P 7647-01-0 HCl
 PRO Y 303191-80-2
 SOL 123-91-1 Dioxane

RX(12) OF 27 AB ==> AC

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

PAGE 2-A


AB

(12)

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

RX(12) RCT AB 303191-79-9
 RGT P 7647-01-0 HCl
 PRO AC 303191-82-4
 SOL 123-91-1 Dioxane


L12 ANSWER 9 OF 18 CASREACT COPYRIGHT 2007 ACS ON STM
 ACCESSION NUMBER: 127:331716 CASREACT Full-text
 TITLE: Folding types of dipeptides containing the
 diastereoisomeric cyclopropane analogs of
 phenylalanine
 AUTHOR(S): Jimenez, Ana I.; Vandersee, Regis; Marraud, Michel;
 Aubry, Andre; Cativiela, Carlos
 CORPORATE SOURCE: Unite de Recherche Associees au CNRS, Laboratoire de
 Chimie Physique Macromoleculaire ENSIC-INPL, Nancy,
 54001, Fr.
 SOURCE: Tetrahedron Letters (1997), 38(43), 7559-7562
 CODEN: TETL; ISSN: 0040-4039
 PUBLISHER: Elsevier
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI



AB In order to consider the possible influence of the orientation of a side chain on the peptide backbone, the mol. structure of four model dipeptides Piv-Pro-c3Phe-NHMe (Piv = Me₃CCO; c3Phe = 2,3-methanophenylalanine residue I) were studied by IR and 1H NMR. All four derivs. are β-folded, but the folding type depends on the stereochem. of the cyclopropane moiety.

REFERENCE COUNT: 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


RX(9) OF 24 ...K ==> P



RX(9) RCT K 197776-19-1
 RGT Q 74-87-3 MeCl, R 76-05-1 F₃CCO₂H, D 109-02-4


N-Methylmorpholine
PRO P 197778-08-8
SOL 75-09-2 CH₂Cl₂

RX(10) OF 24 ...M ==> T




RX(10) RCT M 197778-20-4
RGT Q 74-87-3 MeCl, R 76-05-1 F3CCO₂H, D 109-02-4
N-Methylmorpholine
PRO T 197778-09-9
SOL 75-09-2 CH₂Cl₂

RX(11) OF 24 ...N ==> U



RX(11) RCT N 197778-21-5
RGT Q 74-87-3 MeCl, R 76-05-1 F3CCO₂H, D 109-02-4
N-Methylmorpholine
PRO U 197778-12-4
SOL 75-09-2 CH₂Cl₂



RX(12) OF 24 ...O ==> V



RX(12) RCT O 197778-22-6
RGT Q 74-87-3 MeCl, R 76-05-1 F3CCO₂H, D 109-02-4
N-Methylmorpholine
PRO V 197778-13-5
SOL 75-09-2 CH₂Cl₂

RX(17) OF 24 COMPOSED OF RX(5), RX(9)

RX(17) C + J ==> P





RX(5) RCT C 197778-16-8, J 74-89-5
PRO K 197778-19-1
SOL 67-56-1 MeOH

RX(9) RCT K 197778-19-1
RGT Q 74-87-3 MeCl, R 76-05-1 F3CCO₂H, D 109-02-4
N-Methylmorpholine
PRO P 197778-08-8
SOL 75-09-2 CH₂Cl₂

RX(18) OF 24 COMPOSED OF RX(6), RX(10)

RX(18) F + J ==> T





RX(6) RCT F 197778-17-9, J 74-89-5
PRO M 197778-20-4
SOL 67-56-1 MeOH

RX(10) RCT M 197778-20-4
RGT Q 74-87-3 MeCl, R 76-05-1 F3CCO₂H, D 109-02-4
N-Methylmorpholine
PRO T 197778-09-9
SOL 75-09-2 CH₂Cl₂

RX(19) OF 24 COMPOSED OF RX(7), RX(11)

RX(19) H + J ==> U





RX(7) RCT H 213996-57-7, J 74-89-5
PRO N 197778-21-5
SOL 67-56-1 MeOH

RX(11) RCT N 197778-21-5
RGT Q 74-87-3 MeCl, R 76-05-1 F3CCO₂H, D 109-02-4
N-Methylmorpholine
PRO U 197778-12-4
SOL 75-09-2 CH₂Cl₂

RX(20) OF 24 COMPOSED OF RX(8), RX(12)

RX(20) I + J ==> V



RX(8) RCT I 213996-58-8, J 74-89-5
PRO O 197778-22-6
SOL 67-56-1 MeOH

RX(12) RCT O 197778-22-6
 ROT Q 74-87-3 MeCl, R 74-05-1 P3CCO2H, D 109-02-4
 N-Methylmorpholine
 PRO V 197778-11-5
 SOL 75-09-2 CH2Cl2

L12 ANSWER 10 OF 18 CASREACT COPYRIGHT 2007 ACS on STN


ACCESSION NUMBER: 116:230602 CASREACT [Full-text](#)

TITLE: Synthesis and properties of D-glucosamine N-peptidyl derivatives as substrate analog inhibitors of papain and cathepsin B
 AUTHOR(S): Giordano, C.; Gallina, C.; Consalvi, V.; Scandurra, R.
 CORPORATE SOURCE: Cent. Stud. Chim. Farm., Univ. La Sapienza, Rome, 00185, Italy
 SOURCE: European Journal of Medicinal Chemistry (1991), 26(8), 753-62
 CODEN: EJMCAS; ISSN: 0223-5234


DOCUMENT TYPE: Journal
 LANGUAGE: English

AB N-Peptidyl deriva. of D-glucosamine were synthesized and tested as reversible, substrate analog inhibitors of cysteine and serine proteases. D-Glucosamine itself showed fair inhibiting properties against cysteine proteases. Derive. designed to improved binding at the papain active site, displayed reversible inhibition with Ki 67-860 μM for papain and 111-2400 μM for cathepsin B. Representative serine proteases were unaffected. No inhibitory activity against human leukocyte elastase was observed for 2 deriva. bearing very effective peptidyl recognizing units for this enzyme.

RX(15) OF 26 ...O ==> X



O (15)



X
YIELD 70%

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS) SINCE FILE ENTRY TOTAL
 CA SUBSCRIBER PRICE -9.49 -10.22

IN FILE 'CASREACT' AT 08:51:09 ON 30 MAY 2007

=> d hist

(FILE 'HOME' ENTERED AT 08:18:24 ON 30 MAY 2007)

L1 FILE 'REGISTRY' ENTERED AT 08:18:37 ON 30 MAY 2007
 STRUCTURE UPLOADED

L2 FILE 'CASREACT' ENTERED AT 08:19:13 ON 30 MAY 2007
 0 S L1 SSS SAM
 L3 1 S L1 SSS FULL

L4 FILE 'REGISTRY' ENTERED AT 08:21:37 ON 30 MAY 2007
 STRUCTURE UPLOADED

L5 FILE 'CASREACT' ENTERED AT 08:21:56 ON 30 MAY 2007
 50 S L4

L6 FILE 'CASREACT' ENTERED AT 08:24:06 ON 30 MAY 2007
 28 S L5 NOT PY>2003

L7 FILE 'REGISTRY' ENTERED AT 08:28:44 ON 30 MAY 2007
 STRUCTURE UPLOADED

L8 FILE 'CASREACT' ENTERED AT 08:29:13 ON 30 MAY 2007
 5 S L7
 L9 3 S L6 NOT PY > 2003
 L10 46 S L7 SSS FULL
 L11 21 S L10 NOT PY > 2003
 L12 18 S L11 NOT L9

=> d ibib abs hit 11-18

L12 ANSWER 11 OF 18 CASREACT COPYRIGHT 2007 ACS on STN

ACCESSION NUMBER: 111:115724 CASREACT [Full-text](#)


TITLE: The incorporation of sugar moieties to neuropeptides: comparative study of different methods
 AUTHOR(S): Torres, J. L.; Haro, I.; Bardeji, E.; Valencia, G.; Garcia-Anton, J. M.; Reig, F.
 CORPORATE SOURCE: Dep. Biol. Org. Chem., CSIC, Barcelona, 08034, Spain
 SOURCE: Tetrahedron (1988), 44(19), 6131-6
 CODEN: TETRAH; ISSN: 0040-4020

DOCUMENT TYPE: Journal
 LANGUAGE: English


AB By using both β-N-glycosylation and β-O-glycosylation procedures, different methods for the incorporation of glucose moieties to proline-, hydroxyproline- or glutamic acid-containing protected neuropeptides have been examined. As far as glycosylation of Glu and Hyp containing fragments is concerned, the incorporation of either γ-β-N-glycosylated glutamic acid or 4-β-O-glycosylated hydroxyproline to the rest of the peptide have been chosen. However, in the case of C-terminal proline containing peptide fragments, direct β-N-glycosylation of the full peptide has been preferred. Acetyl protecting groups on the sugar moiety led to better yields than the bulkier benzyl groups.

RX(15) RCT O 141280-20-8
 PRO X 141266-00-4

RX(16) OF 26 ...Q ==> Y



Q (16)




Y
YIELD 71%


RX(16) RCT Q 141280-21-9
 PRO Y 141266-01-5

=> d cost		SINCE FILE		TOTAL	
COST IN U.S. DOLLARS		ENTRY	SESSION		
CONNECT CHARGES		14.43	21.99		
NETWORK CHARGES		2.22	3.42		
SEARCH CHARGES		116.94	231.96		
DISPLAY CHARGES		82.06	88.79		
FULL ESTIMATED COST		215.65	346.16		


RX(7) OF 41 ...M + S ==> R



M



S (7)



S
YIELD 5%

RX(7) RCT M 122350-56-7

STAGE(1)


RGT T 1333-74-0 H2
 CAT 7440-05-3 Pd
 SOL 67-56-1 MeOH

STAGE(2)

RCT S 115615-64-0
 RGT S 538-75-0 DCC, F 2592-95-2 1-Benzotriazolol


PRO R 115730-55-7

L12 ANSWER 12 OF 18 CASREACT COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 109:6942 CASREACT Full-text
 TITLE: Amino acid derivatives that stabilize secondary structures of polypeptides. III. β -Enamino nitriles as analogs of secondary amides. The MCC group [1-(acylamino)-2-(aminoalkyl)-3-cyano-2-cyclopentenes] as amino acid analogs
 AUTHOR(S): Kemp, D. S.; Carter, Jeffery S.
 CORPORATE SOURCE: Dep. Chem., Massachusetts Inst. Technol., Cambridge, MA, 02139, USA
 SOURCE: Tetrahedron Letters (1987), 28(40), 4641-4
 CODEN: TELEAY; ISSN: 0040-4039
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 GI



AB The properties of β -cyanoenamines as analogs of amides are studied with deriva. of a rigid amino acid equivalent I (Boc = Me₃CO₂C), i.e., Boc-DL-Mcc-CH₂CO₂CH₂Ph, prepared in two steps from Me 2-(N-Boc-amino)-5- cyano-pentanoate. Racemation of Mcc and incorporation into cyclic pentapeptide cyclo(Pro-Gly-Pro-DL-Mcc-Gly) are described.


RX(4) OF 21 ...J + L ==> N...




(4) →

RX(5) RCT M 114542-74-4
 ROT O 1333-74-0 H2
 PRO N 114518-88-6
 CAT 7440-05-3 Pd
 SOL 64-19-7 AcOH


RX(10) OF 21 COMPOSED OF RX(4), RX(5)
 RX(10) J + L ==> N



2 STEPS →


N
YIELD 93%

RX(4) RCT J 114542-73-3
 STAGE(1)
 ROT K 76-05-1 F3CCO₂H, H 75-09-2 CH₂Cl₂
 STAGE(2)
 RCT L 36254-59-8
 PRO M 114542-74-4
 RX(5) RCT M 114542-74-4
 ROT O 1333-74-0 H2
 PRO N 114518-88-6
 CAT 7440-05-3 Pd


M
YIELD 61%

RX(4) RCT J 114542-73-3
 STAGE(1)
 ROT K 76-05-1 F3CCO₂H, H 75-09-2 CH₂Cl₂
 STAGE(2)
 RCT L 36254-59-8
 PRO M 114542-74-4

RX(5) OF 21 ...M ==> N...



(5) →

N
YIELD 93%


SOL 64-19-7 AcOH

L12 ANSWER 13 OF 18 CASREACT COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 108:112956 CASREACT Full-text
 TITLE: Preparation of amino acid and peptide arylamides as chromogenic reagents in enzyme determinations
 INVENTOR(S): Bajusz, Sándor; Juhasz, Attila; Barabas, Eva; Bagdi, Daniel; Mohai, László, Mrs.
 PATENT ASSIGNER(S): Gyógyszerkutató Intézet, Hung.
 SOURCE: Hung. Teljes, 37 pp.
 CODEN: HUXKBU
 DOCUMENT TYPE: Patent
 LANGUAGE: Hungarian
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
HU 40615	A2	19870128	HU 1985-2330	19850613
HU 198172	B	19890828		
SU 1512484	A3	19890930	SU 1986-4028098	19860908
			HU 1985-2330	19850613


PRIORITY APPLN. INFO.:
 AB The title compds. RXA (R = H, acyl, aminoacyl, acylaminoacyl, peptidyl, acylpeptidyl; X = L- α -amino acid radical; A = p-nitroanilino, 4-methylcoumarinyl-7-amino) are prepared by reacting the protected corresponding amino acid or peptide with PCl₃ and the aryl amide, in <0.2% water-containing pyridine, followed by deprotection. The arylamine/PCl₃ mol. ratio is 1:0.6-1.5, which is more PCl₃ than the conventional amount. A solution of 6.8 g N2-tert-butyloxycarbonyl-L-arginine-HCl.H₂O and 2.8 g p-nitroaniline in 20 mL pyridine, was treated, at -20°, with 2.26 mL PCl₃, to give N2-tert-butyloxycarbonyl-L-arginine-p-nitroanilide-HCl.H₂O, which was treated with HCl in EtOH to give L-arginine-p-nitroanilide-2HCl.H₂O. The products are chromogenic and fluorogenic reagents in the determination of proteases and transpeptidases.

RX(16) OF 67 ...Y + X ==> Z



X


(16) →



Z
YIELD 90%


RX(16) RCT Y 110-15-6, X 113277-38-6
PRO Z 113277-36-4

L12 ANSWER 14 OF 18 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 10618224 CASREACT Full-text
TITLE: Total synthesis of neothramycin
AUTHOR(S): Mori, Miwako; Uozumi, Yasuhiro; Ban, Yoshio
CORPORATE SOURCE: Fac. Pharm. Sci., Hokkaido Univ., Sapporo, 060, Japan
SOURCE: Journal of the Chemical Society, Chemical Communications (1986), (11), 841-2
CODEN: JCCCAT; ISSN: 0022-4936
DOCUMENT TYPE: Journal
LANGUAGE: English
GI



AB Neothramycin A and B (I; R = H, R1 = OH; R = OH, R1 = H, resp.), from Streptomyces, were prepared in 12 steps from 5,4,2-(4-MeC6H4SO3)(MeO)BrC6H2NH2 and the hydroxyproline II, via the key intermediate III, which was obtained by Pd(PPh3)4-catalyzed carbonylation of the secondary amine IV followed by deprotection.

RX(2) OF 112 ...C ==> D...






RX(2) RCT C 105842-35-1
RGT E 144-55-8 NaHCO3
PRO D 105823-07-2
SOL 7732-18-5 Water

RX(3) RCT D 105823-07-2, G 107-30-2
RGT I 7087-68-5 EtN(Pr-1)2
PRO H 105823-08-3


L12 ANSWER 15 OF 18 CASREACT COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 10560932 CASREACT Full-text
TITLE: Peptidyl carbamates incorporating amino acid isosteres as novel elastase inhibitors
AUTHOR(S): Digenis, George A.; Agha, Bushra J.; Tsuji, Kiyoshi; Kato, Masayuki; Shinogi, Masaki
CORPORATE SOURCE: Coll. Pharm., Univ. Kentucky, Lexington, KY, 40536-0053, USA
SOURCE: Journal of Medicinal Chemistry (1986), 29(8), 1468-76
CODEN: JMCHAM; ISSN: 0022-2623
DOCUMENT TYPE: Journal
LANGUAGE: English
AB Title peptidyl carbamates MeO2CCH2CH2CO-Ala-Ala-Pro-NH2O2CNRR1 [I; Z = p-C6H4, R = H, R1 = Ph, CHMe2; Z = p-C6H4, R = R1 = Me; Z = o-C6H4, CH(CHMe2)CH2, R = H, R1 = Ph] and MeO2CCH2CH2CO-Ala-Ala-Pro-CH2N(CHMe2)COXR2 [II; X = O, R2 = C6H4NO2-p, Ph, C6F5, CH2CF2CF2CF3; X = S, R2 = CH2Ph, 1-methyl-5-tetrazolyl, 1-phenyl-5-tetrazolyl] were prepared and they were tested as inhibitors of elastase, trypsin, and chymotrypsin. Thus, Boc-Pro-NH2OH (Boc = Me3CO2C) were treated with RR1NCOCl or RR1NCO to give Boc-Pro-NH2O2CNRR1, which were Boc-deblocked and then coupled with MeO2CCH2CH2CO-Ala-Ala-OH (III) by ClCO2CH2CHMe2 to give I. Boc-Pro-CH2Cl was treated with H2NCHMe2 to give Boc-Pro-CH2NCHMe2, which was treated with R2XCOCl to give Boc-Pro-CH2N(CHMe2)COXR2, which were Boc-deblocked and then coupled with III to give II. Six peptidyl carbamates specifically inhibited elastase without affecting trypsin and chymotrypsin. Kinetic studies showed that the inhibition was competitive. The inhibition is reversible and proceeds via the rapid formation of a strong enzyme-inhibitor complex, followed by slow acylation of the serine residue on the active site of the enzyme.

RX(16) OF 173 ...G + AE ==> AL




RX(2) RCT C 105842-35-1
RGT E 144-55-8 NaHCO3
PRO D 105823-07-2
SOL 7732-18-5 Water


RX(18) OF 112 COMPOSED OF RX(2), RX(3)
RX(18) C + 2 G ==> H



2
STEPS







AE ● HCl (16)



RX(16) RCT G 102284-27-5, AE 102284-36-6
RGT AM 543-27-1 ClCO2Bu-1, AN 109-02-4 N-Methylmorpholine
PRO AL 92279-27-1
SOL 109-99-9 THF

RX(17) OF 173 ...G + AH ==> AO









AO

RX(17) RCT G 102284-27-5, AH 102284-37-7
 RGT AM 543-27-1 ClCO2Bu-1, AN 109-02-4 N-Methylmorpholine, AP
 10416-59-8 Me3SiN:CMOSiMe3
 PRO AO 92279-28-2
 SOL 109-99-9 THF

RX(18) OF 173 ...O + AI ==> AQ




O



AQ

RX(18) RCT G 102284-27-5, AI 102284-38-8
 RGT AM 543-27-1 ClCO2Bu-1
 PRO AQ 92279-29-3


RX(19) OF 173 ...O + AJ ==> AR



O

AJ

(19)




AR

RX(19) RCT G 102284-27-5, AJ 102284-39-9
 RGT AM 543-27-1 ClCO2Bu-1
 PRO AR 92279-30-6


RX(42) OF 173 COMPOSED OF RX(2), RX(16)

RX(42) C + F + AE ==> AL




C

F



AE

\bullet HCl
 3 STEPS




AL

RX(2) RCT C 52787-46-9, F 1948-31-8
 RGT H 144-55-8 NaHCO3
 PRO G 102284-27-5
 SOL 7732-18-5 Water, 67-64-1 Me2CO

RX(16) RCT G 102284-27-5, AR 102284-36-6
 RGT AM 543-27-1 ClCO2Bu-1, AN 109-02-4 N-Methylmorpholine
 PRO AL 92279-27-1
 SOL 109-99-9 THF


RX(43) OF 173 COMPOSED OF RX(2), RX(17)

RX(43) C + F + AH ==> AO




C

F



AH

\bullet HCl
 2 STEPS




AO

RX(2) RCT C 52787-46-9, F 1948-31-8
 RGT H 144-55-8 NaHCO₃
 PRO G 102284-27-5
 SOL 7732-18-5 Water, 67-64-1 Me₂CO


RX(17) RCT G 102284-27-5, AH 102284-37-7
 RGT AM 543-27-1 ClCO₂Bu-i, AN 109-02-4 N-Methylmorpholine, AP
 10416-59-8 Me₃SiN:CMOSiMe₃
 PRO AO 92279-29-2
 SOL 109-99-9 THF

RX(44) OF 173 COMPOSED OF RX(2), RX(18)
 RX(44) C + F + AI ==> AQ




C

F



AI

● HCl


2
STEPS

AQ

RX(2) RCT C 52787-46-9, F 1948-31-8
 RGT H 144-55-8 NaHCO₃
 PRO G 102284-27-5
 SOL 7732-18-5 Water, 67-64-1 Me₂CO


RX(18) RCT G 102284-27-5, AI 102284-38-8
 RGT AM 543-27-1 ClCO₂Bu-i
 PRO AQ 92279-29-3

RX(45) OF 173 COMPOSED OF RX(2), RX(19)
 RX(45) C + F + AJ ==> AR




C

F



AJ

● HCl


2
STEPS

AR


RX(2) RCT C 52787-46-9, F 1948-31-8
 RGT H 144-55-8 NaHCO₃
 PRO G 102284-27-5
 SOL 7732-18-5 Water, 67-64-1 Me₂CO

RX(19) RCT G 102284-27-5, AJ 102284-39-9
 RGT AM 543-27-1 ClCO₂Bu-i
 PRO AR 92279-30-6


RX(64) OF 173 COMPOSED OF RX(11), RX(16)
 RX(64) V + G ==> AL



V



G


2
STEPS

AL


RX(11) RCT V 102284-31-1
 RGT AP 7647-01-0 HCl
 PRO AR 102284-36-6
 SOL 109-99-9 THF, 64-18-6 HCO₂H

RX(16) RCT G 102284-27-5, AR 102284-36-6
 RGT AM 543-27-1 ClCO₂Bu-i, AN 109-02-4 N-Methylmorpholine
 PRO AL 92279-27-1
 SOL 109-99-9 THF

RX(65) OF 173 COMPOSED OF RX(12), RX(17)
 RX(65) Z + G ==> AO




Z



G

2
STEPS




AQ

RX(12) RCT Z 102284-32-2
RGT AF 7647-01-0 HCl
PRO AH 102284-37-7


RX(17) RCT G 102284-27-5, AH 102284-37-7
RGT AM 543-27-1 ClCO2Bu-1, AN 109-02-4 N-Methylmorpholine, AP 10416-59-8 Me3SiN:CMOSiMe3
PRO AO 92279-28-2
SOL 109-99-9 THF

RX(66) OF 173 COMPOSED OF RX(13), RX(18)

RX(66) AB + G ==> AQ




AB



G

2
STEPS




AQ

RX(13) RCT AB 102284-33-3
RGT AF 7647-01-0 HCl
PRO AI 102284-38-8


RX(18) RCT G 102284-27-5, AI 102284-38-8
RGT AM 543-27-1 ClCO2Bu-1
PRO AQ 92279-29-3

RX(67) OF 173 COMPOSED OF RX(14), RX(19)

RX(67) AC + G ==> AR




AC



G

2
STEPS




AR

RX(14) RCT AC 102284-34-4
RGT AF 7647-01-0 HCl
PRO AJ 102284-39-9

RX(19) RCT G 102284-27-5, AJ 102284-39-9
RGT AM 543-27-1 ClCO2Bu-1
PRO AR 92279-30-6

RX(81) OF 173 COMPOSED OF RX(1), RX(2), RX(16)


RX(81) A + B + F + AE ==> AL



A

B


F



AE

● HCl

3
STEPS



AL


RX(1) RCT A 1490-25-1, B 6066-82-6
RGT D 121-44-8 Et3N
PRO C 52787-46-9
SOL 141-78-6 AcOEt

RX(2) RCT C 52787-46-9, F 1948-31-8
RGT H 144-55-8 NaHCO3
PRO G 102284-27-5
SOL 7732-18-5 Water, 67-64-1 Me2CO

RX(16) RCT G 102284-27-5, AR 102284-36-6
RGT AM 543-27-1 ClCO2Bu-1, AN 109-02-4 N-Methylmorpholine
PRO AL 92279-27-1
SOL 109-99-9 THF

RX(82) OF 173 COMPOSED OF RX(1), RX(2), RX(17)


RX(82) A + B + F + AH ==> AO



A

B


F



AH

● HCl

3
STEPS




AO

RX(1) RCT A 1490-25-1, B 6066-82-6
 RGT D 121-44-8 Et3N
 PRO C 52787-46-9
 SOL 141-78-6 AcOEt


RX(2) RCT C 52787-46-9, F 1948-31-8
 RGT H 144-55-8 NaHCO3
 PRO G 102284-27-5
 SOL 7732-18-5 Water, 67-64-1 Me2CO

RX(17) RCT G 102284-27-5, AH 102284-37-7
 RGT AM 543-27-1 ClCO2Bu-i, AN 109-02-4 N-Methylmorpholine, AP 10416-59-8 Me3SiN:CMOSiMe3
 PRO AO 92279-28-2
 SOL 109-99-9 THF


RX(83) OF 173 COMPOSED OF RX(1), RX(2), RX(18)
 RX(83) A + B + F + AI ==> AQ




A



B




F



AI

● HCl

3 STEPS




AQ

RX(1) RCT A 1490-25-1, B 6066-82-6
 RGT D 121-44-8 Et3N
 PRO C 52787-46-9
 SOL 141-78-6 AcOEt


RX(2) RCT C 52787-46-9, F 1948-31-8
 RGT H 144-55-8 NaHCO3
 PRO G 102284-27-5
 SOL 7732-18-5 Water, 67-64-1 Me2CO

RX(18) RCT G 102284-27-5, AI 102284-38-8
 RGT AM 543-27-1 ClCO2Bu-i
 PRO AQ 92279-29-3


RX(84) OF 173 COMPOSED OF RX(1), RX(2), RX(19)
 RX(84) A + B + F + AI ==> AR




A



B




F



AJ

● HCl

3 STEPS



AR

RX(1) RCT A 1490-25-1, B 6066-82-6
 RGT D 121-44-8 Et3N
 PRO C 52787-46-9
 SOL 141-78-6 AcOEt


RX(2) RCT C 52787-46-9, F 1948-31-8
 RGT H 144-55-8 NaHCO3
 PRO G 102284-27-5
 SOL 7732-18-5 Water, 67-64-1 Me2CO

RX(19) RCT G 102284-27-5, AJ 102284-39-9
 RGT AM 543-27-1 ClCO2Bu-i
 PRO AR 92279-30-6

RX(93) OF 173 COMPOSED OF REACTION SEQUENCE RX(1), RX(16)
 AND REACTION SEQUENCE RX(2), RX(16)


...V ==> AE...

... C + F + AE ==> AL



V


2 STEPS




AE

● HCl


START NEXT REACTION SEQUENCE



C




F



AE

● HCl

2 STEPS



AL


RX(11) RCT V 102284-31-1
 RGT AF 7647-01-0 HCl
 PRO AS 102284-36-6
 SOL 109-99-9 THF, 64-18-6 HCO₂H

RX(2) RCT C 52787-46-9, F 1948-31-8
 RGT H 144-55-8 NaHCO₃
 PRO G 102284-27-5
 SOL 7732-18-5 Water, 67-64-1 Me₂CO


RX(16) RCT G 102284-27-5, AH 102284-36-6
 RGT AM 543-27-1 ClCO₂Bu-1, AN 109-02-4 N-Methylmorpholine
 PRO AL 92279-27-1
 SOL 109-99-9 THF

RX(94) OF 173 COMPOSED OF REACTION SEQUENCE RX(12), RX(17)
 AND REACTION SEQUENCE RX(2), RX(17)

...Z ==> AH...
 ... C + F + AH ==> AO




2
 STEPS



● HCl


AH

START NEXT REACTION SEQUENCE



C


F



● HCl

AH

2
 STEPS



AO


RX(12) RCT Z 102284-32-2
 RGT AF 7647-01-0 HCl
 PRO AH 102284-37-7

RX(2) RCT C 52787-46-9, F 1948-31-8
 RGT H 144-55-8 NaHCO₃
 PRO G 102284-27-5
 SOL 7732-18-5 Water, 67-64-1 Me₂CO


RX(17) RCT G 102284-27-5, AH 102284-37-7
 RGT AM 543-27-1 ClCO₂Bu-1, AN 109-02-4 N-Methylmorpholine, AP
 10416-59-8 Me₃SiN:CMesOSiMe₃
 PRO AO 92279-28-2
 SOL 109-99-9 THF

RX(95) OF 173 COMPOSED OF REACTION SEQUENCE RX(13), RX(18)
 AND REACTION SEQUENCE RX(2), RX(18)

...AB ==> AI...
 ... C + F + AI ==> AQ




2
 STEPS



● HCl


AI

START NEXT REACTION SEQUENCE



C


F



● HCl

AI

2
 STEPS



AQ


RX(13) RCT AB 102284-33-3
 RGT AF 7647-01-0 HCl
 PRO AI 102284-38-8

RX(2) RCT C 52787-46-9, F 1948-31-8
 RGT H 144-55-8 NaHCO₃
 PRO G 102284-27-5
 SOL 7732-18-5 Water, 67-64-1 Me₂CO




RX(18) RCT G 102284-27-5, AI 102284-38-8
 RGT AM 543-27-1 ClCO₂Bu-1
 PRO AQ 92279-29-3

RX(96) OF 173 COMPOSED OF REACTION SEQUENCE RX(14), RX(19)
 AND REACTION SEQUENCE RX(2), RX(19)

...AC ==> AJ...
... C + F + AJ ==> AR



START NEXT REACTION SEQUENCE





RX(14) RCT AC 102284-34-4
RGT AF 7647-01-0 HCl
PRO AJ 102284-39-9

RX(2) RCT C 52787-46-9, F 1948-31-8
RGT H 144-55-8 NaHCO3
PRO G 102284-27-5
SOL 7732-18-5 Water, 67-64-1 Me2CO




RX(19) RCT G 102284-27-5, AJ 102284-39-9
RGT AM 543-27-1 ClCO2Bu-1
PRO AR 92279-30-6

RX(102) OF 173 COMPOSED OF REACTION SEQUENCE RX(11), RX(16)
AND REACTION SEQUENCE RX(1), RX(2), RX(16)

...V ==> AE...
... A + B + F + AE ==> AL



START NEXT REACTION SEQUENCE



RX(11) RCT V 102284-31-1
RGT AP 7647-01-0 HCl
PRO AR 102284-36-6
SOL 109-99-9 THF, 64-18-6 HCO2H

RX(1) RCT A 1490-25-1, B 6066-82-6
RGT D 121-44-8 Et3N
PRO C 52787-46-9
SOL 141-78-6 AcOEt



RX(2) RCT C 52787-46-9, F 1948-31-8
RGT H 144-55-8 NaHCO3
PRO G 102284-27-5
SOL 7732-18-5 Water, 67-64-1 Me2CO

RX(16) RCT G 102284-27-5, AE 102284-36-6
RGT AM 543-27-1 ClCO2Bu-1, AN 109-02-4 N-Methylmorpholine


PRO AL 92279-27-1
SOL 109-99-9 THF



RX(103) OF 173 COMPOSED OF REACTION SEQUENCE RX(12), RX(17)
AND REACTION SEQUENCE RX(1), RX(2), RX(17)

...Z ==> AH...
... A + B + F + AH ==> AO



START NEXT REACTION SEQUENCE






RX(12) RCT Z 102284-32-2
 RGT AP 7647-01-0 HCl
 PRO AH 102284-37-7

RX(1) RCT A 1490-25-1, B 6066-82-6
 RGT D 121-44-8 Et3N
 PRO C 52787-46-9
 SOL 141-78-6 AcOEt

RX(2) RCT C 52787-46-9, F 1948-31-8
 RGT H 144-55-8 NaHCO3
 PRO G 102284-27-5
 SOL 7732-18-5 Water, 67-64-1 Me2CO

RX(17) RCT G 102284-27-5, AH 102284-37-7
 RGT AM 543-27-1 ClCO2Bu-1, AN 109-02-4 N-Methylmorpholine, AP 10416-59-8 Me3SiN:CMOSiMe3
 PRO AO 92279-28-2
 SOL 109-99-9 THF

RX(104) OF 173 COMPOSED OF REACTION SEQUENCE RX(13), RX(18)
 AND REACTION SEQUENCE RX(1), RX(2), RX(18)
 ...AB ----> AI...
 ... A + B + F + AI ----> AQ




RX(13) RCT AB 102284-33-3
 RGT AP 7647-01-0 HCl
 PRO AI 102284-38-8

RX(1) RCT A 1490-25-1, B 6066-82-6
 RGT D 121-44-8 Et3N
 PRO C 52787-46-9
 SOL 141-78-6 AcOEt


RX(2) RCT C 52787-46-9, F 1948-31-8
 RGT H 144-55-8 NaHCO3
 PRO G 102284-27-5
 SOL 7732-18-5 Water, 67-64-1 Me2CO

RX(18) RCT G 102284-27-5, AI 102284-38-8
 RGT AM 543-27-1 ClCO2Bu-1
 PRO AQ 92279-29-3




RX(105) OF 173 COMPOSED OF REACTION SEQUENCE RX(14), RX(19)
 AND REACTION SEQUENCE RX(1), RX(2), RX(19)
 ...AC ----> AJ...
 ... A + B + F + AJ ----> AR



START NEXT REACTION SEQUENCE



START NEXT REACTION SEQUENCE



RX(14) RCT AC 102284-34-4
 RGT AP 7647-01-0 HCl
 PRO AJ 102284-39-9




RX(1) RCT A 1490-25-1, B 6066-82-6
 RGT D 121-44-8 Et3N
 PRO C 52787-46-9
 SOL 141-78-6 AcOEt

RX(2) RCT C 52787-46-9, F 1948-31-8
 RGT H 144-55-8 NaHCO3
 PRO G 102284-27-5
 SOL 7732-18-5 Water, 67-64-1 Me2CO

RX(19) RCT G 102284-27-5, AJ 102284-39-9
 RGT AM 543-27-1 ClCO2Bu-1

PRO AR 92279-30-6

RX(116) OF 173 COMPOSED OF RX(6), RX(11), RX(16)
 RX(116) M + U + G ==> AL






RX(6) RCT M 102284-28-6, U 103-71-9
 RGT W 110-86-1 Pyridine
 PRO V 102284-31-1
 SOL 68-12-2 DMF

RX(11) RCT V 102284-31-1
 RGT AF 7647-01-0 HCl
 PRO AE 102284-36-6
 SOL 109-99-9 THF, 64-18-6 HCO2H

RX(16) RCT G 102284-27-5, AE 102284-36-6
 RGT AM 543-27-1 ClCO2Bu-i, AN 109-02-4 N-Methylmorpholine, AP
 PRO AL 92279-27-1
 SOL 109-99-9 THF

RX(118) OF 173 COMPOSED OF RX(7), RX(12), RX(17)
 RX(118) M + Y + G ==> AO






RX(7) RCT M 102284-28-6, Y 79-44-7
 PRO Z 102284-32-2
 SOL 110-86-1 Pyridine

RX(12) RCT Z 102284-32-2
 RGT AF 7647-01-0 HCl
 PRO AH 102284-37-7

RX(17) RCT G 102284-27-5, AH 102284-37-7
 RGT AM 543-27-1 ClCO2Bu-i, AN 109-02-4 N-Methylmorpholine, AP
 10416-59-8 Me3SiN:CMOSiMe3
 PRO AO 92279-28-2
 SOL 109-99-9 THF

RX(120) OF 173 COMPOSED OF RX(8), RX(13), RX(18)
 RX(120) M + AA + G ==> AQ






RX(8) RCT M 102284-28-6, AA 1795-48-8
 RGT W 110-86-1 Pyridine
 PRO AB 102284-33-3
 SOL 68-12-2 DMF

RX(13) RCT AB 102284-33-3
 RGT AF 7647-01-0 HCl
 PRO AI 102284-38-8

RX(18) RCT G 102284-27-5, AI 102284-38-8
 RGT AM 543-27-1 ClCO2Bu-i
 PRO AQ 92279-29-3

RX(122) OF 173 COMPOSED OF RX(9), RX(14), RX(19)
 RX(122) Q + U + G ==> AR





RX(9) RCT Q 102284-29-7, U 103-71-9
 PRO AC 102284-34-4







RX(14) RCT AC 102284-34-4
 RGT AF 7647-01-0 HCl
 PRO AJ 102284-39-9

RX(19) RCT G 102284-27-5, AJ 102284-39-9
 RGT AM 543-27-1 ClCO2Bu-i
 PRO AR 92279-30-6

RX(126) OF 173 COMPOSED OF REACTION SEQUENCE RX(2), RX(16)
 AND REACTION SEQUENCE RX(6), RX(11), RX(16)
 ...C + F ==> G...
 ...M + U + G ==> AL



START NEXT REACTION SEQUENCE



RX(2) RCT C 52787-46-9, F 1948-31-8
RGT H 144-55-8 NaHCO₃
PRO G 102284-27-5
SOL 7732-18-5 Water, 67-64-1 Me₂CO

RX(7) RCT M 102284-28-6, Y 79-44-7
PRO Z 102284-32-2
SOL 110-86-1 Pyridine

RX(12) RCT Z 102284-32-2
RGT AF 7647-01-0 HCl
PRO AH 102284-37-7

RX(17) RCT G 102284-27-5, AH 102284-37-7
RGT AM 543-27-1 ClCO₂Bu-1, AN 109-02-4 N-Methylmorpholine, AP 10416-59-8 Me₃SiN:CH₃OSiMe₃
PRO AO 92279-28-2
SOL 109-99-9 THF

RX(128) OF 173 COMPOSED OF REACTION SEQUENCE RX(2), RX(18)



RX(2) RCT C 52787-46-9, F 1948-31-8
RGT H 144-55-8 NaHCO₃
PRO G 102284-27-5
SOL 7732-18-5 Water, 67-64-1 Me₂CO

RX(6) RCT M 102284-28-6, U 103-71-9
RGT W 110-86-1 Pyridine
PRO V 102284-31-1
SOL 68-12-2 DMP

RX(11) RCT V 102284-31-1
RGT AF 7647-01-0 HCl
PRO AR 102284-36-6
SOL 109-99-9 THF, 64-18-6 HCO₂H



RX(16) RCT G 102284-27-5, AR 102284-36-6
RGT AM 543-27-1 ClCO₂Bu-1, AN 109-02-4 N-Methylmorpholine
PRO AL 92279-27-1
SOL 109-99-9 THF

RX(127) OF 173 COMPOSED OF REACTION SEQUENCE RX(2), RX(17)
AND REACTION SEQUENCE RX(7), RX(12), RX(17)
...C + F ==> G...
... M + Y + G ==> AO






START NEXT REACTION SEQUENCE

AND REACTION SEQUENCE RX(8), RX(13), RX(18)
...C + F ==> G...
... M + AA + G ==> AQ



START NEXT REACTION SEQUENCE







AQ


- RX(2) RCT C 52787-46-9, F 1948-31-8
 RGT H 144-55-8 NaHCO₃
 PRO G 102284-27-5
 SOL 7732-18-5 Water, 67-64-1 Me₂CO
- RX(8) RCT M 102284-26-6, AA 1795-48-8
 RGT W 110-86-1 Pyridine
 PRO AB 102284-33-3
 SOL 68-12-2 DMF
- RX(13) RCT AB 102284-33-3
 RGT AF 7647-01-0 HCl
 PRO AI 102284-38-8
- RX(18) RCT G 102284-27-5, AI 102284-38-8
 RGT AM 543-27-1 ClCO₂Bu-1
 PRO AQ 92279-29-3

RX(129) OF 173 COMPOSED OF REACTION SEQUENCE RX(2), RX(19)
 AND REACTION SEQUENCE RX(9), RX(14), RX(19)
 ...C + F ==> Q...
 ...Q + U + G ==> AR




G

...M + U ==> AR...
 ...A + B + F + AR ==> AL




M




● HCl
 AR

START NEXT REACTION SEQUENCE



A




F




F

START NEXT REACTION SEQUENCE




Q

U



G


3 STEPS



AR

- RX(2) RCT C 52787-46-9, F 1948-31-8
 RGT H 144-55-8 NaHCO₃
 PRO G 102284-27-5
 SOL 7732-18-5 Water, 67-64-1 Me₂CO
- RX(9) RCT Q 102284-29-7, U 103-71-9
 PRO AC 102284-34-4
- RX(14) RCT AC 102284-34-4
 RGT AF 7647-01-0 HCl
 PRO AJ 102284-39-9
- RX(19) RCT G 102284-27-5, AJ 102284-39-9
 RGT AM 543-27-1 ClCO₂Bu-1
 PRO AR 92279-30-6


RX(147) OF 173 COMPOSED OF REACTION SEQUENCE RX(6), RX(11), RX(16)
 AND REACTION SEQUENCE RX(1), RX(2), RX(16)



AL

- RX(6) RCT M 102284-28-6, U 103-71-9
 RGT M 110-86-1 Pyridine
 PRO V 102284-31-1
 SOL 68-12-2 DMF
- RX(11) RCT V 102284-31-1
 RGT AF 7647-01-0 HCl
 PRO AR 102284-36-6
 SOL 109-99-9 THF, 64-18-6 HCO₂H
- RX(1) RCT A 1490-25-1, B 6066-82-6
 RGT D 121-44-8 Et₃N
 PRO C 52787-46-9
 SOL 141-78-6 AcOEt
- RX(2) RCT C 52787-46-9, F 1948-31-8
 RGT H 144-55-8 NaHCO₃
 PRO G 102284-27-5
 SOL 7732-18-5 Water, 67-64-1 Me₂CO
- RX(16) RCT G 102284-27-5, AR 102284-36-6
 RGT AM 543-27-1 ClCO₂Bu-1, AN 109-02-4 N-Methylmorpholine
 PRO AL 92279-27-1
 SOL 109-99-9 THF


RX(148) OF 173 COMPOSED OF REACTION SEQUENCE RX(7), RX(12), RX(17)
 AND REACTION SEQUENCE RX(1), RX(2), RX(17)
 ...M + Y ==> AR...
 ...A + B + F + AR ==> AO



M

Y


3 STEPS




● HCl

AH


START NEXT REACTION SEQUENCE




A



B




F




● HCl

AH


3
STEPS

AO


RX (7) RCT M 102284-28-6, Y 79-44-7
PRO Z 102284-32-2




A



B




F



● HCl

AI

3
STEPS

AQ

RX (8) RCT M 102284-28-6, AA 1795-48-8
RGT W 110-86-1 Pyridine
PRO AB 102284-33-3
SOL 68-12-2 DMP

RX (13) RCT AB 102284-33-3
RGT AF 7647-01-0 HCl
PRO AI 102284-38-8

RX (1) RCT A 1490-25-1, B 6066-82-6
RGT D 121-44-8 Et3N
PRO C 52787-46-9
SOL 141-78-6 AcOEt

RX (2) RCT C 52787-46-9, F 1948-31-8
RGT H 144-55-8 NaHCO3
PRO G 102284-27-5
SOL 7732-18-5 Water, 67-64-1 Me2CO

SOL 110-86-1 Pyridine

RX (12) RCT Z 102284-32-2
RGT AF 7647-01-0 HCl
PRO AH 102284-37-7


RX (1) RCT A 1490-25-1, B 6066-82-6
RGT D 121-44-8 Et3N
PRO C 52787-46-9
SOL 141-78-6 AcOEt

RX (2) RCT C 52787-46-9, F 1948-31-8
RGT H 144-55-8 NaHCO3
PRO G 102284-27-5
SOL 7732-18-5 Water, 67-64-1 Me2CO

RX (17) RCT G 102284-27-5, AH 102284-37-7
RGT AM 543-27-1 ClCO2Bu-1, AN 109-02-4 N-Methylmorpholine, AP
10416-59-8 Me3SiN:CMOSiMe3
PRO AO 92279-28-2
SOL 109-99-9 THF


RX (149) OF 173 COMPOSED OF REACTION SEQUENCE RX (8), RX (13), RX (18)
AND REACTION SEQUENCE RX (1), RX (2), RX (18)

...M + AA ==> AI...
... A + B + F + AI ==> AQ



M

AA

3
STEPS

● HCl


AI

START NEXT REACTION SEQUENCE

RX (18) RCT G 102284-27-5, AI 102284-38-8
RGT AM 543-27-1 ClCO2Bu-1
PRO AQ 92279-29-3


RX (150) OF 173 COMPOSED OF REACTION SEQUENCE RX (9), RX (14), RX (19)
AND REACTION SEQUENCE RX (1), RX (2), RX (19)

...Q + U ==> AJ...
... A + B + F + AJ ==> AR



Q

U

3
STEPS

● HCl




AJ

START NEXT REACTION SEQUENCE



A

B



RX(9) RCT Q 102284-29-7, U 103-71-9
PRO AC 102284-34-4

RX(14) RCT AC 102284-34-4
RGT AF 7647-01-0 HCl
PRO AJ 102284-39-9


RX(1) RCT A 1490-25-1, B 6066-82-6
RGT D 121-44-8 Et3N
PRO C 52787-46-9
SOL 141-78-6 AcOEt

RX(2) RCT C 52787-46-9, F 1948-31-8
RGT H 144-55-8 NaHCO3
PRO G 102284-27-5
SOL 7732-18-5 Water, 67-64-1 Me2CO

RX(19) RCT G 102284-27-5, AJ 102284-39-9
RGT AM 543-27-1 ClCO2Bu-i
PRO AR 92279-30-6

RX(154) OF 173 COMPOSED OF REACTION SEQUENCE RX(3), RX(6), RX(11), RX(16)
AND REACTION SEQUENCE RX(2), RX(16)

...K + L + U ==> AR...
...C + F + AR ==> AL



RX(3) RCT K 15761-39-4, L 123-30-8
RGT N 538-75-0 DCC
PRO M 102284-28-6
SOL 109-99-9 THF

RX(6) RCT M 102284-28-6, U 103-71-9
RGT W 110-86-1 Pyridine
PRO V 102284-31-1
SOL 68-12-2 DMF




RX(11) RCT V 102284-31-1
RGT AF 7647-01-0 HCl
PRO AR 102284-36-6
SOL 109-99-9 THF, 64-18-6 HCO2H

RX(2) RCT C 52787-46-9, F 1948-31-8
RGT H 144-55-8 NaHCO3
PRO G 102284-27-5
SOL 7732-18-5 Water, 67-64-1 Me2CO

RX(16) RCT G 102284-27-5, AR 102284-36-6
RGT AM 543-27-1 ClCO2Bu-i, AN 109-02-4 N-Methylmorpholine
PRO AL 92279-27-1
SOL 109-99-9 THF



RX(155) OF 173 COMPOSED OF REACTION SEQUENCE RX(3), RX(7), RX(12), RX(17)
AND REACTION SEQUENCE RX(2), RX(17)

...K + L + Y ==> AH...
...C + F + AH ==> AO




AR

START NEXT REACTION SEQUENCE





AR




AH

START NEXT REACTION SEQUENCE



AH



RX(3) RCT K 15761-39-4, L 123-30-8
RGT N 538-75-0 DCC
PRO M 102284-28-6
SOL 109-99-9 THF



RX(7) RCT M 102284-28-6, Y 79-44-7
PRO Z 102284-32-2
SOL 110-86-1 Pyridine

RX(12) RCT Z 102284-32-2
RGT AF 7647-01-0 HCl
PRO AH 102284-37-7




RX(2) RCT C 52787-46-9, F 1948-31-8
RGT H 144-55-8 NaHCO₃
PRO G 102284-27-5
SOL 7732-18-5 Water, 67-64-1 Me₂CO

RX(17) RCT G 102284-27-5, AH 102284-37-7
RGT AM 543-27-1 ClCO₂Bu-1, AN 109-02-4 N-Methylmorpholine, AP
10416-59-8 Me₃SiN:MeOSiMe₃
PRO AO 92279-28-2
SOL 109-99-9 THF

RX(156) OF 173 COMPOSED OF REACTION SEQUENCE RX(3), RX(8), RX(13), RX(18)
AND REACTION SEQUENCE RX(2), RX(18)
...K + L + AA ==> AI...
...C + F + AI ==> AQ



START NEXT REACTION SEQUENCE



RX(3) RCT K 15761-39-4, L 123-30-8
RGT N 538-75-0 DCC
PRO M 102284-28-6
SOL 109-99-9 THF

RX(8) RCT M 102284-28-6, AA 1795-48-8
RGT W 110-86-1 Pyridine
PRO AB 102284-33-3
SOL 68-12-2 DMF



RX(13) RCT AB 102284-33-3
RGT AF 7647-01-0 HCl
PRO AI 102284-38-8

RX(2) RCT C 52787-46-9, F 1948-31-8
RGT H 144-55-8 NaHCO₃




PRO G 102284-27-5
SOL 7732-18-5 Water, 67-64-1 Me₂CO

RX(18) RCT G 102284-27-5, AI 102284-38-8
RGT AM 543-27-1 ClCO₂Bu-1
PRO AQ 92279-29-3

RX(157) OF 173 COMPOSED OF REACTION SEQUENCE RX(4), RX(9), RX(14), RX(19)
AND REACTION SEQUENCE RX(2), RX(19)
...K + P + U ==> AJ...
...C + F + AJ ==> AR



START NEXT REACTION SEQUENCE



RX(4) RCT K 15761-39-4, P 95-55-6
RGT N 538-75-0 DCC
PRO Q 102284-29-7
SOL 109-99-9 THF



RX(9) RCT Q 102284-29-7, U 103-71-9
PRO AC 102284-34-4

RX(14) RCT AC 102284-34-4
RGT AF 7647-01-0 HCl
PRO AJ 102284-39-9

RX(2) RCT C 52787-46-9, F 1948-31-8
RGT H 144-55-8 NaHCO₃
PRO G 102284-27-5
SOL 7732-18-5 Water, 67-64-1 Me₂CO

RX(19) RCT G 102284-27-5, AJ 102284-39-9
RGT AM 543-27-1 ClCO₂Bu-1
PRO AR 92279-30-6


RX(161) OF 173 COMPOSED OF REACTION SEQUENCE RX(3), RX(6), RX(11), RX(16)
AND REACTION SEQUENCE RX(1), RX(2), RX(16)
...K + L + U ==> AR...
...A + B + F + AR ==> AL



● HCl


AR

START NEXT REACTION SEQUENCE



A

B




F

AR

● HCl

3 STEPS



AL

RX(3) RCT K 15761-39-4, L 123-30-8
RGT N 538-75-0 DCC
PRO M 102284-28-6
SOL 109-99-9 THF

RX(6) RCT M 102284-28-6, U 103-71-9
RGT W 110-86-1 Pyridine
PRO V 102284-31-1
SOL 68-12-2 DMP

RX(11) RCT V 102284-31-1
RGT AF 7647-01-0 HCl
PRO AR 102284-36-6
SOL 109-99-9 THF, 64-18-6 HCO2H


RX(1) RCT A 1490-25-1, B 6066-82-6
RGT D 121-44-8 Et3N
PRO C 52787-46-9
SOL 141-78-6 AcOEt

RX(2) RCT C 52787-46-9, F 1948-31-8
RGT H 144-55-8 NaHCO3
PRO G 102284-27-5
SOL 7732-18-5 Water, 67-64-1 Me2CO

RX(16) RCT G 102284-27-5, AR 102284-36-6
RGT AM 543-27-1 ClCO2Bu-1, AN 109-02-4 N-Methylmorpholine
PRO AL 92279-27-1
SOL 109-99-9 THF

RX(162) OF 173 COMPOSED OF REACTION SEQUENCE RX(3), RX(7), RX(12), RX(17)
AND REACTION SEQUENCE RX(1), RX(2), RX(17)

...K + L + Y ==> AR...
...A + B + F + AR ==> AO




K

L

Y


3 STEPS



● HCl


AR

START NEXT REACTION SEQUENCE



A

B




F

AR

● HCl

3 STEPS



AO

RX(3) RCT K 15761-39-4, L 123-30-8
RGT N 538-75-0 DCC
PRO M 102284-28-6
SOL 109-99-9 THF

RX(7) RCT M 102284-28-6, Y 79-44-7
PRO Z 102284-32-2
SOL 110-86-1 Pyridine

RX(12) RCT Z 102284-32-2
RGT AF 7647-01-0 HCl
PRO AR 102284-37-7


RX(1) RCT A 1490-25-1, B 6066-82-6
RGT D 121-44-8 Et3N
PRO C 52787-46-9
SOL 141-78-6 AcOEt

RX(2) RCT C 52787-46-9, F 1948-31-8
RGT H 144-55-8 NaHCO3
PRO G 102284-27-5
SOL 7732-18-5 Water, 67-64-1 Me2CO

RX(17) RCT G 102284-27-5, AR 102284-37-7
RGT AM 543-27-1 ClCO2Bu-1, AN 109-02-4 N-Methylmorpholine, AP 10416-59-8 Me3SiN:CMOSiMe3
PRO AO 92279-28-2
SOL 109-99-9 THF

RX(163) OF 173 COMPOSED OF REACTION SEQUENCE RX(3), RX(8), RX(13), RX(18)
AND REACTION SEQUENCE RX(1), RX(2), RX(18)

...K + L + AA ==> AI...
...A + B + F + AI ==> AQ




K

L

AA


3 STEPS




AI

● HCl


START NEXT REACTION SEQUENCE



A




B



F

AI

● HCl

3
STEPS


AQ

RX(3) RCT K 15761-39-4, L 123-30-8
RGT N 538-75-0 DCC
PRO M 102284-28-6
SOL 109-99-9 THF

RX(8) RCT M 102284-28-6, AA 1795-48-8
RGT M 110-86-1 Pyridine
PRO AB 102284-33-3
SOL 68-12-2 DMF

RX(13) RCT AB 102284-33-3
RGT AF 7647-01-0 HCl
PRO AI 102284-38-8


RX(1) RCT A 1490-25-1, B 6066-82-6
RGT D 121-44-8 Et3N
PRO C 52787-46-9
SOL 141-78-6 AcOEt



F

AJ

● HCl

3
STEPS

AR

RX(4) RCT K 15761-39-4, P 95-55-6
RGT N 538-75-0 DCC
PRO Q 102284-29-7
SOL 109-99-9 THF

RX(9) RCT Q 102284-29-7, U 103-71-9
PRO AC 102284-34-4

RX(14) RCT AC 102284-34-4
RGT AF 7647-01-0 HCl
PRO AJ 102284-39-9

RX(1) RCT A 1490-25-1, B 6066-82-6
RGT D 121-44-8 Et3N
PRO C 52787-46-9
SOL 141-78-6 AcOEt

RX(2) RCT C 52787-46-9, F 1948-31-8
RGT H 144-55-8 NaHCO3
PRO Q 102284-27-5
SOL 7732-18-5 Water, 67-64-1 Me2CO

RX(19) RCT Q 102284-27-5, AJ 102284-39-9
RGT AM 543-27-1 ClCO2Bu-i
PRO AR 92279-30-6


L12 ANSWER 16 OF 18 CASREACT COPYRIGHT 2007 ACS on STM
ACCESSION NUMBER: 98:143062 CASREACT Full-text
TITLE: The enantioselective Michael addition of thiole to

RX(2) RCT C 52787-46-9, F 1948-31-8
RGT H 144-55-8 NaHCO3
PRO Q 102284-27-5
SOL 7732-18-5 Water, 67-64-1 Me2CO

RX(18) RCT G 102284-27-5, AI 102284-39-8
RGT AM 543-27-1 ClCO2Bu-i
PRO AQ 92279-29-3

RX(164) OF 173 COMPOSED OF REACTION SEQUENCE RX(4), RX(9), RX(14), RX(19)
AND REACTION SEQUENCE RX(1), RX(2), RX(19)


...K + P + U ==> AJ...
... A + B + F + AJ ==> AR



K

P


U

3
STEPS

AJ

● HCl


START NEXT REACTION SEQUENCE



A


B

cycloalkenones by using (2S,4S)-2-(anilinoethyl)-1-ethyl-4-hydroxypyrrolidine as chiral catalyst
Suzuki, Katsuko; Ikegawa, Akihiko; Mukaiyama, Teruaki
Fac. Sci., Univ. Tokyo, Tokyo, 113, Japan
Bulletin of the Chemical Society of Japan (1982),
55(10), 3277-82
CODEN: BCSJAB; ISSN: 0009-2673
DOCUMENT TYPE: Journal
LANGUAGE: English
OI



AB Catalytic asym. addition of thiole 4-RC6H4(CH2)nSH (R = H, Me, Cl, MeO, Me3C, n = 0; R = H, n = 1) to 2-cycloalkenones I (X = bond, CH2, CH2CH2, CMe2) to give II was studied by using the chiral amino alcs. III (R1 = H, R2 = Ph, cyclohexyl, 2,6-Me2C6H3, 1-naphthyl, R3 = Me; R1 = R3 = H, R2 = Ph, R3 = Me3C) IV (R4 = H, OH), and V derived from L-hydroxyproline or (S)-proline, as base catalysts. The effects of the structure of the catalyst, the reaction medium, the temperature, and the concentration on the enantioselectivity of the reaction was determined. Good optical yields (47-88%) were achieved by the reaction of arenethiole and 2-cyclohexen-1-one in toluene at -5°, by using the catalyst (2S,4S)-2-anilinoethyl-1-ethyl-4-hydroxypyrrolidine.

RX(8) OF 16 ...P + E ==> Q...



P


E

Q

YIELD 74%

RX(8) RCT P 64030-43-9, E 108-24-7
PRO Q 84046-41-3


RX(14) OF 16 COMPOSED OF RX(7), RX(8)
RX(14) O + E ==> Q



RX(7) RCT O 64030-42-8
PRO P 64030-43-9

RX(8) RCT P 64030-43-9, E 108-24-7
PRO Q 84846-41-3

L12 ANSWER 17 OF 18 CASREACT COPYRIGHT 2007 ACS ON STN
ACCESSION NUMBER: 97:71598 CASREACT Full-text
TITLE: Asymmetric synthesis based on chiral diamines having a pyrrolidine ring
AUTHOR(S): Mukaiyama, Teruaki
CORPORATE SOURCE: Dep. Chem., Univ. Tokyo, Tokyo, 113, Japan
SOURCE: Tetrahedron (1981), 37(23), 4111-19
CODEN: TETRAH; ISSN: 0040-4020
DOCUMENT TYPE: Journal
LANGUAGE: English
OI





AB The chiral diamines I-III were used in the asym. preparation of chiral aldehydes and secondary alcs. and in enantioselective addition reactions of thiols with cyclohexenone. Treatment of I with LiAlH₄ in Et₂O at room temperature for 1 h followed by addition of PhCOMe in Et₂O at -100° gave 87% (S)-PhCH(OH)Me. The preparation of the marine antibiotic, (-)-malyngolide (IV), through reaction of I with MeOCH(OH)CO₂Me to give the intermediate V is also reported.

US 4119620	A	19781010	US 1976-733343	19761018
SE 7612064	A	19770501	SE 1976-12064	19761029
NL 7612030	A	19770503	NL 1976-12030	19761029
FR 2329646	A1	19770527	FR 1976-32830	19761029
FR 2329646	B1	19810710		
GB 1518207	A	19780719	GB 1976-45200	19761029
CA 1080216	A1	19800624	CA 1976-264754	19761029
SU 786853	A3	19801207	SU 1976-2415363	19761029
US 4191808	A	19800304	US 1978-897043	19780417
			JP 1975-130809	19751030
			US 1976-733343	19761018

PRIORITY APPLN. INFO.:

AB H-X-Pro-NHCH₂CH₂CH₂ (I; X = Gly, Ala, Asp, Glu, Lys, Arg; R = NO₂; X = Gly, R = N₂NPh) and their HCl or tosylate salts were prepared as enzyme substrates for the diagnostic determination of enzymic activities in various diseases. Thus, 2-Pr-OH (Z = PhCH₂CO₂C) was amidated with p-nitroaniline by P(O)Cl₃ in THF to give the anilide which was Z-deblocked with HBr/HOAc and coupled to 2-Gly-OSu (Su = succinimido) to give 2-Gly-Pro-NHCH₂CH₂CH₂NO₂ (II). The latter was Z-deblocked with HBr/HOAc to give I (X = Gly, R = NO₂) (II) which was treated with p-MeC₆H₄SO₃H to give II tosylate (III). III was used as a substrate for the determination of the enzymic activity of human serum by the photometric determination of the resulting p-nitroaniline. The enzymic activity of the serum from patients suffering from various diseases (e.g., hepatitis) was measured with II.

RX(2) OF 3 ...C + D ==> E




RX(2) RCT C 21027-63-4, D 2699-60-7
PRO E 60189-43-7

=> FIL STNGUIDE
COST IN U.S. DOLLARS


SINCE FILE TOTAL
ENTRY SESSION

RX(2) OF 205 ...D + E ==> F...



RX(2) RCT D 77937-78-1, E 108-24-7
PRO F 77937-79-2
CAT 110-86-1 Pyridine

RX(59) OF 205 COMPOSED OF RX(16), RX(2)
RX(59) AR + E ==> F



RX(16) RCT AR 77937-77-0
PRO D 77937-78-1

RX(2) RCT D 77937-78-1, E 108-24-7
PRO F 77937-79-2
CAT 110-86-1 Pyridine

L12 ANSWER 18 OF 18 CASREACT COPYRIGHT 2007 ACS ON STN
ACCESSION NUMBER: 88:7373 CASREACT Full-text
TITLE: Di-peptide derivatives and method for measuring enzyme activity using these derivatives
INVENTOR(S): Nagatsu, Toshiharu; Sakakibara, Shunpei
PATENT ASSIGNEE(S): Ajinomoto Co., Inc., Japan
SOURCE: Ger. Offen., 35 pp.
CODEN: GWXXBX
DOCUMENT TYPE: Patent
LANGUAGE: German
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 2649171	A1	19770518	DE 1976-2649171	19761028
JP 5205593	A	19770507	JP 1975-130809	19751030
JP 56020839	B	19810515		

FULL ESTIMATED COST	273.09	403.60
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
CA SUBSCRIBER PRICE	ENTRY	SESSION
	-15.33	-16.06

FILE 'STNGUIDE' ENTERED AT 08:56:10 ON 30 MAY 2007
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY, JAPAN SCIENCE AND TECHNOLOGY CORPORATION, AND FACHINFORMATIONSZENTRUM KARLSRUHE

FILE CONTAINS CURRENT INFORMATION.
LAST RELOADED: May 25, 2007 (20070525/UP).

=> file reg		
COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	0.76	404.38
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
CA SUBSCRIBER PRICE	ENTRY	SESSION
	0.00	-16.06

FILE 'REGISTRY' ENTERED AT 09:04:03 ON 30 MAY 2007
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGE TERMS" FOR DETAILS.
COPYRIGHT (C) 2007 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 28 MAY 2007 HIGHEST RN 935999-19-2
DICTIONARY FILE UPDATES: 28 MAY 2007 HIGHEST RN 935999-19-2

New CAS Information Use Policies, enter HELP USAGE TERMS for details.



TS/CA INFORMATION NOW CURRENT THROUGH December 2, 2006

Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stndoc/properties.html>

>> Uploading C:\Program Files\Stnexp\Queries\10.561754\formula VII.str




chain nodes :
1 2 8 9 10 11 14 15 16 17
ring nodes :
3 4 5 6 7
ring/chain nodes :
12 13
chain bonds :
1-4 1-2 1-12 3-8 8-9 8-10 10-11 12-13 13-14 14-15 14-16 16-17
ring bonds :
3-4 3-7 4-5 5-6 6-7
exact/norm bonds :
1-4 1-2 1-12 3-4 3-7 3-8 4-5 5-6 6-7 8-9 8-10 10-11 12-13 13-14 14-15
14-16 16-17

G1:C,S

Match level :
1:CLASS 2:CLASS 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:CLASS 9:CLASS 10:CLASS
11:Atom 12:CLASS 13:CLASS 14:CLASS 15:CLASS 16:CLASS 17:CLASS

L13 STRUCTURE UPLOADED

>> d
L13 HAS NO ANSWERS
L13 STR



Structure attributes must be viewed using STN Express query preparation.

>> s l13 sss sam
SAMPLE SEARCH INITIATED 09:04:29 FILE 'REGISTRY'
SAMPLE SCREEN SEARCH COMPLETED 23 TO ITERATE
100.0% PROCESSED 23 ITERATIONS 4 ANSWERS
SEARCH TIME: 00.00.01


FULL FILE PROJECTIONS: ONLINE **COMPLETE**
BATCH **COMPLETE**
PROJECTED ITERATIONS: 173 TO 747
PROJECTED ANSWERS: 4 TO 200

L14 4 SEA SSS SAM L13

>> d scan

L14 4 ANSWERS REGISTRY COPYRIGHT 2007 ACS on STN
IN L-Prolineamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-2-pyridinyl-
MP C18 H26 N4 O4

Absolute stereochemistry.




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):1

L14 4 ANSWERS REGISTRY COPYRIGHT 2007 ACS on STN
IN L-Prolineamide, (2R)-2-butyl-N-formyl-N-(phenylmethoxy)-β-alanyl-N-(4-

methoxy-2-pyridinyl)- (9CI)
MF C26 H34 N4 O5

Absolute stereochemistry.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):0

>> file hcplus
COST IN U.S. DOLLARS SINCE FILE TOTAL
ENTRY SESSION
FULL ESTIMATED COST 0.45 404.83
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS) SINCE FILE TOTAL
ENTRY SESSION
CA SUBSCRIBER PRICE 0.00 -16.06

FILE 'HCPLUS' ENTERED AT 09:04:43 ON 30 MAY 2007
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 30 May 2007 VOL 146 ISS 23
FILE LAST UPDATED: 29 May 2007 (20070529/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

>> s l14
L15 4 L14

>> d ibib

L15 ANSWER 1 OF 4 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2006.1177562 HCAPLUS Full-text
DOCUMENT NUMBER: 146.92498
TITLE: Peptide deformylase inhibitors as potent antimicrobial agents
AUTHOR(S): Teo, Jeanette W. P.; Thayalan, Pamela; Beer, David; Yap, Amelia S. L.; Nanjundappa, Mahesh; Ngew, Kinyi; Duraiswamy, Jeyarej; Liung, Sarah; Dartois, Veronique; Schreiber, Mark; Hasan, Samiul; Cynamon, Michael; Ryder, Neil S.; Yang, Xia; Weidmann, Beat; Bracken, Kathryn; Dick, Thomas; Mukherjee, Kakoli
CORPORATE SOURCE: Novartis Institute for Tropical Diseases, Singapore, 138670, Singapore
SOURCE: Antimicrobial Agents and Chemotherapy (2006), 50(11), 3665-3673
CODEN: AMACQJ; ISSN: 0066-4804
PUBLISHER: American Society for Microbiology
DOCUMENT TYPE: Journal
LANGUAGE: English
REFERENCE COUNT: 64 THERE ARE 64 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

>> d ibib 2-4

L15 ANSWER 2 OF 4 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2006.111167 HCAPLUS Full-text
DOCUMENT NUMBER: 144.131762
TITLE: Method for increasing the susceptibility of peptide deformylase inhibitors by using efflux pump inhibitors
INVENTOR(S): Dean, Charles Richard; Ryder, Neil Stewart
PATENT ASSIGNEE(S): Novartis AG, Swiss; Novartis Pharma GmbH
SOURCE: PCT Int. Appl., 51 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2006002896	A1	20060112	WO 2005-EP7008	20050629
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SI, SM, SN, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RM: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CP, CG, CI, CM, GA, GN, GQ, GM, ML, MR, NE, SN, TD, TO, BM, OH, OM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KO, KZ, MD, RU, TJ, TM				
AU 2005259488	A1	20060112	AU 2005-259488	20050629
CA 2569681	A1	20061206	CA 2005-2569681	20050629
EP 1763348	A1	20070321	EP 2005-772146	20050629
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR				
PRIORITY APPL. INFO:			US 2004-584023P	P 20040630
			WO 2005-EP7008	W 20050629
OTHER SOURCE(S):		MARPAT 144:121762		

10/561754 385/447 Robert Havlin
REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT
L15 ANSWER 3 OF 4 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2005:714250 HCAPLUS Full-text
DOCUMENT NUMBER: 143:322091
TITLE: Role of the AcrAB-TolC efflux pump in determining susceptibility of Haemophilus influenzae to the novel peptide deformylase inhibitor LBM415
AUTHOR(S): Dean, Charles R.; Narayan, Shubha; Daigle, Denise M.; Dzink-Fox, JoAnn L.; Puyang, Xiaoling; Bracken, Kathryn R.; Dean, Karl E.; Weidmann, Beat; Yuan, Zhengyu; Jain, Rakesh; Ryder, Neil
CORPORATE SOURCE: Novartis Institutes for Biomedical Research, Inc., Cambridge, MA, 02139, USA
SOURCE: Antimicrobial Agents and Chemotherapy (2005), 49(8), 3129-3135
CODEN: AMACQ; ISSN: 0066-4804
PUBLISHER: American Society for Microbiology
DOCUMENT TYPE: Journal
LANGUAGE: English
REFERENCE COUNT: 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L15 ANSWER 4 OF 4 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2002:977804 HCAPLUS Full-text
DOCUMENT NUMBER: 138:55863
TITLE: Preparation of N-formyl-N-hydroxylamino-substituted pyrrolidine derivatives as inhibitors of peptidyl deformylase
INVENTOR(S): Patel, Dinesh V.; Yuan, Zhengyu; Jain, Rakesh K.; Garcia Alvarez, Salvador; Jacobs, Jeffrey
PATENT ASSIGNEE(S): Versicor, Inc., USA; Novartis AG
SOURCE: PCT Int. Appl., 69 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002102790	A1	20021227	WO 2002-EP6504	20020614
M: AR, AG, AL, AM, AT, AU, AZ, BA, BE, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LT, LU, LV, MA, MD, MK, MN, MX, NO, NZ, OM, PH, PL, PT, RO, RU, SE, SG, SI, SK, TJ, TM, TN, TR, TT, UA, US, UZ, VN, YU, ZA, ZW				
RM: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR				
CA 2448526	A1	20021227	CA 2002-2448526	20020614
AU 2002321062	A1	20030102	AU 2002-321062	20020614
US 2003048479	A1	20030306	US 2002-171706	20020614
US 7148242	B2	20061212		
EP 1401828	A1	20040331	EP 2002-754681	20020614
EP 1401828	B1	20060412		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
HU 200400208	A2	20040628	HU 2004-208	20020614
CN 1511152	A	20040707	CN 2002-810596	20020614
BR 2002010377	A	20040810	BR 2002-10377	20020614
JP 200502606	T	20050127	JP 2003-506263	20020614
NZ 529489	A	20051028	NZ 2002-529489	20020614

10/561754 387/447 Robert Havlin
FILE 'CASREACT' ENTERED AT 08:21:56 ON 30 MAY 2007
L5 50 S L4
FILE 'CASREACT' ENTERED AT 08:24:06 ON 30 MAY 2007
L6 28 S L5 NOT PY-2003
FILE 'REGISTRY' ENTERED AT 08:28:44 ON 30 MAY 2007
L7 STRUCTURE UPLOADED
FILE 'CASREACT' ENTERED AT 08:29:13 ON 30 MAY 2007
L8 5 S L7
L9 3 S L8 NOT PY > 2003
L10 46 S L7 SSS FULL
L11 21 S L10 NOT PY > 2003
L12 18 S L11 NOT L9
FILE 'STNGUIDE' ENTERED AT 08:56:10 ON 30 MAY 2007
FILE 'REGISTRY' ENTERED AT 09:04:03 ON 30 MAY 2007
L13 STRUCTURE UPLOADED
L14 4 S L13 SSS SAM
FILE 'HCAPLUS' ENTERED AT 09:04:43 ON 30 MAY 2007
L15 4 S L14
FILE 'REGISTRY' ENTERED AT 09:05:46 ON 30 MAY 2007
-- s l13 sss full
FULL SEARCH INITIATED 09:05:55 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 444 TO ITERATE
100.0% PROCESSED 444 ITERATIONS 67 ANSWERS
SEARCH TIME: 00.00.01
L16 67 SRA SSS FUL L13
-- file hcaplus
COST IN U.S. DOLLARS SINCE FILE ENTRY TOTAL
FULL ESTIMATED COST 172.10 586.85
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS) SINCE FILE ENTRY TOTAL
CA SUBSCRIBER PRICE 0.00 -16.06
FILE 'HAPLUS' ENTERED AT 09:06:01 ON 30 MAY 2007
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.
FILE COVERS 1907 - 30 May 2007 VOL 146 ISS 23

10/561754 386/447 Robert Havlin
AT 323081 T 20060415 AT 2002-754681 20020614
PT 1401828 T 20060931 PT 2002-754681 20020614
ES 2262824 T3 20061201 ES 2002-2754681 20020614
ZA 2003008379 A 20040521 ZA 2003-8379 20031028
IN 2003CN01963 A 20060106 IN 2003-CN1963 20031210
NO 2003005571 A 20040216 NO 2003-5571 20031212
HK 1064370 A1 20061020 HK 2004-107013 20040914
PRIORITY APPLN. INFO.: US 2001-296419P P 20010615
US 2002-360313P P 20020227
WO 2002-EP6604 W 20020614
OTHER SOURCE(S): MARPAT 138:55863
REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

SINCE FILE ENTRY	TOTAL
ENTRY	SESSION
9.92	414.75
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE ENTRY
0.00	SESSION
CA SUBSCRIBER PRICE	-16.06

FILE 'REGISTRY' ENTERED AT 09:05:46 ON 30 MAY 2007
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2007 AMERICAN CHEMICAL SOCIETY (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 28 MAY 2007 HIGHEST RN 935999-19-2
DICTIONARY FILE UPDATES: 28 MAY 2007 HIGHEST RN 935999-19-2

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TECA INFORMATION NOW CURRENT THROUGH December 2, 2006

Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stndoc/properties.html>

-- d his

(FILE 'HOME' ENTERED AT 08:18:24 ON 30 MAY 2007)

FILE 'REGISTRY' ENTERED AT 08:18:37 ON 30 MAY 2007
L1 STRUCTURE UPLOADED

FILE 'CASREACT' ENTERED AT 08:19:13 ON 30 MAY 2007
L2 0 S L1 SSS SAM
L3 1 S L1 SSS FULL

FILE 'REGISTRY' ENTERED AT 08:21:37 ON 30 MAY 2007
L4 STRUCTURE UPLOADED

10/561754 388/447 Robert Havlin
FILE LAST UPDATED: 29 May 2007 (20070529/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

-- s l16

L17 32 L16

-- d ibib abs hitstr tot

L17 ANSWER 1 OF 32 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2007:269020 HCAPLUS Full-text
DOCUMENT NUMBER: 146:418201


TITLE: Reduced susceptibility of Haemophilus influenzae to the peptide deformylase inhibitor LBM415 can result from target protein overexpression due to amplified chromosomal def gene copy number

AUTHOR(S): Dean, Charles R.; Narayan, Shubha; Richards, Joel; Daigle, Denise M.; Esterow, Stacy; Leeds, Jennifer A.; Kemp, Heather; Puyang, Xiaoling; Wiedmann, Brigitte; Mueller, Dieter; Voshol, Hans; van Oostrum, Jan; Wall, Daniel; Koehn, James; Dzink-Fox, JoAnn; Ryder, Neil S.
CORPORATE SOURCE: Infectious Diseases, Novartis Institute for Biomedical Research, Cambridge, MA, 02139, USA
SOURCE: Antimicrobial Agents and Chemotherapy (2007), 51(3), 1004-1010
CODEN: AMACQ; ISSN: 0066-4804
PUBLISHER: American Society for Microbiology
DOCUMENT TYPE: Journal
LANGUAGE: English

AB Previous genetic anal. of Haemophilus influenzae revealed two mechanisms associated with decreased susceptibility to the novel peptide deformylase inhibitor LBM415: AcrAB-TolC-mediated efflux and Fmt bypass, resulting from mutations in the pump repressor gene acrR and in the fmt gene, resp. The authors have isolated an addnl. mutant, CDS23 (LBM415 MIC 64 µg/mL vs. 4 µg/mL against the parent strain NB65044) that lacks mutations in the acrR or fmt structural genes or in the gene encoding Def, the intracellular target of LBM415. Western immunoblot anal., two-dimensional gel electrophoresis, and tryptic digestion combined with mass spectrometric identification showed that the Def protein was highly overexpressed in the mutant strain. Consistent with this, real-time reverse transcription-PCR revealed a significant increase in def transcript titer. No mutations were found in the region upstream of def that might account for altered expression; however, pulsed-field gel electrophoresis suggested that a genetic rearrangement of the region containing def had occurred. Using a combination of PCR, sequencing, and Southern blot analyses, it was determined that the def gene had undergone copy number amplification, explaining the high level of target protein expression. Inactivation of the AcrAB-TolC efflux pump in this mutant increased susceptibility 16-fold, highlighting the role of efflux in exacerbating the overall reduced susceptibility resulting from target overexpression.


IT 475913-91-6, LBM415
RL: BSU (Biological study, unclassified); PAC (Pharmacological activity); BIOL (Biological study)
(reduced susceptibility of Haemophilus influenzae to peptide deformylase inhibitor LBM415 can result from target protein overexpression due to amplified chromosomal def gene copy number)
RN 475913-91-6 HCAPLUS
CN L-Prolineamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 34 THERE ARE 34 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L17 ANSWER 2 OF 32 HCAPLUS COPYRIGHT 2007 ACS ON STN
ACCESSION NUMBER: 2007:148028 HCAPLUS Full-text
DOCUMENT NUMBER: 146:180650
TITLE: Multistep resistance selection and postantibiotic-effect studies of the antipneumococcal activity of LBM415 compared to other agents
AUTHOR(S): Kosowska-Shick, Klaudia; Credito, Kim L.; Pankuch, Glenn A.; DeWasse, Bonifacio; McGhee, Pamela; Appelbaum, Peter C.
CORPORATE SOURCE: Department of Pathology, Hershey Medical Center, Hershey, PA, 17033, USA
SOURCE: Antimicrobial Agents and Chemotherapy (2007), 51(2), 770-773
CODEN: AMACQJ; ISSN: 0066-4804
PUBLISHER: American Society for Microbiology
DOCUMENT TYPE: Journal
LANGUAGE: English
AB LBM415 is a peptide deformylase inhibitor active against gram-pos. bacterial species and some gram-neg. species. In multiselection studies, LBM415 had low MICs against all Streptococcus pneumoniae strains tested, regardless of their genotype, and selected resistant clones after 14 to 50 days. MIC increases correlated with changes mostly in the 700KAAKQ77 motif in peptide deformylase. The postantibiotic effect of LBM415 ranged from 0.3 to 1.4 h.
IT 478913-91-6, LBM415
RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(postantibiotic effect of peptide deformylase inhibitor LBM415 against Streptococcus pneumoniae)
RN 478913-91-6 HCAPLUS
CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)
Absolute stereochemistry.



REFERENCE COUNT: 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS

Absolute stereochemistry.



● 1/2 Mg

REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L17 ANSWER 4 OF 32 HCAPLUS COPYRIGHT 2007 ACS ON STN
ACCESSION NUMBER: 2006:1250645 HCAPLUS Full-text
DOCUMENT NUMBER: 146:45733
TITLE: Preparation of N-formylhydroxylamine-containing peptides
INVENTOR(S): Bracken, Kathryn Rene; Bushell, Simon; Dean, Karl; Francavilla, Charles; Jain, Rakesh K.; Lee, Kwangho; Seepersaud, Mohindra; Shu, Lei; Sundaram, Arathia; Yuan, Zhengyu
PATENT ASSIGNEE(S): Novartis A.-G., Switz.; Novartis Pharma G.m.b.H.; Vicuron Pharmaceuticals, Inc
SOURCE: PCT Int. Appl., 61pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2006127576	A2	20061130	WO 2006-US19688	20060522
WO 2006127576	A3	20070125		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NO, NI, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, DZ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TO, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			


PRIORITY APPL. INFO.: US 2005-663655P P 20050523
OTHER SOURCE(S): MARPAT 146:45733
OI

RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L17 ANSWER 3 OF 32 HCAPLUS COPYRIGHT 2007 ACS ON STN
ACCESSION NUMBER: 2006:1263091 HCAPLUS Full-text
DOCUMENT NUMBER: 146:184267
TITLE: β-Amino amides from β-lactams: application to the formal synthesis of a peptide-deformylase inhibitor
AUTHOR(S): Jiang, Xinglong; Prasad, Kape; Prasad, Mahavir; Slade, Joel; Repic, Oljan; Blacklock, Thomas J.
CORPORATE SOURCE: Process Research & Development, Novartis Pharmaceuticals Corporation, East Hanover, NJ, 07936, USA
SOURCE: Synlett (2006), (18), 3179-3181
CODEN: SYNLES; ISSN: 0936-5214
PUBLISHER: Georg Thieme Verlag
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 146:184267
OI




AB A facile and a practical synthesis of I, an intermediate for a peptide-deformylase inhibitor, is described using an acid-catalyzed aminolysis of a β-lactam with a pyrrolidine derivative as the key transformation. In addition, simplified conditions for the conversion of a β-hydroxy acid to a β-lactam are reported.
IT 478913-92-7P 771478-93-2P
RL: SPN (Synthetic preparation); PREP (Preparation)
(formal synthesis of peptide-deformylase inhibitor via aminolysis of β-lactam)
RN 478913-92-7 HCAPLUS
CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-(phenylmethoxy)-β-alanyl-N-(5-fluoro-2-pyridinyl)- (9CI) (CA INDEX NAME)
Absolute stereochemistry. Rotation (-).




RN 771478-83-2 HCAPLUS
CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-oxido-2-pyridinyl)-, magnesium salt (2:1) (9CI) (CA INDEX NAME)



AB The invention relates to novel N-formyl hydroxylamine compds. I (R1 is H, alkyl, heteroalkyl, heterocycloalkyl, aryl, or heteroaryl; R3 is H, halo, or alkoxy; R4 is aryl or heteroaryl; n is 0-3) or their salts or prodrugs that inhibit peptidyl deformylase (PDF), an enzyme present in prokaryotes, and are useful as antimicrobials and antibiotics. Examples describe syntheses of title compds. and intermediates, e.g., for the preparation of I (n = 1, R1 = cyclopentyl, R3 = H, R4 = 5-fluoro-N-oxido-2-pyridyl). Compds. of the invention were assayed for inhibition of PDF and for antimicrobial activity (e.g., min. inhibitory concns. approx. 0.25-32 μg/mL against H. influenza).
IT 915280-68-1P 915280-69-2P 915280-72-7P 915280-75-0P 915280-77-2P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation of N-formylhydroxylamine-containing peptides as inhibitors of peptidyl deformylase)
RN 915280-68-1 HCAPLUS
CN 2-Pyrrolidinecarboxamide, 4-fluoro-N-(5-fluoro-2-pyridinyl)-1-[(2R)-2-[[formyl(phenylmethoxy)amino]methyl]-1-oxohexyl]-, (2S,4R)- (CA INDEX NAME)
Absolute stereochemistry.




RN 915280-69-2 HCAPLUS
CN 2-Pyrrolidinecarboxamide, 4-fluoro-N-(5-fluoro-1-oxido-2-pyridinyl)-1-[(2R)-2-[[formyl(phenylmethoxy)amino]methyl]-1-oxohexyl]-, (2S,4R)- (CA INDEX NAME)
Absolute stereochemistry.




RN 915280-72-7 HCAPLUS
CN 2-Pyrrolidinecarboxamide, 1-[(2R)-2-[[[formyl][(tetrahydro-2H-pyran-2-yl)oxy]amino]methyl]-1-oxohexyl]-N-3-pyridazinyl-, (2S)- (CA INDEX NAME)

Absolute stereochemistry.




RN 915280-75-0 HCAPLUS
CN 2-Pyrrolidinecarboxamide, 1-[(2R)-3-cyclobutyl-2-[[[formyl][(tetrahydro-2H-pyran-2-yl)oxy]amino]methyl]-1-oxopropyl]-4-fluoro-N-(2-oxido-3-pyridazinyl)-, (2S,4R)- (CA INDEX NAME)

Absolute stereochemistry.



RN 915280-77-2 HCAPLUS
CN 2-Pyrrolidinecarboxamide, 1-[(2R)-3-cyclobutyl-2-[[[formyl][(tetrahydro-2H-pyran-2-yl)oxy]amino]methyl]-1-oxopropyl]-4-fluoro-N-(2-oxido-3-pyridazinyl)-, (2S,4R)- (CA INDEX NAME)


Absolute stereochemistry.



IT 478912-97-9, PDF 709
RL: PAC (Pharmacological activity); PKT (Pharmacokinetics); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(peptide deformylase inhibitors as antimycobacterial agents)

RN 478912-97-9 HCAPLUS
CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(4-ethyl-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 64 THERE ARE 64 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L17 ANSWER 6 OF 32 HCAPLUS COPYRIGHT 2007 ACS ON STN
ACCESSION NUMBER: 2006:446747 HCAPLUS [Full-text](#)
DOCUMENT NUMBER: 145:184169
TITLE: Activity of LBM415 compared to those of 11 other agents against Haemophilus species
AUTHOR(S): Bogdanovich, Tatiana; Smith, Kathy A.; Clark, Catherine; Pankuch, Glenn A.; Lin, Gengrong; McGehee, Pamela; Dewasse, Bonifacio; Appelbaum, Peter C.
CORPORATE SOURCE: Hershey Medical Center, Hershey, PA, 17033, USA
SOURCE: Antimicrobial Agents and Chemotherapy (2006), 50(7), 2323-2329
CODEN: AMACQJ; ISSN: 0066-4804
PUBLISHER: American Society for Microbiology
DOCUMENT TYPE: Journal
LANGUAGE: English

AB When tested against 254 Haemophilus influenzae strains, LBM415, a peptide deformylase inhibitor, gave MIC50 and MIC90 values of 2.0 µg/mL and 8.0 µg/mL, resp. The MICs were independent of β-lactam or quinolone susceptibility and the presence or absence of macrolide efflux or ribosomal protein mutations. The MICs of LBM415 against 23 H. parainfluenzae strains were similar to those against H. influenzae. In contrast, erythromycin, azithromycin, and clarithromycin gave unimodal MIC distributions, and apart from β-lactamase-neg., ampicillin-resistant strains, all strains were susceptible to the β-lactams tested. Apart from selected quinolone-resistant strains, all strains were susceptible to ciprofloxacin, levofloxacin, gemifloxacin, and gemifloxacin. Resistance to trimethoprim-sulfamethoxazole was common. The potencies of all drugs



L17 ANSWER 5 OF 32 HCAPLUS COPYRIGHT 2007 ACS ON STN
ACCESSION NUMBER: 2006:1177562 HCAPLUS [Full-text](#)
DOCUMENT NUMBER: 146:92698

TITLE: Peptide deformylase inhibitors as potent antimycobacterial agents
AUTHOR(S):

Teo, Jeanette W. P.; Thayalan, Pamela; Beer, David; Yap, Amelia S. L.; Nanjundappa, Mahesh; Ngew, Xinyi; Duraiswamy, Jeyaraj; Liang, Sarah; Dartois, Veronique; Schreiber, Mark; Hasan, Samiul; Cynamon, Michael; Ryder, Neil S.; Yang, Xia; Weidmann, Beat; Bracken, Kathryn; Dick, Thomas; Mukherjee, Kakoli

CORPORATE SOURCE: Novartis Institute for Tropical Diseases, Singapore, 138670, Singapore
SOURCE: Antimicrobial Agents and Chemotherapy (2006), 50(11), 3665-3673

CODEN: AMACQJ; ISSN: 0066-4804
PUBLISHER: American Society for Microbiology
DOCUMENT TYPE: Journal
LANGUAGE: English

AB Peptide deformylase (PDF) catalyzes the hydrolytic removal of the N-terminal formyl group from nascent proteins. This is an essential step in bacterial protein synthesis, making PDF an attractive target for antibacterial drug development. Essentiality of the def gene, encoding PDF from Mycobacterium tuberculosis, was demonstrated through genetic knockout expts. with Mycobacterium bovis BCG. PDF from M. tuberculosis strain H37Rv was cloned, expressed, and purified as an N-terminal histidine-tagged recombinant protein in Escherichia coli. A novel class of PDF inhibitors (PDF-I), the N-alkyl urea hydroxamic acids, were synthesized and evaluated for their activities against the M. tuberculosis PDF enzyme as well as their antimycobacterial effects. Several compds. from the new class had 50% inhibitory concentration (IC50) values of <100 nM. Some of the PDF-I displayed antibacterial activity against M. tuberculosis, including MDR strains with MIC90 values of <1 µM. Pharmacokinetic studies of potential leads showed that the compds. were orally bioavailable. Spontaneous resistance towards these inhibitors arose at a frequency of 5.5-10⁻⁷ in M. bovis BCG. DNA sequence anal. of several spontaneous PDF-I-resistant mutants revealed that half of the mutants had acquired point mutations in their formyl transferase gene (fmt), which formylated Met-tRNA. The results from this study validate M. tuberculosis PDF as a drug target and suggest that this class of compds. have the potential to be developed as novel antimycobacterial agents.

IT 478912-45-7, LBM-611
RL: PAC (Pharmacological activity); PKT (Pharmacokinetics); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(PDF-611; peptide deformylase inhibitors as antimycobacterial agents)

RN 478912-45-7 HCAPLUS
CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(2-pyridinyl)- (CA INDEX NAME)


Absolute stereochemistry.

against 23 H. parainfluenzae strains were similar to those against H. influenzae. Time-kill studies with 10 Haemophilus strains showed LBM415 to be bactericidal at 2 × the MIC against 8 of 10 strains after 24 h. For comparison, the macrolides and β-lactams were bactericidal against 8 to 10 strains each at 2 × the MIC after 24 h. Quinolones were bactericidal against all 10 strains tested at 2 × the MIC after 24 h. Against six H. influenzae strains, postantibiotic effects for LBM415 lasted between 0.8 and 2.2 h. In multi-step resistance selection studies, LBM415 produced resistant clones in 7 of the 10 strains tested, with MICs ranging from 4 to 64 µg/mL. No mutations in deformylase (def) and formyltransferase (fmt) genes were detected in any of the LBM415-resistant mutants.

IT 478913-91-6, LBM415
RL: BSU (Biological study, unclassified); BIOL (Biological study)
(activity of LBM415 compared to other agents against Haemophilus species)

RN 478913-91-6 HCAPLUS
CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 32 THERE ARE 32 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L17 ANSWER 7 OF 32 HCAPLUS COPYRIGHT 2007 ACS ON STN
ACCESSION NUMBER: 2006:446747 HCAPLUS [Full-text](#)
DOCUMENT NUMBER: 144:484688
TITLE: Preclinical study of peptide deformylase inhibition in Streptococcus pneumoniae and Staphylococcus aureus
AUTHOR(S): Wang, Wen; White, Richard; Yuan, Zhengyu
CORPORATE SOURCE: Vicuron Pharmaceuticals, Fremont, CA, 94555, USA
SOURCE: Antimicrobial Agents and Chemotherapy (2006), 50(5), 1656-1663
CODEN: AMACQJ; ISSN: 0066-4804
PUBLISHER: American Society for Microbiology
DOCUMENT TYPE: Journal
LANGUAGE: English

AB Peptide deformylase (PDF) is an essential enzyme in both gram-neg. and gram-pos. bacteria. It hydrolyzes formylated N-terminal peptides to generate free N-terminal peptides during the process of protein maturation. Inhibition of this enzyme results in cessation of bacterial growth. We have examined the effect of a potent PDF inhibitor, LBM-415 (also known as VIC-104959), on the proteomes of Staphylococcus aureus and Streptococcus pneumoniae using two-dimensional electrophoresis. Both S. aureus and S. pneumoniae showed accumulation of many N-terminal formylated peptides/proteins upon PDF inhibition. In S. pneumoniae, formylated peptide/protein accumulation was time dependent. Following inhibition, subsequent removal of the inhibitor resulted in deformation of formylated peptides/proteins; this recovery process was also time dependent. If instead the inhibited cells were maintained in the presence of sub-MIC levels of the PDF inhibitor, the formylated peptides/proteins remained for a much longer time, which correlated with a prolonged postantibiotic effect in vitro. These observations may have broader implications for the application of this class of antibiotics in vivo.

IT 478913-91-6, LBM-415

RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (PDF inhibitor; proteomic study of peptide deformylase inhibition in
 Streptococcus pneumoniae and Staphylococcus aureus)
 RN 478913-91-6 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-
 oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 22 THERE ARE 22 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L17 ANSWER 8 OF 32 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2006:332553 HCAPLUS [Full-text](#)
 DOCUMENT NUMBER: 145:79591
 TITLE: Antimicrobial activity of a novel peptide deformylase inhibitor, LBM415, tested against respiratory tract and cutaneous infection pathogens: a global surveillance report (2003-2004)
 AUTHOR(S): Watters, Amy A.; Jones, Ronald N.; Leeds, Jennifer A.; Denys, Gerald; Sader, Helio S.; Fritsche, Thomas R.
 CORPORATE SOURCE: JMI Laboratories, North Liberty, IA, 52317, USA
 SOURCE: Journal of Antimicrobial Chemotherapy (2006), 57(5), 914-923
 CODEN: JACHDX; ISSN: 0305-7453
 PUBLISHER: Oxford University Press
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB To evaluate the spectrum of activity and potency of LBM415, the 1st of the peptide deformylase inhibitor (PDFI) class to be developed for treatment of community-acquired respiratory tract infections and uncomplicated skin and soft tissue infections (uSTTI), against a large, contemporary international collection of targeted pathogens collected during 2003-2004. A total of 21 636 isolates were tested by reference broth microdilution methods as part of a longitudinal international antimicrobial resistance surveillance study. Characteristics of the organism collection included resistance to oxacillin among 35.0% of *Staphylococcus aureus* and 76.0% of coagulase-neg. staphylococci (CONS); resistance to penicillin (MIC ≥ 2 mg/L) among 18.0% of *Streptococcus pneumoniae*; vancomycin resistance among 20.0% of *Enterococcus* spp. and ampicillin resistance among 22.0% of *Haemophilus influenzae*. LBM415 displayed potent activity against staphylococci, streptococci, *Enterococcus faecium* and *Moraxella catarrhalis*, with 299.0% of strains being inhibited at 54 mg/L; 97.0% of *Enterococcus faecalis* isolates and 92.0% of *H. influenzae* isolates were also inhibited at this concentration. Seventy-seven % of *Burkholderia cepacia* and 82.0% of *Stenotrophomonas maltophilia* were inhibited at 58 mg/L. No differences in LBM415 activity against *S. aureus*, CONS, *S. pneumoniae*, *Enterococcus* spp. and *H. influenzae* were detected for subsets susceptible or resistant to antimicrobials such as oxacillin, penicillin, ampicillin, macrolides, vancomycin and fluoroquinolones. While regional differences were apparent with some comparator agents, sensitivity to LBM415 did not vary significantly among strains from the various geog. areas sampled. One isolate of *S. aureus* displayed high-level resistance to LBM415 owing to multiple sequence changes in resistance phenotype genes (*defB* and *fat*), despite the absence of the compound

in clin. practice. This isolate remained susceptible to all other antimicrobials tested except for penicillin. With few differences detected among strains from various geog. regions, the 1st PDFI class agent to enter clin. development has consistently demonstrated a broad spectrum of activity against commonly isolated pathogens associated with uncomplicated respiratory and cutaneous infections. These compds. represent a significant therapeutic advance owing to their novel mechanism of action and antibacterial spectrum, including activity against resistant organisms, should pharmacokinetic and pharmacodynamic parameters support their continued development. Given the detection of a pre-existing PDFI-resistant isolate of *S. aureus* as demonstrated here, surveillance for resistance among the PDFI-targeted pathogens following introduction of this class of agent into clin. usage will be an important component of future studies.
 IT 478913-91-6, LBM415
 RL: BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (antibiotic activity of peptide deformylase inhibitor LBM415 against respiratory tract and cutaneous infection pathogens)
 RN 478913-91-6 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L17 ANSWER 9 OF 32 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2006:206088 HCAPLUS [Full-text](#)
 DOCUMENT NUMBER: 144:424953
 TITLE: Natural products - the future scaffolds for novel antibiotics?
 AUTHOR(S): Butler, Mark S.; Buss, Antony D.
 CORPORATE SOURCE: MerLion Pharmaceuticals, Singapore, 117528, Singapore
 SOURCE: Biochemical Pharmacology (2006), 71(7), 919-929
 CODEN: BCPAC6; ISSN: 0006-2952
 PUBLISHER: Elsevier B.V.
 DOCUMENT TYPE: Journal; General Review
 LANGUAGE: English
 AB A review. Natural products have played a pivotal role in antibiotic drug discovery with most antibacterial drugs being derived from a natural product or natural product lead. However, the rapid onset of resistance to most antibacterial drugs diminishes their effectiveness considerably and necessitates a constant supply of new antibiotics for effective treatment of infections. The natural product templates of actinonin, pleuromutilin, ramoplanin and tiamcinacin B, which are compds. undergoing clin. evaluation, represent templates not found in currently marketed antibacterial drugs. In addition, the new templates present in the recently discovered lead antibacterials arylomycin, GE23077, manogepinycin, muraymycin/caprazamycin, nocathiacin and ECO-0501, are discussed. Despite extensive efforts to identify antibiotic leads from mol. targets, only the peptide deformylase inhibitor LBM-415 is currently in clin. trials. It is proposed that new antibacterial assays which combine cell-based screening with mol. targets could offer better prospects for lead discovery.
 IT 478913-91-6, LBM 415

RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (natural products)
 RN 478913-91-6 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 100 THERE ARE 100 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L17 ANSWER 10 OF 32 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2006:31167 HCAPLUS [Full-text](#)
 DOCUMENT NUMBER: 144:121762
 TITLE: Method for increasing the susceptibility of peptide deformylase inhibitors by using efflux pump inhibitors
 INVENTOR(S): Dean, Charles Richard; Ryder, Neil Stewart
 PATENT ASSIGNEE(S): Novartis AG, Swiss.; Novartis Pharma GmbH
 SOURCE: PCT Int. Appl., 51 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2006002896	A1	20060112	NO 2005-EP7008	20050629
M:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BU, CF, CG, CI, CM, GA, GN, GO, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
AU 2005259488	A1	20060112	AU 2005-259488	20050629
CA 2569681	A1	20061206	CA 2005-2569681	20050629
EP 1763348	R	20070321	EP 2005-72146	20050629
R:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR			
PRIORITY APPLN. INFO.:			US 2004-584023P	P 20040630
			WO 2005-EP7008	W 20050629
OTHER SOURCE(S):			MARPAT 144:121762	


AB The invention provides methods and compds. for increasing the susceptibility of peptide deformylase inhibitors against Gram-neg. organisms by using efflux pump inhibitors. Related methods for the treatment of Gram-neg. bacterial infections are also provided.
 IT 478912-45-7, LBR 611 478913-91-6
 RL: BSU (Biological study, unclassified); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (method for increasing the susceptibility of peptide deformylase inhibitors by using efflux pump inhibitors)
 RN 478912-45-7 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-2-pyridinyl- (CA INDEX NAME)

Absolute stereochemistry.



RN 478913-91-6 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L17 ANSWER 11 OF 32 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2005:1093778 HCAPLUS [Full-text](#)
 DOCUMENT NUMBER: 143:359507
 TITLE: Evaluation of LBM415 (NVP PDF-713), a novel peptide deformylase inhibitor, for treatment of experimental Mycoplasma pneumoniae pneumonia
 AUTHOR(S): Fonseca-Aten, Monica; Salvatore, Christine M.; Mejias, Anuncion; Rios, Ana M.; Chavez-Bueno, Susana; Katz, Kathy; Gomez, Ana M.; McCracken, George H., Jr.; Hardy, R. Doug
 CORPORATE SOURCE: Department of Pediatrics, University of Texas Southwestern Medical Center, Dallas, TX, 75390-9063, USA
 SOURCE: Antimicrobial Agents and Chemotherapy (2005), 49(10), 4128-4136


CODEN: AMACQ; ISSN: 0066-4804
 PUBLISHER: American Society for Microbiology
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB Mycoplasma pneumoniae is a major cause of community-acquired pneumonia. We evaluated the efficacy of LBM415, a novel peptide deformylase inhibitor antimicrobial agent, for the treatment of M. pneumoniae pneumonia in a mouse model. Eight-week-old BALB/c mice were intranasally inoculated once with 10⁷ CFU of M. pneumoniae. Groups of mice were treated with LBM415 (50 mg/kg of body weight) or placebo s.c. daily for 13 days, starting 24 h after inoculation. Groups of mice were evaluated at the baseline; at days of treatment 1, 3, 6, and 13; and at 7 days after treatment. The MIC of LBM415 against M. pneumoniae was <0.005 µg/mL. LBM415-treated mice had significantly lower bronchoalveolar lavage fluid M. pneumoniae concns. than placebo-treated mice on days 6 and 13 of treatment. Compared with placebo treatment, therapy with LBM415 significantly decreased lung histopathol. scores at days 3, 6, and 13 of treatment and at 7 days after treatment. Airway obstruction was significantly lower in LBM415-treated mice than in placebo-treated mice on days 1, 3, and 6 of treatment and after 7 days of therapy, while airway hyperresponsiveness was significantly lower only on day 3 of therapy. The bronchoalveolar lavage fluid concns. of tumor necrosis factor alpha, gamma interferon (IFN-γ), interleukin-6 (IL-6), IL-12, KC (functional IL-8), monocyte chemoattractant protein 1, macrophage inflammatory protein 1α, monokine induced by IFN-γ, and IFN-inducible protein 10 were significantly reduced in LBM415-treated mice compared with the levels in placebo-treated mice. There were no differences in the bronchoalveolar lavage fluid concns. of granulocyte-macrophage colony-stimulating factor, IL-1β, IL-2, IL-4, IL-5, and IL-10 between the two groups of mice. LBM415 therapy had beneficial microbiol., histol., respiratory, and immunol. effects on acute murine M. pneumoniae pneumonia.

IT 478913-91-6, LBM415
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USBS (Uses)
 (LBM415 (NVP PDF-713), a novel peptide deformylase inhibitor, for treatment of exptl. Mycoplasma pneumoniae pneumonia)

RN 478913-91-6 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 45 THERE ARE 45 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L17 ANSWER 12 OF 32 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2005:714250 HCAPLUS [Full-text](#)
 DOCUMENT NUMBER: 143:322091
 TITLE: Role of the AcrAB-TolC efflux pump in determining susceptibility of Haemophilus influenzae to the novel peptide deformylase inhibitor LBM415
 AUTHOR(S): Dean, Charles R.; Narayan, Shubha; Daigle, Denis M.; Dzink-Fox, JoAnn L.; Puyang, Xiaoling; Bracken, Kathryn R.; Dean, Karl E.; Weidmann, Beat; Yuan, Zhengyu; Jain, Rakesh; Ryder, Neil S.

CORPORATE SOURCE: Novartis Institutes for Biomedical Research, Inc., Cambridge, MA, 02139, USA
 SOURCE: Antimicrobial Agents and Chemotherapy (2005), 49(8), 3129-3135
 CODEN: AMACQ; ISSN: 0066-4804


PUBLISHER: American Society for Microbiology
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB Haemophilus influenzae isolates vary widely in their susceptibilities to the peptide deformylase inhibitor LBM415 (MIC range, 0.06 to 32 µg/mL); however, on average, they are less susceptible than gram-pos. organisms, such as Staphylococcus aureus and Streptococcus pneumoniae. Insertional inactivation of the H. influenzae acrB or tolC gene in strain NB65044 (Rd strain KM20) increased susceptibility to LBM415, confirming a role for the AcrAB-TolC pump in determining resistance. Consistent with this, sequencing of a PCR fragment generated with primers flanking the acrA region from an LBM415-hypersusceptible H. influenzae clin. isolate revealed a genetic deletion of acrA. Inactivation of acrB or tolC in several clin. isolates with atypically reduced susceptibility to LBM415 (MIC of 16 µg/mL or greater) significantly increased susceptibility, confirming that the pump is also a determinant of decreased susceptibility in these clin. isolates. Examination of acrR, encoding the putative repressor of pump gene expression, from several of these strains revealed mutations introducing frameshifts, stop codons, and amino acid changes relative to the published sequence, suggesting that loss of pump repression leads to decreased susceptibility. Supporting this, NB65044 acrR mutants selected by exposure to LBM415 at 8 µg/mL had susceptibilities to LBM415 and other pump substrates comparable to the least sensitive clin. isolates and showed increased expression of pump genes.

IT 478912-45-7, LSK 611
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (AcrR is repressor of AcrAB expression, and mutations in acrR are related to susceptibility to LBM415)

RN 478912-45-7 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-2-pyridinyl- (CA INDEX NAME)

Absolute stereochemistry.



IT 478913-91-6, LBM415
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (role of AcrAB-TolC efflux pump in determining susceptibility of Haemophilus influenzae to novel peptide deformylase inhibitor LBM415 and structurally related LSK611)

RN 478913-91-6 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


CODEN: AMACQ; ISSN: 0066-4804
 PUBLISHER: American Society for Microbiology
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB Objectives: To compare the in vitro activity of NVP-LMB415 (formerly referred to as NVP-PDF 713) with that of other agents with anti-anaerobic activity against clin. anaerobic isolates, with emphasis on the Bacteroides fragilis group. The MICs for 405 B. fragilis group and 102 Gram-pos. anaerobic isolates were determined using NCCLS-recommended procedures. The activity of NVP-LMB415 was compared with that of cefoxitin, clindamycin, isipenem, garenoxacin, linezolid, moxifloxacin and tigecycline. Vancomycin was included in the evaluation of the Gram-pos. organisms. NVP-LMB415 showed excellent in vitro activity against all the species of the B. fragilis group isolates (MIC range ≤ 0.03-0.5 mg/L and MIC90 0.5 mg/L). NVP-LMB415 was active against B. fragilis group strains resistant to β-lactams, quinolones or clindamycin, and the MICs were much lower than those of newer agents such as linezolid, tigecycline and garenoxacin. The MICs of NVP-LMB415 (2-4 mg/L) for Clostridium species were higher than the MICs for other anaerobes. Given the frequency of isolation of anaerobic bacteria and their increasing resistance to all classes of antibiotics, NVP-LMB415 is an ideal agent for potential use against mixed infections caused by resistant anaerobic pathogens such as B. fragilis and Gram-pos. aerobic strains such as methicillin-resistant staphylococci, streptococci and enterococci.

IT 478913-91-6, NVP-PDF 713
 RL: BSU (Biological study, unclassified); DMA (Drug mechanism of action); THU (Therapeutic use); BIOL (Biological study); USBS (Uses)
 (Evaluation of the in vitro activity of NVP-LMB415 against clin. anaerobic isolates with emphasis on the Bacteroides fragilis group)

RN 478913-91-6 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L17 ANSWER 13 OF 32 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2005:602838 HCAPLUS [Full-text](#)
 DOCUMENT NUMBER: 143:169417
 TITLE: Evaluation of the in vitro activity of NVP-LMB415 against clinical anaerobic isolates with emphasis on the Bacteroides fragilis group
 AUTHOR(S): Snyderman, David R.; Jacobus, Nilda V.; McDermott, Laura A.
 CORPORATE SOURCE: Tufts-New England Medical Center, Boston, MA, 02111, USA
 SOURCE: Journal of Antimicrobial Chemotherapy (2005), 55(6), 1024-1028
 CODEN: JACHDX; ISSN: 0305-7453
 PUBLISHER: Oxford University Press
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB A review. Peptide deformylase (PDF) is a prokaryotic metalloenzyme that is essential for bacterial growth but is not required by mammalian cells. Thus, it represents a selective and promising target for the development of new antibacterial agents. Since deformylase inhibitors have yet to be used clin. as antibacterial drugs, compds. targeting this enzyme should avoid cross-resistance with currently used antibacterial agents. The PDF enzyme is a ferrous ion-containing metallohydrolase, but a nickel-containing surrogate is routinely used in the laboratory for testing inhibitors due to its better stability. Enzymes from several bacterial species have been cloned and both their three-dimensional structures and co-crystal structures with bound inhibitor have been determined. As a metallo enzyme, PDF lends itself to the well-precedented mechanism-based rational drug design approach. Using structural and mechanistic information together with high throughput screening, several types of potent PDF inhibitors have been identified. PDF inhibitors identified to date share a common structural feature of a "chelator + peptidomimetic" scaffold. Although compds. with many different chelators inhibit the cell free enzyme, only compds. containing hydroxamic acid or N-formyl hydroxylamine exhibit appreciable antibacterial activity. Several lead inhibitors have demonstrated in vivo efficacy and an excellent safety profile. Two PDF inhibitors, VIC-104959 (LBM415) and BB-83698, have progressed to Phase I clin. trials. In this review, different PDF inhibitors are compared and their biol. activities are discussed. Structure-activity relationships have been established and the implications of this work in the design of future PDF inhibitors are considered.

IT 478913-91-6, VIC 104959
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USBS (Uses)
 (bacterial peptide deformylase inhibitors as a new class of antibacterial agents)

RN 478913-91-6 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)

CORPORATE SOURCE: Vicuron Pharmaceuticals, Fremont, CA, 94555, USA
 SOURCE: Current Medicinal Chemistry (2005), 12(14), 1607-1621
 CODEN: CMCHET; ISSN: 0929-8673
 PUBLISHER: Bentham Science Publishers Ltd.
 DOCUMENT TYPE: Journal; General Review
 LANGUAGE: English


L17 ANSWER 14 OF 32 HCAPLUS COPYRIGHT 2007 ACS on STN
 ACCESSION NUMBER: 2005:553883 HCAPLUS [Full-text](#)
 DOCUMENT NUMBER: 143:221622
 TITLE: Bacterial peptide deformylase inhibitors: A new class of antibacterial agents
 AUTHOR(S): Jain, R.; Chen, D.; White, R. J.; Patel, D. V.; Yuan, Z.
 CORPORATE SOURCE: Vicuron Pharmaceuticals, Fremont, CA, 94555, USA
 SOURCE: Current Medicinal Chemistry (2005), 12(14), 1607-1621
 CODEN: CMCHET; ISSN: 0929-8673
 PUBLISHER: Bentham Science Publishers Ltd.
 DOCUMENT TYPE: Journal; General Review
 LANGUAGE: English


AB A review. Peptide deformylase (PDF) is a prokaryotic metalloenzyme that is essential for bacterial growth but is not required by mammalian cells. Thus, it represents a selective and promising target for the development of new antibacterial agents. Since deformylase inhibitors have yet to be used clin. as antibacterial drugs, compds. targeting this enzyme should avoid cross-resistance with currently used antibacterial agents. The PDF enzyme is a ferrous ion-containing metallohydrolase, but a nickel-containing surrogate is routinely used in the laboratory for testing inhibitors due to its better stability. Enzymes from several bacterial species have been cloned and both their three-dimensional structures and co-crystal structures with bound inhibitor have been determined. As a metallo enzyme, PDF lends itself to the well-precedented mechanism-based rational drug design approach. Using structural and mechanistic information together with high throughput screening, several types of potent PDF inhibitors have been identified. PDF inhibitors identified to date share a common structural feature of a "chelator + peptidomimetic" scaffold. Although compds. with many different chelators inhibit the cell free enzyme, only compds. containing hydroxamic acid or N-formyl hydroxylamine exhibit appreciable antibacterial activity. Several lead inhibitors have demonstrated in vivo efficacy and an excellent safety profile. Two PDF inhibitors, VIC-104959 (LBM415) and BB-83698, have progressed to Phase I clin. trials. In this review, different PDF inhibitors are compared and their biol. activities are discussed. Structure-activity relationships have been established and the implications of this work in the design of future PDF inhibitors are considered.

IT 478913-91-6, VIC 104959
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USBS (Uses)
 (bacterial peptide deformylase inhibitors as a new class of antibacterial agents)

RN 478913-91-6 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.






REFERENCE COUNT: 76 THERE ARE 76 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L17 ANSWER 15 OF 32 HCAPLUS COPYRIGHT 2007 ACS ON STN
 ACCESSION NUMBER: 2005:494346 HCAPLUS [Full-text](#)
 DOCUMENT NUMBER: 143:149764
 TITLE: Comparative in vitro activities of investigational peptide deformylase inhibitor NVP LBM-415 and other agents against human mycoplasmas and ureaplasmas
 AUTHOR(S): Waites, Ken B.; Reddy, Nipun B.; Crabb, Donna M.; Duffy, Lynn B.
 CORPORATE SOURCE: Department of Pathology, University of Alabama at Birmingham, Birmingham, AL, 35249, USA
 SOURCE: Antimicrobial Agents and Chemotherapy (2005), 49(6), 2541-2542
 CODEN: AMACCO; ISSN: 0066-4804
 PUBLISHER: American Society for Microbiology
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB Peptide deformylase inhibitor LBM-415 and seven other drugs were tested against Mycoplasma pneumoniae (100 isolates), Mycoplasma hominis (20 isolates), Mycoplasma fermentans (10 isolates), and Ureaplasma species (50 isolates). LBM-415 was active against M. pneumoniae (MICs, 50.008 µg/mL). It showed no activity against M. hominis and M. fermentans and modest activity against Ureaplasma spp.
 IT 478913-91-6, LBM-415
 RL: BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)
 (comparative in vitro activities of investigational peptide deformylase inhibitor NVP LBM-415 and other agents against human mycoplasmas and ureaplasmas)
 RN 478913-91-6 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

activity has been widely evaluated in preclin. studies against multiple pathogens, including drug-resistant strains. In vitro studies using recent clin. isolates have demonstrated potent activity against streptococcal and staphylococcal strains responsible for community-acquired respiratory tract infections and skin infections. LBM-415 is also active against medically important groups of drug-resistant pathogens, including methicillin-resistant Staphylococcus aureus (MRSA), penicillin-resistant Streptococcus pneumoniae, vancomycin-resistant enterococci and clarithromycin-resistant Helicobacter pylori. The efficacy of LBM-415 has been demonstrated in mouse models of infection, where it was active against Mycoplasma pneumoniae-induced pneumonia, and had comparable efficacy to linezolid and vancomycin against systemic MRSA and methicillin-susceptible S. aureus (MSSA). Pharmacokinetic studies, including single- and multiple-dose studies in humans, demonstrated linear kinetics, with rapid absorption of LBM-415 and no evidence of accumulation. The compound is advancing to phase II/III clin. trials.

IT 478913-91-6, LBM 415
 RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (peptide deformylase inhibitor LBM-415 was active against drug-resistant pathogens, was effective in mouse models of infection and had linear kinetics with rapid absorption and no evidence of accumulation in human)

RN 478913-91-6 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 41 THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L17 ANSWER 18 OF 32 HCAPLUS COPYRIGHT 2007 ACS ON STN
 ACCESSION NUMBER: 2005:325026 HCAPLUS [Full-text](#)
 DOCUMENT NUMBER: 143:22918
 TITLE: Comparative antimicrobial characterization of LBM415 (NVP PDF-713), a new peptide deformylase inhibitor of clinical importance
 AUTHOR(S): Fritsche, Thomas R.; Sader, Helio S.; Cleeland, Roy; Jones, Ronald N.
 CORPORATE SOURCE: The JONES Group/JMI Laboratories, North Liberty, IA, 53117, USA
 SOURCE: Antimicrobial Agents and Chemotherapy (2005), 49(4), 1468-1476
 CODEN: AMACCO; ISSN: 0066-4804
 PUBLISHER: American Society for Microbiology
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 QT


L17 ANSWER 16 OF 32 HCAPLUS COPYRIGHT 2007 ACS ON STN
 ACCESSION NUMBER: 2005:494343 HCAPLUS [Full-text](#)
 DOCUMENT NUMBER: 143:149763
 TITLE: In vitro and intracellular activities of LBM415 (NVP PDF-713) against Legionella pneumophila
 AUTHOR(S): Edelman, Paul H.; Hu, Baofeng; Edelman, Martha A. C.
 CORPORATE SOURCE: Departments of Pathology and Laboratory Medicine, University of Pennsylvania School of Medicine, Philadelphia, PA, 19104-4283, USA
 SOURCE: Antimicrobial Agents and Chemotherapy (2005), 49(6), 2533-2535
 CODEN: AMACCO; ISSN: 0066-4804
 PUBLISHER: American Society for Microbiology
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB LBM415 activity against extracellular and intracellular L. pneumophila was studied. The LBM415 MIC50 for 20 Legionella sp. strains was 4 µg/mL, vs. 0.06, 0.25, and ≤ 0.03 µg/mL for azithromycin, erythromycin, and levofloxacin, resp. LBM415 (0.5 and 16 µg/mL) reversibly prevented intracellular growth of 2 L. pneumophila strains and was less active than erythromycin.
 IT 478913-91-6, LBM415
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (in vitro and intracellular activities of LBM415 against Legionella pneumophila)
 RN 478913-91-6 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L17 ANSWER 17 OF 32 HCAPLUS COPYRIGHT 2007 ACS ON STN
 ACCESSION NUMBER: 2005:375159 HCAPLUS [Full-text](#)
 DOCUMENT NUMBER: 142:475186
 TITLE: LBM-415: Antibacterial peptide deformylase inhibitor
 AUTHOR(S): McIntyre, J. A.; Castaner, J.; Martin, L.
 CORPORATE SOURCE: Prous Science, Barcelona, 08080, Spain
 SOURCE: Drugs of the Future (2005), 30(1), 23-28
 CODEN: DRFUD4; ISSN: 0377-8282
 PUBLISHER: Prous Science
 DOCUMENT TYPE: Journal; General Review
 LANGUAGE: English
 AB A review. Resistance among bacterial pathogens has necessitated the search for novel targets in antimicrobial research. The peptide deformylase inhibitors are a novel and unique class of antimicrobial agents in development for the treatment of respiratory tract and skin infections. LBM-415 is the first such compound to enter clin. development. Its



AB LBM415 (NVP PDF-713) (I) is the first member of the peptide deformylase (PDF) inhibitor class being developed for clin. trials as a parenteral and oral agent for treatment of community-acquired respiratory tract disease and serious infections caused by antimicrobial-resistant gram-pos. cocci. In this study, susceptibility testing results from 1,306 recent clin. isolates selected to overrepresent resistance trends among the species were summarized. All staphylococci (153 strains; MIC at which 90% of isolates were inhibited (MIC90), 2 µg/mL), Streptococcus pneumoniae (170 strains; MIC90, 1 µg/mL), other streptococci (150 strains; MIC90, 1 µg/mL), enterococci (104 strains; MIC90, 4 µg/mL), Moraxella catarrhalis (103 strains; MIC90, 0.5 µg/mL), and Legionella pneumophila (50 strains; MIC90, 0.12 µg/mL) were inhibited at ≤ 8 µg of LBM415/mL, as were 97% of Haemophilus influenzae isolates (300 strains; MIC90, 4 to 8 µg/mL). Among other bacterial groups, 100% of gram-pos. and -neg. anaerobes, including 22 Bacteroides spp. strains (31 strains total; MIC90, 1 µg/mL), were inhibited by 54 µg/mL, whereas Enterobacteriaceae (112 strains) and most nonfermentative bacilli (107 strains) were not inhibited at readily achievable concns. The compound was found to have a dominantly bacteriostatic action, and spontaneous single-step mutational rates occurred at low levels (10⁻⁶ to <10⁻⁸). Drug interaction studies failed to identify any class-specific synergistic interactions, nor were antagonistic interactions observed. Variations in broth and agar MIC test conditions demonstrated that, whereas the agar-based method trended towards a 1-log₂ dilution-higher MIC than the broth method and was inoculum dependent, other variations in incubation environment, medium supplements, pH, or calcium concentration had little influence on LBM415 MIC results. Use of the efflux inhibitor phe-arg-β-naphthylamide showed an average of 1 log₂ dilution decrease in H. influenzae MICs, demonstrating the contribution of efflux pumps in influencing susceptibility to PDF inhibitors. The in vitro activity of LBM415 against targeted bacterial species, including resistant subsets, and other laboratory characteristics of this novel compound demonstrate the potential of PDF inhibitors as a new class of antimicrobial agents.

IT 478913-91-6, LBM415
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (comparative antibacterial characterization of LBM415 as new peptide deformylase inhibitor of clin. importance)
 RN 478913-91-6 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 34 THERE ARE 34 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L17 ANSWER 19 OF 32 HCAPLUS COPYRIGHT 2007 ACS ON STN
 ACCESSION NUMBER: 2005:109471 HCAPLUS Full-text
 DOCUMENT NUMBER: 142:459965
 TITLE: Antimicrobial activity of LBM415 (NVP PDF-713) tested against pathogenic *Neisseria* spp. (*Neisseria gonorrhoeae* and *Neisseria meningitidis*)
 AUTHOR(S): Jones, Ronald N.; Sader, Helio S.; Fritsche, Thomas R.
 CORPORATE SOURCE: The JONES Group, JMI Laboratories Inc., North Liberty, IA, 52117, USA
 SOURCE: Diagnostic Microbiology and Infectious Disease (2005), 51(2), 139-141
 CODEN: DMIIDZ; ISSN: 0732-6893
 PUBLISHER: Elsevier Inc.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB LBM415 (NVP PDF-713), a novel peptide deformylase inhibitor, was tested by reference methods against 2 collections of pathogenic *Neisseria*, *N. gonorrhoeae* (157 strains) and *N. meningitidis* (100 strains). The collection included strains resistant to penicillin, tetracycline, and fluoroquinolones and were also tested against ceftriaxone, ciprofloxacin, penicillin, and tetracycline. The 50% and 90% min. inhibitory concentration values for LBM415 were 1 and 2 µg/mL, and 4 and 8 µg/mL for *N. meningitidis* and *N. gonorrhoeae*, resp. All comparison agents were more active than this peptide deformylase inhibitor against this genus.
 IT 478913-91-6, LBM 415
 RL: BSU (Biological study, unclassified); DMA (Drug mechanism of action); THU (Therapeutic use); BIOL (Biological study); USXS (Uses)
 (antimicrobial activity of deformylase inhibitor LBM415 (NVP PDF-713) against pathogenic *Neisseria gonorrhoeae* and *N. meningitidis*)
 RN 478913-91-6 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 19 THERE ARE 19 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L17 ANSWER 20 OF 32 HCAPLUS COPYRIGHT 2007 ACS ON STN
 ACCESSION NUMBER: 2005:88905 HCAPLUS Full-text
 DOCUMENT NUMBER: 143:93836
 TITLE: Activity of a peptide deformylase inhibitor LBM415 (NVP PDF-713) tested against recent clinical isolates from Japan
 AUTHOR(S): Bell, Jan M.; Turnidge, John D.; Inoue, Matsuhisa; Kohno, Shigeru; Hirakata, Yoichi; Ono, Yasuo; Jones, Ronald N.
 CORPORATE SOURCE: Women's and Children's Hospital, Adelaide, Australia
 SOURCE: Journal of Antimicrobial Chemotherapy (2005), 55(2),

276-278
 CODEN: JACHDK; ISSN: 0305-7453
 PUBLISHER: Oxford University Press
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB The potency of LBM415, a new peptide deformylase (PDF) inhibitor, against key Gram-pos. pathogens, as well as *Haemophilus influenzae*, from Japan was assessed. A total of 695 clin. isolates originally collected in Japan included *Staphylococcus aureus* (n=222), *Haemophilus influenzae*, *Streptococcus pneumoniae* (n=122), coagulase-neg. staphylococci (CoNS, n=119), *Enterococcus* spp. (n=65) and *Streptococcus* spp. (n=48). Oxacillin-resistant *S. aureus* had slightly lower LBM415 MIC values than oxacillin-susceptible strains. MIC50 and MIC90 values of LBM415 against oxacillin-resistant *S. aureus* were 2 log10 dilns. lower than previous findings. CoNS had similar MIC results to *S. aureus*, although oxacillin-resistant strains appeared to be less susceptible than oxacillin-susceptible strains. All enterococci were inhibited at ≤8 mg/L of LBM415, all *S. pneumoniae* were inhibited at ≤2 mg/L of the PDF inhibitor, regardless of penicillin or multi-drug resistance. The LBM415 MIC90 for the β-hemolytic streptococci was 0.5 mg/L; all streptococci were inhibited at ≤4 mg/L. These results indicate that LBM415 appears to be an active agent that may be suitable for the treatment of infections, caused by Gram-pos. organisms.
 IT 478913-91-6, LBM 415
 RL: BSU (Biological study, unclassified); BIOL (Biological study)
 (activity of peptide deformylase inhibitor LBM415 (NVP PDF-713) tested against recent clin. isolates from Japan)
 RN 478913-91-6 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L17 ANSWER 21 OF 32 HCAPLUS COPYRIGHT 2007 ACS ON STN
 ACCESSION NUMBER: 2005:14391 HCAPLUS Full-text
 DOCUMENT NUMBER: 142:113909
 TITLE: Process for preparation of N-[oxidopyridinyl] L-prolinamide derivatives
 INVENTOR(S): Slade, Joel; Vivaldo, James Anthony; Chen, Guang-Pei; Bajwa, Joginder Singh; Parker, David John
 PATENT ASSIGNER(S): Novartis A.-G., Swiss.; Novartis Pharma G.m.b.H.
 SOURCE: PCT Int. Appl., 34 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
NO 2005000835	A1	20050106	NO 2004-EP6915	20040625

W: AR, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, GU, HD, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SV, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW
 RM: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TO
 AU 2004251876 A1 20050106 AU 2004-251876 20040625
 CA 2530142 A1 20050106 CA 2004-2530142 20040625
 EP 1641778 A1 20060405 EP 2004-740324 20040625
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IS, SI, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK
 BR 2004011921 A 20060815 BR 2004-11921 20040625
 CN 1829719 A 20060906 CN 2004-80021438 20040625
 US 2007060753 A1 20070315 US 2005-561754 20051221
 PRIORITY APPL. INFO.: US 2003-482686P P 20030626
 WO 2004-EP6915 W 20040625
 OTHER SOURCE(S): CASREACT 142:113909; MARPAT 142:113909
 OI




AB A process for the preparation of title compds. of formula I [Y = a OH protecting group; R1 = (hetero)aryl; R2-R5 = independently H or alkyl, or R2R3 and/or R4R5 = cycloalkyl; X = CH2, S, CH(OH), etc.; n = 0-3] is disclosed. For example, contacting IiTeOH with 1N Na2CO3 in EtOAc to move TeOH and oxidation by H2O2 gave III (R = H). Formylation of III with formic acetic anhydride gave III (R = CHO). Reaction of III with HBr salt of N-(5-fluoro-2-pyridinyl)-2-pyrrolidinedicarboxamide, followed by oxidation, gave IV. Thus, the present invention provides a process producing the title compound, which are useful to prepare certain antibacterial N-formyl hydroxylamine compds. as peptide deformylase inhibitors.
 IT 478913-92-7P
 RL: IMP (Industrial manufacture); RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (preparation of N-[oxidopyridinyl] L-prolinamide derive.)
 RN 478913-92-7 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-(phenylmethoxy)-β-alanyl-N-(5-fluoro-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).

IT 478913-93-8P
 RL: IMP (Industrial manufacture); SPN (Synthetic preparation); PREP (Preparation)
 (preparation of N-[oxidopyridinyl] L-prolinamide derive.)
 RN 478913-93-8 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-(phenylmethoxy)-β-alanyl-N-(5-fluoro-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L17 ANSWER 22 OF 32 HCAPLUS COPYRIGHT 2007 ACS ON STN
 ACCESSION NUMBER: 2004:915738 HCAPLUS Full-text
 DOCUMENT NUMBER: 142:35071
 TITLE: Commercial broth microdilution panel validation and reproducibility trials for NVP PDF-713 (LBM 415), a novel inhibitor of bacterial peptide deformylase
 AUTHOR(S): Fritsche, T. R.; Moet, G. J.; Jones, R. N.
 CORPORATE SOURCE: The JONES Group/JMI Laboratories, North Liberty, IA, USA
 SOURCE: Clinical Microbiology and Infection (2004), 10(9), 857-860
 CODEN: CMINFM; ISSN: 1198-743X
 PUBLISHER: Blackwell Publishing Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB NVP PDF-713 (LBM 415) is a peptide deformylase inhibitor being progressed into clin. trials. Dry-form broth microdilution panels of NVP PDF-713 were compared to reference MIC panels of 552 recent clin. isolates. Most (99.2%) dry-form MIC results were within ± 1 log2 dilution of the reference panel MICs. Of the bacteria tested, *Streptococcus pneumoniae* and *Haemophilus influenzae* showed a bias towards higher and lower MICs, resp. Same-day and between-day reproducibility tests showed that 98.9% and 96.7% of MIC values, resp., were within ± 1 log2 dilution step, thereby demonstrating a high degree of reliability of the dry-form MIC product for clin. studies.

IT 478913-91-6
 RL: BSU (Biological study, unclassified); DMA (Drug mechanism of action);
 THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (antimicrobial activity of peptide deformylase inhibitor NVP PDP-713
 against respiratory tract pathogens)
 RN 478913-91-6 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-
 oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L17 ANSWER 23 OF 32 HCAPLUS COPYRIGHT 2007 ACS ON STN
 ACCESSION NUMBER: 2004:857380 HCAPLUS [Full-text](#)
 DOCUMENT NUMBER: 141:337761
 TITLE: Crystalline N-formylhydroxylamine compounds for
 pharmaceuticals
 INVENTOR(S): Mueller, Martin; Liu, Hui; Bajwa, Joginder Singh
 PATENT ASSIGNEE(S): Novartis Ag, Switz.; Novartis Pharma GmbH; Slade, Joel
 SOURCE: PCT Int. Appl., 36 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
MO 2004087133	A1	20041014	MO 2004-EP3478	20040401
W: AE, AQ, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MM, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BM, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KO, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2004226815	A1	20041014	AU 2004-226815	20040401
CA 2520682	A1	20041014	CA 2004-2520682	20040401
EP 1613305	A1	20060111	EP 2004-725014	20040401
R: AT, BR, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IS, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR				
BR 2004009009	A	20060338	BR 2004-9009	20040401
CN 1764450	A	20060426	CN 2004-80007872	20040401
JP 2006522054	T	20060928	JP 2006-504952	20040401
NO 2005005097	A	20051222	NO 2005-5097	20051101

PRIORITY APPLN. INFO.: US 2003-459726P P 20030402
 WO 2004-EP3478 A 20040401
 OTHER SOURCE(S): MARPAT 141:337761
 AB Certain N-formylhydroxylamine compds., such as N-(1-oxo-2-alkyl-3-(N-
 hydroxyformamido)propyl)[carbonylaminoaryl]azacycloalkanes are useful in the treatment of
 bacterial infections. Disclosed are crystalline salts of such compds. Thus, a capsule
 contained a N-formylhydroxylamine 200, spray-dried lactose 148, and Mg stearate 2 mg. 1-
 [2-R-[[[formylhydroxylamino)methyl]hexanoyl]pyrrolidone-2S-carboxylic acid-(4-
 ethylpyridin-2-yl)amide was converted to its calcium salt by reaction with CaCl2 solution
 in NaOH solution
 IT 771478-83-2P
 RL: PRP (Properties); EPN (Synthetic preparation); THU (Therapeutic use);
 BIOL (Biological study); PREP (Preparation); USES (Uses)
 (crystalline N-formylhydroxylamine compds. for pharmaceuticals)
 RN 771478-83-2 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-
 oxido-2-pyridinyl)-, magnesium salt (2:1) (9CI) (CA INDEX NAME)


Absolute stereochemistry.



●1/2 Mg

IT 478912-97-9 478913-91-6
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (crystalline N-formylhydroxylamine compds. for pharmaceuticals)
 RN 478912-97-9 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(4-ethyl-2-
 pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 478913-91-6 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-
 oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

IT 771478-82-1P 771478-84-3P 771478-85-4P
 RL: EPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological
 study); PREP (Preparation); USES (Uses)
 (crystalline N-formylhydroxylamine compds. for pharmaceuticals)
 RN 771478-82-1 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(4-ethyl-2-
 pyridinyl)-, calcium salt (2:1) (9CI) (CA INDEX NAME)


Absolute stereochemistry.



●1/2 Ca

RN 771478-84-3 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-
 oxido-2-pyridinyl)-, zinc salt (2:1) (9CI) (CA INDEX NAME)


Absolute stereochemistry.



●1/2 Zn

RN 771478-85-4 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-
 oxido-2-pyridinyl)-, calcium salt (2:1) (9CI) (CA INDEX NAME)

Absolute stereochemistry.



●1/2 Ca


REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L17 ANSWER 24 OF 32 HCAPLUS COPYRIGHT 2007 ACS ON STN
 ACCESSION NUMBER: 2004:848356 HCAPLUS [Full-text](#)
 DOCUMENT NUMBER: 142:19798
 TITLE: Antistaphylococcal activity of LBM415, a new peptide
 deformylase inhibitor, compared with those of other
 agents
 AUTHOR(S): Credito, Kim; Lin, Gengrong; Ednie, Lois M.;
 Appelbaum, Peter C.
 CORPORATE SOURCE: Department of Pathology, Hershey Medical Center,
 Hershey, PA, USA
 SOURCE: Antimicrobial Agents and Chemotherapy (2004), 48(10),
 4033-4036
 CODEN: AMACQJ; ISSN: 0066-4804
 PUBLISHER: American Society for Microbiology
 DOCUMENT TYPE: Journal
 LANGUAGE: English

AB The MICs of LBM415, a new peptide deformylase inhibitor, were 50.06 to 4.0 μg/mL for 258
 isolates of Staphylococcus aureus and coagulase-neg. staphylococci. LBM415 MICs were
 similar irres. of whether the strains were methicillin susceptible or resistant. All
 strains were also susceptible to vancomycin, linezolid, ranbezolid, daptomycin,
 oritavancin, and quinupristin-dalfopristin. LBM415 at the MIC was bacteriostatic after 24
 h.

IT 478913-91-6, LBM 415
 RL: BSU (Biological study, unclassified); PRP (Properties); BIOL
 (Biological study)
 (antistaphylococcal activity of LBM415, new peptide deformylase
 inhibitor, compared with those of other agents)
 RN 478913-91-6 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-
 oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L17 ANSWER 25 OF 32 HCAPLUS COPYRIGHT 2007 ACS ON STN
 ACCESSION NUMBER: 2004:848355 HCAPLUS [Full-text](#)
 DOCUMENT NUMBER: 142:3338
 TITLE: Antipneumococcal activity of LBM415, a new peptide diformylase inhibitor, compared with those of other agents
 AUTHOR(S): Ednie, Lois M.; Pankuch, Glenn; Appelbaum, Peter C.
 CORPORATE SOURCE: Department of Pathology, Hershey Medical Center, Hershey, PA, USA
 SOURCE: Antimicrobial Agents and Chemotherapy (2004), 48(10), 4027-4032
 CODEN: AMACQ; ISSN: 0066-4804
 PUBLISHER: American Society for Microbiology
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB The MICs of LBM415, a new peptide diformylase inhibitor, were evaluated and ranged from 0.03 to 4.0 µg/mL for 300 pneumococci, irres. of their β-lactam, macrolide, and quinolone susceptibilities. By comparison, vancomycin, teicoplanin, linezolid, and quinupristin-dalfopristin were also active, with MICs 2.0 µg/mL. Gatifloxacin and moxifloxacin were the most active quinolones tested, while the MICs of the β-lactams rose with those of penicillin G. LBM415 at two times the MIC was bactericidal (99.9% killing) against six strains after 24 h.
 IT 478913-91-6, LBM 415
 RL: BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)
 (antipneumococcal activity of LBM415, new peptide diformylase inhibitor, compared with those of other agents)
 RN 478913-91-6 HCAPLUS
 CN L-Prolineamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 25 THERE ARE 25 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


AB β-Amino acid deriva. I (R is alkyl, R1-R3 are H or alkyl or R2R3C are cycloalkyl, Y is a protecting group), intermediates in the synthesis of aminoacyl azacycloalkanes II [same R-R3 and Y, R4 is aryl or heteroaryl, n is 0-3, X is CH2, S, CHOH, CH(OR), CH(SH), CF2, C=N(OR) or CHF] were prepared by hydrogenation of corresponding α-alkylidene deriva. in the presence of a chiral ligand and a catalytic amount of a hydrogenation catalyst. Thus, a mixture of 2-[(phenylmethoxy)amino]methyl-2-hexenoic acid Me ester (apprx. 1:1 E/Z, preparation given), bis(norbornadiene)rhodium(I) tetrafluoroborate and (1S,1'S,2R,2'R)-TangPhos in deoxygenated methanol in a Parr bottle is hydrogenated under H2 (45-55 psi) at room temperature for 24 h to afford 94 % 2-[(phenylmethoxy)amino]methyl-(2S)-hexanoic acid Me in 95 % yield (R:S = 98:2).
 IT 478913-93-9
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (preparation of β-amino acid intermediates in synthesis of aminoacylpyrrolidinecarboxamides and related antibacterial compds.)
 RN 478913-93-9 HCAPLUS
 CN L-Prolineamide, (2R)-2-butyl-N-formyl-N-(phenylmethoxy)-β-alanyl-N-(5-fluoro-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



IT 478912-56-0P
 RL: SPN (Synthetic preparation); PRP (Preparation)
 (preparation of β-amino acid intermediates in synthesis of aminoacylpyrrolidinecarboxamides and related antibacterial compds.)
 RN 478912-56-0 HCAPLUS
 CN L-Prolineamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L17 ANSWER 27 OF 32 HCAPLUS COPYRIGHT 2007 ACS ON STN
 ACCESSION NUMBER: 2004:383872 HCAPLUS [Full-text](#)
 DOCUMENT NUMBER: 141:170760
 TITLE: Antimicrobial spectrum and activity of NVP PDF-713, a

L17 ANSWER 26 OF 32 HCAPLUS COPYRIGHT 2007 ACS ON STN
 ACCESSION NUMBER: 2004:740215 HCAPLUS [Full-text](#)
 DOCUMENT NUMBER: 141:261060
 TITLE: Process for preparing β-amino acid intermediates in the synthesis of aminoacylpyrrolidinecarboxamides and related antibacterial compounds
 INVENTOR(S): Prashad, Mahavir; Kim, Hang-yong; Hu, Bin; Slade, Joel; Kapa, Prasad Koteswara; Girgis, Michael John
 PATENT ASSIGNER(S): Novartis Ag, Switz.; Novartis Pharma GmbH
 SOURCE: PCT Int. Appl., 52 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004076053	A2	20040910	WO 2004-US5159	20040220
WO 2004076053	A3	20041202		
M: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NA, NI, NO, NZ, OM, PA, PE, PG, PH, PK, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, SM, SN, SV, TC, TD, TM, TR, TT, TZ, UA, UG, UZ, VC, VE, VU, WO, XG, YU, ZA, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, CA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2004216178	A1	20040910	AU 2004-216178	20040220
CA 2516465	A1	20040910	CA 2004-2516465	20040220
EP 1599440	A2	20051130	EP 2004-713381	20040220
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IS, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
BR 2004007448	A	20060131	BR 2004-7448	20040220
CN 1759097	A	20060412	CN 2004-8006326	20040220
JP 2006519786	T	20060831	JP 2006-503764	20040220
PRIORITY APPLN. INFO.:				
US 2003-449015P			P 20030221	
US 2003-449016P			P 20030221	
US 2003-449017P			P 20030221	
WO 2004-US5159			A 20040220	

OTHER SOURCE(S): CASREACT 141:261060; MARPAT 141:261060
 GI



novel peptide deformylase inhibitor, tested against 1,837 recent gram-positive clinical isolates
 Jones, Ronald N.; Fritsche, Thomas R.; Sader, Helio S.
 CORPORATE SOURCE: The JONES Group/JMI Laboratories, North Liberty, IA, USA
 SOURCE: Diagnostic Microbiology and Infectious Diseases (2004), 49(1), 63-65
 CODEN: DMIID2; ISSN: 0732-8893
 PUBLISHER: Elsevier Science Inc.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 AB Continued emergence of antimicrobial resistances among gram-pos. pathogens requires further development of compds. with novel modes of action. The peptide deformylase inhibitor NVP PDF-713 was tested against 1,837 recent strains of Gram-pos. organisms. All NVP PDF-713 MICs were at ≤4 µg/mL except for 6 enterococci (0.3% of strains overall). NVP PDF-713 MIC90 results were: Staphylococcus aureus, β-haemolytic and viridans group streptococci and Streptococcus bovis at 1 µg/mL; coagulase-neg. staphylococci, Streptococcus pneumoniae, and Listeria spp. at 2 µg/mL; and the enterococci at 4 µg/mL. NVP PDF-713 appears to be a promising new agent worthy of continued in vivo development.
 IT 478913-91-6, NVP PDF-713
 RL: BSU (Biological study, unclassified); DNA (Drug mechanism of action); THU (Therapeutic use); BIOL (Biological study); USBS (Uses)
 (antibiotic spectrum and activity of peptide deformylase inhibitor NVP PDF-713 gram-pos. pathogens)
 RN 478913-91-6 HCAPLUS
 CN L-Prolineamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L17 ANSWER 28 OF 32 HCAPLUS COPYRIGHT 2007 ACS ON STN
 ACCESSION NUMBER: 2004:361959 HCAPLUS [Full-text](#)
 DOCUMENT NUMBER: 141:136885
 TITLE: Potential utility of a peptide deformylase inhibitor (NVP PDF-713) against oxazolidinone-resistant or streptogramin-resistant Gram-positive organism isolates
 AUTHOR(S): Jones, Ronald N.; Moet, Gary J.; Sader, Helio S.; Fritsche, Thomas R.
 CORPORATE SOURCE: The JONES Group/JMI Laboratories, North Liberty, IA, 52317, USA
 SOURCE: Journal of Antimicrobial Chemotherapy (2004), 53(5), 804-807
 CODEN: JACHDX; ISSN: 0305-7453
 PUBLISHER: Oxford University Press
 DOCUMENT TYPE: Journal

```

GH, HR, HU, ID, IL, IN, IS, JP, KR, KO, KP, KR, KZ, LC, LK, LT,
LU, LV, MA, MD, MK, MN, MX, NI, NO, NZ, OM, PG, PH, PL, PT, RO,
RU, SC, SE, SG, SK, SY, TJ, TM, TN, TR, TT, UA, US, UZ, VC, VN,
YU, ZA, ZW
RM: AM, AZ, BY, KG, KZ, MD, RO, TJ, TM, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE,
SI, SK, TR
CA 2499426 A1 20040401 CA 2003-2499426 20030918
AU 2003273404 A1 20040408 AU 2003-273404 20030918
EP 1542968 A1 20050622 EP 2003-755559 20030918
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK
BR 20034592 A 20040809 BR 2003-145918 20030918
CN 1684945 A 20051019 CN 2003-822471 20030918
JP 2004503053 T 20060126 JP 2004-537129 20030918
ZA 2005001923 A 20050912 ZA 2005-1923 20050307
IN 2005CN00346 A 20070406 IN 2005-CN346 20050308
NO 2005001867 A 20050418 NO 2005-1867 20050418
US 2005261504 A1 20051124 US 2005-527628 20050525
PRIORITY APPLN. INFO.: US 2002-411920P P 20030919
US 2003-480242P P 20030520
WO 2003-EP10416 W 20030918
OTHER SOURCE(S): CASREACT 140:287275; MARPAT 140:287275


```

OTHER SOURCE(S): CASREACT 140:287275; MARPAT 140:287275
GI

CCCC(=O)N1CCC[C@H]1C(=O)N(C2=CC=CC=C2)C3=CC=CC=C3

AB	Title compds. X = [Y protective group; R1 = aryl-, heteroaryl-; R2-R5 = H; aliph-; R3R3, R4R5 = alkylene; X = CH2, S, (un)substituted CH(OH), CH(SH), CF2, C(NOH), CHF, n = 0-3] were prepared for use as intermediates to prepare certain antitumor N-formyl hydroxylamine compds. which are peptide deformylase inhibitors. Thus, HOCH2CHBuCO2H was treated with PhCH2ONH2, followed by MeSO3Cl to give MeSO3CH2CHBuCONHOC2H5, which was cyclized to the β -lactam and treated with (S)-N-(5-fluoro-2-pyridinyl)pyrrolidine- 2-carboxamide, followed by formylation to give the pyrrolidine II.
IT	478913-92-7P RL: SPW (Synthetic preparation); PREP (Preparation) (process for preparing benzyloxyminoacylpyrrolidinecarboxamides)
RN	478913-92-7 HCAPLUB
CN	L-Prolineamide, (2R)-2-butyl-N-formyl-N-(phenylmethoxy)- β -alanlyl-N-(5-fluoro-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



L17 ANSWER 31 OF 32 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2003:883789 HCAPLUS Full-text
DOCUMENT NUMBER: 141:20308
TITLE: Antibacterial susceptibility of a vancomycin-resistant
Staphylococcus aureus strain isolated at the Hershey
Medical Center
AUTHOR(S): Bozdogan, Buelent; Seel, Duygu; Whitener, Cynthia;
Browne, Frederick A.; Appelbaum, Peter C.
CORPORATE SOURCE: Department of Pathology, Hershey Medical Center,
Hershey, PA, 17033, USA
SOURCE: Journal of Antimicrobial Chemotherapy (2003), 52(5),
864-868
CODEN: JACHDX; ISSN: 0305-7453
PUBLISHER: Oxford University Press
DOCUMENT TYPE: Journal


AB Staphylococcus aureus strain HM03 isolated at the Hershey Medical Center, was resistant to vancomycin (VRSA) through the presence of the vanA resistance gene; it also contained mecA, erm(A), erm(B), tet(K) and sac(6')-aph(2''), conferring resistance to licensed β -lactams, macrolides, tetracyclines and aminoglycosides. HM03 also had alterations in GyrA and GrlB and was resistant to available quinolones. Exptl. drug MICs (\pm 2 mg/L) for HM03 included cephalosporins BA954 and cefazolin; glycopeptides telavancin and dalbavancin; the lipopeptide daptomycin; the glycolipodepsipeptide ceftazidime; fluoroquinolones MCK 771 A, MCK 1153, DK-507K and sitafloxacin; and the DNA topoisomerase II inhibitors WVP-PDF713 and GS02-12. These agents were all bactericidal as were trimethoprim/sulfamethoxazole and teicoplanin (MIC 4 mg/L). Oxazolidinones linezolid and rambexidol; the injectable streptogramin quinupristin/dalfopristin; DNA topoisomerases GS2-10547 and GS02-104; peptide deformylase inhibitors WVP-PDF713 and GS02-12; tetracycline derivative tigecycline; the antiolate iclaprim; mupirocin and fusidic acid were all active in vitro but bacteriostatic.

IT 478913-91-6
RL: BSU (Biological study, unclassified); PAC (Pharmacological activity);
THU (Therapeutic use); BIOL (Biological study); USRS (Uses)
(antibacterial susceptibility of a vancomycin-resistant *Staphylococcus aureus* strain isolated at the Hershey Medical Center)

RN 478913-91-6 NCAPLUS

CN L-Prolinamide, (2R)-2-butyl-L-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-oxydo-2-nwridjnyll, [9C]l, (#A.WNDXPX NAWM).



Absolute stereochemistry



L17 ANSWER 32 OF 32 HCAPLUS COPYRIGHT 2007 ACS on STN
ACCESSION NUMBER: 2002:977804 HCAPLUS Full-text

DOCUMENT NUMBER: 138:55863
 TITLE: Preparation of N-formyl-N-hydroxylamino-substituted pyrrolidine derivatives as inhibitors of peptidyl deformylase
 INVENTOR(S): Patel, Dinesh V.; Yuen, Zhengyu; Jain, Rakesh K.; Garcia Alvarez, Salvador; Jacobs, Jeffrey
 PATENT ASSIGNEE(S): Versicor, Inc., USA; Novartis AG
 SOURCE: PCT Int. Appl., 69 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002102790	A1	20021227	WO 2002-EP6604	20020614
M: AB, AG, AL, AM, AT, AU, AZ, BA, BB, BO, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LT, LU, LV, MA, MD, ME, MN, MO, NP, OM, PH, PL, PT, RO, RU, SE, SG, SI, SK, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ZA, ZW				
RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR				
CA 2448526	A1	20021227	CA 2002-2448526	20020614
AU 2002312062	A1	20030102	AU 2002-312062	20020614
US 2003045479	A1	20030306	US 2002-171706	20020614
US 7148242	B2	20061212		
EP 1401828	A1	20040331	EP 2002-754681	20020614
EP 1401828	B1	20060412		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
HU 200400208	A2	20040628	HU 2004-208	20020614
CN 1511152	A	20040707	CN 2002-810596	20020614
BR 2002010377	A	20040810	BR 2002-10377	20020614
JP 2005502606	T	20050127	JP 2003-506263	20020614
NZ 529489	A	20051028	NZ 2002-529489	20020614
AT 323081	T	20060415	AT 2002-754681	20020614
PT 1401828	T	20060831	PT 2002-754681	20020614
ES 2262824	T3	20061201	ES 2002-2754681	20020614
ZA 2003008379	A	20040521	ZA 2003-8379	20031028
IN 2003CN01963	A	20060106	IN 2003-CN1963	20031210
NO 2003005871	A	20040216	NO 2003-5871	20031212
HK 1064370	A1	20061020	HK 2004-107013	20040914
PRIORITY APPL. INFO.:				
OTHER SOURCE(S):	MARPAT 138:55863			
GI				




carboxylic acid N-(4,6-dimethylpyridin-2-yl)amide 478912-97-9P,
 (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(4-ethylpyridin-2-yl)amide 478913-05-2P,
 (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(3-hydroxypyridin-2-yl)amide 478913-12-1P,
 (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(isoquinolin-1-yl)amide 478913-16-5P,
 (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(quinolin-3-yl)amide 478913-21-2P,
 (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]azetidine-2-carboxylic acid N-(4-methylpyridin-2-yl)amide 478913-24-5P,
 (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]azetidine-2-carboxylic acid N-(5-methylpyridin-2-yl)amide 478913-27-8P,
 (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]azetidine-2-carboxylic acid N-(5-fluoropyridin-2-yl)amide 478913-30-3P,
 (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(5-trifluoromethyl-1-oxopyridin-2-yl)amide
 478913-37-0P, (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(4-phenylpyridin-2-yl)amide
 478913-41-6P, (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(4-phenyl-1-oxopyridin-2-yl)amide
 478913-45-0P, (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(4-trifluoromethylpyridin-2-yl)amide
 478913-46-3P, (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(4-trifluoromethyl-1-oxopyridin-2-yl)amide
 478913-51-8P, (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(8-hydroxyquinolin-2-yl)amide
 478913-55-2P, (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(3-methoxy-6-methylpyridin-2-yl)amide
 478913-59-6P, (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(4-methoxypyridin-2-yl)amide
 478913-64-3P, (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(3-methoxypyridin-2-yl)amide
 478913-68-7P, (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(3-methoxypyridin-2-yl)amide
 478913-69-6P, (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(6-hydroxypyridin-2-yl)amide
 478913-75-6P, (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(3-hydroxy-1-oxopyridin-2-yl)amide
 478913-83-6P, (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(2,2-difluorobenzo[1,3]dioxol-5-yl)amide
 478913-87-0P, (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(3-phenoxyphenyl)amide
 478913-91-6P, (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(5-fluoro-1-oxopyridin-2-yl)amide
 478913-94-9P 478913-96-1P, (2S,4R)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(4-hydroxypyrrrolidine-2-carboxylic acid N-(5-methylpyridine-2-yl)amide 478913-99-4P,
 (2S,4R)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]-4-hydroxypyrrrolidine-2-carboxylic acid N-(5-methylpyridine-2-yl)amide
 478914-01-1P, (2S,4R)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]-4-hydroxypyrrrolidine-2-carboxylic acid N-(4-ethylpyridine-2-yl)amide
 478914-03-3P, (2S,4R)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]-4-hydroxypyrrrolidine-2-carboxylic acid N-(5-trifluoromethylpyridine-2-yl)amide 478914-05-5P, (2S,4R)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]-4-hydroxypyrrrolidine-2-carboxylic acid N-(5-methylpyridine-2-yl)amide 478914-08-8P,
 (2S,4R)-4-fluoro-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(5-methylpyridine-2-yl)amide 478914-10-2P
 478914-12-4P, (2S,4R)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]-4-methoxypyrrrolidine-2-carboxylic acid N-(5-methylpyridine-2-yl)amide
 478914-17-9P, (2S,4R)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]-4-methoxypyrrrolidine-2-carboxylic acid N-(5-methylpyridine-2-yl)amide

AB Title compds. I [X = CH₂, S, CHOH, CH-alkoxy, CHSH, etc.; R1 = (hetero)aryl; R2-5 = H, alkyl, etc.; n = 0-3 provided that when n = 0, X = CH₂] are prepared For instance, (S)-2-(chlorocarbonyl)pyrrolidine-1-carboxylic acid benzyl ester is used to acylate 2-aminopyridine and the resulting amide deprotected and coupled to (2R)-2-[(benzyloxyformylamino)methyl]hexanoic acid (preparation given; dioxane, HATU, i-Pr₂NEt) to give II. IC50 of selected examples of I against MMP-7 ranges from >10 pM to >100 pM, whereas the IC50 of these same compds. against zinc-containing peptidyl deformylase (PDF) ranges from about 0.005 pM to 5 pM, and against nickel-containing PDF ranges from about 0.001 pM to about 0.3 pM. I are useful for preventing contamination of a cell culture medium.

IT 478913-80-3P
 RL: PAC (Pharmacological activity); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USSS (Uses)
 (preparation of N-formyl-N-hydroxylamino-substituted pyrrolidine derivs. as inhibitors of peptidyl deformylase)
 RN 478913-80-3 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-1,3-benzodioxol-5-yl- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 478912-45-7P, (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(pyridin-2-yl)amide 478912-46-0P,
 (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(3-methylpyridin-2-yl)amide 478912-50-4P,
 (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(6-methylpyridin-2-yl)amide 478912-52-6P,
 (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(4-methylpyridin-2-yl)amide 478912-56-0P,
 (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(5-fluoropyridin-2-yl)amide 478912-59-3P,
 (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(5-methylpyridin-2-yl)amide 478912-63-9P,
 (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(6-ethylpyridin-2-yl)amide 478912-66-2P,
 (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(5-trifluoromethylpyridin-2-yl)amide
 478912-69-5P, (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(6-fluoropyridin-2-yl)amide
 478912-76-4P, (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(4,6-dimethyl-1-oxopyridin-2-yl)amide
 478912-80-0P, (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(4-methyl-1-oxopyridin-2-yl)amide
 478912-85-5P, (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]azetidine-2-carboxylic acid N-(pyridin-2-yl)amide 478912-92-4P,
 (2S)-1-[(2R)-2-[(Formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-

479067-88-4P
 RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USSS (Uses)
 (preparation of N-formyl-N-hydroxylamino-substituted pyrrolidine derivs. as inhibitors of peptidyl deformylase)


RN 478912-45-7 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-2-pyridinyl- (CA INDEX NAME)

Absolute stereochemistry.




RN 478912-48-0 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(3-methyl-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 478912-50-4 HCAPLUS
 CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(6-methyl-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478912-52-6 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy- β -alanyl-N-(4-methyl-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478912-56-0 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy- β -alanyl-N-(5-fluoro-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478912-59-3 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy- β -alanyl-N-(5-methyl-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478912-63-9 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy- β -alanyl-N-(6-ethyl-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478912-80-0 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy- β -alanyl-N-(4-methyl-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478912-85-5 HCAPLUS

CN 2-Azetidinecarboxamide, 1-[(2R)-2-[(formylhydroxyamino)methyl]-1-oxohexyl]-N-2-pyridinyl-, (2S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478912-92-4 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy- β -alanyl-N-(4,6-dimethyl-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478912-66-2 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy- β -alanyl-N-[5-(trifluoromethyl)-2-pyridinyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478912-69-5 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy- β -alanyl-N-(6-fluoro-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478912-76-4 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy- β -alanyl-N-(4,6-dimethyl-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478912-97-9 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy- β -alanyl-N-(4-ethyl-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-05-2 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy- β -alanyl-N-(3-hydroxy-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-12-1 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy- β -alanyl-N-1-isoquinolinyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-16-5 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy- β -alanyl-N-3-quinolinyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-21-2 HCAPLUS

CN 2-Azetidinecarboxamide, 1-[(2R)-2-[(formylhydroxyamino)methyl]-1-oxohexyl]-N-(4-methyl-2-pyridinyl)-, (2S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 478913-24-5 HCAPLUS


CN 2-Azetidinecarboxamide, 1-[(2R)-2-[(formylhydroxyamino)methyl]-1-oxohexyl]-N-(5-methyl-2-pyridinyl)-, (2S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

RN 478913-41-6 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy- β -alanyl-N-[1-oxido-4-phenyl-2-pyridinyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-45-0 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy- β -alanyl-N-[4-(trifluoromethyl)-2-pyridinyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.




RN 478913-46-3 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy- β -alanyl-N-[1-oxido-4-(trifluoromethyl)-2-pyridinyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 478913-51-8 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy- β -alanyl-N-(8-hydroxy-2-quinolinyl)- (9CI) (CA INDEX NAME)

RN 478913-27-8 HCAPLUS

CN 2-Azetidinecarboxamide, N-(5-fluoro-2-pyridinyl)-1-[(2R)-2-[(formylhydroxyamino)methyl]-1-oxohexyl]-, (2S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-30-3 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy- β -alanyl-N-[1-oxido-5-(trifluoromethyl)-2-pyridinyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-37-0 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy- β -alanyl-N-(4-phenyl-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-55-2 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy- β -alanyl-N-(3-methoxy-6-methyl-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-59-6 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy- β -alanyl-N-(4-methoxy-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-64-3 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy- β -alanyl-N-(3-methoxy-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-68-7 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-(phenylmethoxy)-β-alanyl-N-(3-methoxy-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-69-8 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(1,6-dihydro-6-oxo-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-75-6 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-methoxycarbonyl-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-83-6 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(2,2-difluoro-1,3-benzodioxol-5-yl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-87-0 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(4-phenoxyphenyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-91-6 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-fluoro-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-94-9 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(5-methoxycarbonyl-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-96-1 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-4-hydroxy-N-(5-methyl-2-pyridinyl)-, (4R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-99-4 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-4-hydroxy-N-(5-methyl-2-pyridinyl)-, (4S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478914-01-1 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(4-ethyl-2-pyridinyl)-4-hydroxy-, (4R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478914-03-3 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-4-hydroxy-N-[5-(trifluoromethyl)-2-pyridinyl]-, (4R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478914-05-5 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-4-fluoro-N-(5-methyl-2-pyridinyl)-, (4S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478914-08-8 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-4-fluoro-N-(5-methyl-2-pyridinyl)-, (4R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478914-10-2 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-4,4-difluoro-N-(5-methyl-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478914-12-4 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-4-methoxy-N-(5-methyl-2-pyridinyl)-, (4R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478914-17-9 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-4-methoxy-N-(5-methyl-2-pyridinyl)-, (4S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 479067-88-4 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-hydroxy-β-alanyl-N-(4-ethyl-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




IT 478913-11-0P, (2S)-1-[(2R)-2-[(Benzoyloxyformylamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(3-benzoyloxy-2-pyridin-2-yl)amide
 478913-15-4P, (2S)-1-[(2R)-2-[(Benzoyloxyformylamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(4-methoxy-2-pyridin-2-yl)amide
 478913-20-1P, (2S)-1-[(2R)-2-[(Benzoyloxyformylamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(4-methoxy-2-pyridin-2-yl)amide
 478913-20-1P, (2S)-1-[(2R)-2-[(Benzoyloxyformylamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(4-methoxy-2-pyridin-2-yl)amide
 478913-63-2P, (2S)-1-[(2R)-2-[(Benzoyloxyformylamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(4-methoxy-2-pyridin-2-yl)amide
 478913-74-5P, (2S)-1-[(2R)-2-[(Benzoyloxyformylamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(4-methoxy-2-pyridin-2-yl)amide

1)pyrrolidine-2-carboxylic acid N-(6-benzoyloxy-2-pyridin-2-yl)amide
 478913-79-0P, (2S)-1-[(2R)-2-[(Benzoyloxyformylamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(3-benzoyloxy-2-pyridin-2-yl)amide
 478913-92-7P, (2S)-1-[(2R)-2-[(Benzoyloxyformylamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(5-fluoro-2-pyridin-2-yl)amide
 478913-93-8P, (2S)-1-[(2R)-2-[(Benzoyloxyformylamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(5-fluoro-1-oxo-2-pyridin-2-yl)amide
 478913-98-3P, (2S,4R)-4-(Benzoyloxy)-1-[(2R)-2-[(formylhydroxyamino)methyl]hexanoyl]pyrrolidine-2-carboxylic acid N-(5-methyl-2-pyridin-2-yl)amide 478914-16-8P.
 (2S,4R)-1-[(2R)-2-[(Benzoyloxyformylamino)methyl]hexanoyl]-4-methoxy-2-pyridin-2-carboxylic acid N-(5-methyl-2-pyridin-2-yl)amide
 478914-21-5P, (2S,4S)-1-[(2R)-2-[(Benzoyloxyformylamino)methyl]hexanoyl]-4-methoxy-2-pyridin-2-carboxylic acid N-(5-methyl-2-pyridin-2-yl)amide
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (preparation of N-formyl-N-hydroxyamino-substituted pyrrolidine derivs. as inhibitors of peptidyl deformylase)

RN 478913-11-0 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-(phenylmethoxy)-β-alanyl-N-[3-(phenylmethoxy)-2-pyridinyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-15-4 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-(phenylmethoxy)-β-alanyl-N-1-isoquinolinyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-20-1 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-(phenylmethoxy)-β-alanyl-N-1-quinolinyl- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-63-2 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-(phenylmethoxy)-β-alanyl-N-(4-methoxy-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-74-5 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-(phenylmethoxy)-β-alanyl-N-[6-(phenylmethoxy)-2-pyridinyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-79-0 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-(phenylmethoxy)-β-alanyl-N-[1-oxido-3-(phenylmethoxy)-2-pyridinyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-92-7 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-(phenylmethoxy)-β-alanyl-N-(5-fluoro-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 478913-93-8 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-(phenylmethoxy)-β-alanyl-N-(5-fluoro-1-oxido-2-pyridinyl)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478913-98-3 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-(phenylmethoxy)-β-alanyl-N-(5-methyl-2-pyridinyl)-4-(phenylmethoxy)-, (4R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478914-16-8 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-(phenylmethoxy)-β-alanyl-4-methoxy-N-(5-methyl-2-pyridinyl)-, (4R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 478914-21-5 HCAPLUS

CN L-Prolinamide, (2R)-2-butyl-N-formyl-N-(phenylmethoxy)-β-alanyl-4-methoxy-N-(5-methyl-2-pyridinyl)-, (4S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> log hold
COST IN U.S. DOLLARS
FULL ESTIMATED COST

SINCE FILE	TOTAL
ENTRY	SESSION
173.84	760.69

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
CA SUBSCRIBER PRICE	ENTRY	SESSION
	-24.96	-41.02

SESSION WILL BE HELD FOR 120 MINUTES
STN INTERNATIONAL SESSION SUSPENDED AT 09:07:23 ON 30 MAY 2007